

**The influence of He-Ne Laser (632.8 nm) on *Candida albicans* isolated from complete upper dentures: In vitro study**

***Dhilal Mahdi Al-Muathen\****

Date of acceptance 3/3 / 2009

**Abstract:**

The main objective of this study is to determine whether the use of He-Ne Laser (632.8 nm wavelength and power 0.5 mW) is an eligible and effective tool to kill or reduce the cell viability of *Candida albicans* isolated from complete upper dentures.

Twenty one swabs were taken from the complete upper dentures. Only six swabs showed positive cultures for *C. albicans*.

The isolate was divided into two groups, group I was not irradiated (control), and group II was irradiated by He-Ne Laser for different periods (10, 15, 20, and 30) min.

After irradiation, the results showed a significant reduction in the viable cell count and colonies diameters especially at exposure periods 10 and 15 min.

Although the low power He-Ne Laser was not eradicating the cell itself, but it affected on the viable cell count and colonies diameters comparing with the control.

**Key words:** He-Ne Laser, *Candida albicans*, in vitro.

**Introduction:**

*Candida albicans* is a common inhabitant of the human oral cavity [1]. Several reports indicated that this organism can be considered as a part of normal human flora [2] and it is an opportunistic yeast [3].

Presence of denture is one of the factors that could enhance the growth of the organism in the oral cavity of human [4]. Mccourtie and Douglas [5] had been shown that *C. albicans* adheres to acrylic surfaces.

Dentures may change the flora of the oral cavity as a result of food debris and plaque that collects between the mucosal surface of the denture and the palate. In addition to that the saliva which is present between the maxillary denture and the mucosa may have a pH

lower than usual [6]. Atrophic (Erythematous) candidiasis is most commonly found in patients with ill-fitting dentures or in those who wear their dentures continuously [7].

The He-Ne Laser is electrically pumped gas Laser. Low current (in milliamperes) discharged between the cathode and the anode of the tube of high voltage difference (hundreds of volts) applied between them [8].

Laser can enhance the physiological changes without structural changes, so Low Energy Laser Irradiation (LELI) was developed for using in the medical biology [9].

Frucht-Pery [10] used ArF excimer Laser for *C. albicans* and

\*Al-Mustansiriya University-College of Dentistry-Basic Science Department-Biology

explained that this type of laser may be reduce the yeast in the laboratory significantly. And with some bacteriocidals Tolebzadeh et. al. [11] used CO<sub>2</sub> Laser to treat the limited bacterial infections and they found that the survival of cells decrease with the increase of Laser intensity. Ward et. al. [12] demonstrated that the use of Nd-YAG Laser was able to kill some G<sup>+</sup> and G<sup>-</sup> bacteria and 2 types of yeast (*C. albicans*, and *Saccharomyces cerevisiae*). Al-Masraf [13] did a wide study about the effect of He-Ne Laser on *pseudomonas aeruginosa* and she was found that there is a significant differences in the responsiveness of these germs to this type of Laser, thus she also indicated that these germs were resistant to this ray at density of low power while their resistance reduced with the increase of it's power density. Rassam [14] found that He-Ne Laser was the best performing method for elimination of *Staphylococcus aureus* with using (TBO) as photosensitizer.

### Materials and Methods:

#### \* Collection of Samples

Twenty one swabbed samples were taken. Subjects were eligible if they were edentulous and had a complete acrylic upper denture. Subjects with partial dentures or lower dentures only and those who had used a denture cleanser or antibiotics within the previous two weeks not eligible [15].

All of the tissue bearing surface of the complete maxillary dentures was swabbed for 30 seconds with a dry sterile swab to harvest plaque. Each portion of the surface-bearing side was swabbed three times [16].

All of the swabs were soaked in physiological saline in sterile test tubes.

#### \* Laboratory Identification

The swabs were transported to the laboratory to be inoculated on Sabourauds' dextrose agar plates containing penicillin, chloramphenicol, and streptomycin to prevent bacterial growth. The specimens were incubated at 30°C for 4 days [17].

The isolated colonies were identified by the following methods: -

##### 1. Gram-stained film: -

In which *C. albicans* appears as gram positive rounded or oval yeast cells when examined under the light microscope.

##### 2. The morphological appearance of colonies: -

Colonies on Sabourauds' media appear as high convex, soft, cream-colored or off-white, glabrous or membranous and may have a fringe of submerged hyphae with yeast-like odor. The colony became (0.5 mm) in diameter after (18 hr.) and developed into (1.5-2 mm) in (5-7 days) on this medium [18]. Colonies diameters were measured for irradiated and control samples by using of ocular micrometer, after putting the opened dishes containing the colonies on the stage of the microscope.

##### 3. Germ tube formation: -

It is one of the most valuable test used in most laboratories for rapid and presumptive identification of *C. albicans*.

This test was performed by inoculating a small portion of the organism's colony (0.5) ml of human serum contained in sterile tubes and incubated at 37°C for 2-3 hours. Then a drop of the sediment containing the organism in the bottom of the tube is placed on a

slide to be examined microscopically for the presence of germ tubes which sometime described as having (hand mirror) shape [19].

The presence of *C. albicans* was considered when the three tests were positive.

#### \* Laser System

The Laser was used in this study CW Helium Neon gas Laser (Hamburg-Germany) with a measured output power of (0.5 mW), which emits light in a collimated beam with diameter (4 mm), wavelength (632.8 nm), and power density (3.98 mW/cm<sup>2</sup>), which was calculated from this equation [20, 21]: -

Power Density (mW/cm<sup>2</sup>) = output power (mW) / area (cm<sup>2</sup>).

#### \* Samples Preparation and Irradiation Procedure

The yeast was cultured on (9ml) of Sabourauds' dextrose broth in screw cup bottle and incubated at 30°C for 4 days. After that a series of dilutions were made and then (2ml) of each dilution were added to each of eight sterilized test tubes each of them containing (2ml) of (9%) NaCl. Four of these test tubes were irradiated with He-Ne Laser for different exposure time (10, 15, 20, and 30) min, and the latter four tubes were considered as control.

#### \* Microbiological Procedures after Irradiation

For each test tubes (the irradiated test tubes and the control) (0.5ml) was taken and added to sterilized Petri dishes containing SDA. Then, CFUs and colonies diameters were determined after incubation at 30°C for 4 days.

#### \* Statistical Analysis

Statistical analysis was done by two-ways analysis of variance (ANOVA2) to determine the effect of He-Ne Laser on CFUs and colonies diameters of *C. albicans* in different exposure times.


#### Results:

Laboratory identification revealed that out of twenty one subjects six subjects showed positive cultures only. The prevalence of *C. albicans* isolated from complete upper dentures was found to be (28.6%).

The samples were irradiated by He-Ne Laser for different time periods (10, 15, 20, and 30) min, because of the longest irradiation periods do not appear results better than the shortest periods.

The cell viability is considered the important point for studying the effect of low power of He-Ne Laser on *C. albicans* yeast. Comparing with untreated (control), (CFUs) of irradiated samples peaked at (10 min) irradiation period, and then, declined at (15 min), thereafter when the irradiation period was increased more than that we saw that there was a slightly paralleled increase in (CFUs) of the sample, whereas the control samples (CFUs) increased gradually during the time and peaked at (30 min) of the experiment's period.


(fig, 1) demonstrates the decrease in (CFUs) especially in the periods of irradiation (15-30) min, and there is a significant relationship between (CFUs) of irradiated samples and irradiation periods in count ( $P < 0.05$ ,  $0.01$ ).



**Fig. 1: The effect of He-Ne Laser radiation on CFUs for different irradiation periods**

On the other hand, in spite of the irradiation by He-Ne Laser did not kill the cells completely, the survival cells were affected clearly by measuring the colonies diameters of irradiated samples after culturing on solid media and compare with the diameters of non irradiated samples.

(Fig. 2) demonstrates the decrease in colonies diameters especially in shortest period of irradiation (10) min, and there is a significant relationship between colonies diameters of irradiated samples and irradiation periods in count (P<0.05, 0.01).



**Fig. 2: The effect of He-Ne Laser radiation on colonies diameter for different irradiation periods.**

**Discussion: -**

Our previous results indicated that the prevalence of *C. albicans* isolated from complete upper dentures was (28.6%) and this result comes in disagreement with the finding of Tmilade et. al. [22] who recorded out of 8 patients 5 patients (62.5%) showed positive result to *C. albicans* isolated from removable upper dentures. This disparity may be attributed to differences in the sampling techniques, in the sites of sampling, and in the selection of the subjects (4).

As it is illustrated in (fig. 1), there is a significant decrease (P<0.05, 0.01) in (CFUs) especially for the periods of irradiation (15-30) min.

This result shows the effect of He-Ne Laser radiation on the viability of cells which is may be affected on the cell division and do not kill the cell completely, and this is similar to the results of Wilson and Mia [23] and Bown and Lovate [24] on yeast cell. Fedoseyeva et. al. [25] and Karu and Letokhov [26] demonstrated that the irradiation by He-Ne Laser may be cause a decrease in the time of yeast formation. Karu et. al. [27] showed that the irradiation of Hela cells by He-Ne Laser with doses (10-10<sup>3</sup>) Joule/cm<sup>2</sup> will stimulate or increase DNA of the cells in S-phase, which also explain the stimulation of cells in long irradiation periods.

When the irradiation period was increased more than that, we saw that there was an increase in viable cell counts, this result may be pointed to the ability of cells to repair the damage that happened by the irradiation with He-Ne Laser in shortest periods by developing a special mechanism to ignore the damage or repair it rapidly, and this may be due to the absorption of cells to the high energy of heat in

the long irradiation periods that enable the cells to restore their viability, so for these reasons the shortest periods were chosen.

As it is shown in (fig. 2), the colonies diameters are well correlated with the irradiation periods, and there is a significant decreasing ( $P < 0.05$ ,  $0.01$ ) in the colonies diameters of irradiated samples comparing with the control, the same result appeared in the study of Quickenden and Daniels [28] about the effect of the same radiation on *Saccharomyces cerevisiae* yeast.

The reduction of colonies diameters that resulted from irradiated cells may be caused by the effect of He-Ne Laser on growth and proliferation rates of the yeast [29].

#### References:

1. Samaranayake, L.P. 1992. Oral mycoses in HIV infection. *Oral Surg. Oral Med. Oral Pathol.*, 73: 171-180.
2. Taschdjian, C.L., Seeling, M.S. and Kozinn, P.J. 1973. Serological diagnosis of candidal infections. *CBC. Grit. Rev. Clin. Lab. Sci.*, 4: 19-59.
3. Witkin, S.S. 1991. Immunological factors influencing susceptibility to recurrent candidal vaginitis. *Clin. Obstet. Gynecol.*, 34: 662-668.
4. Arendorf, F.M. and Walker, D.M. 1980. The prevalence and intra-oral distribution of *Candida albicans* in man. *Arch. Oral Biol.*, 25: 1-10.
5. McCourtie, J. and Douglas, L.J. 1981. Relationship between cell surface composition of *Candida albicans* and adherence to acrylic after growth on different carbon sources. *Infection and Immunity*. 32: 1234-1241.
6. Dorey, L., Janet. Bruce Blasberg, Mac Entee, I. and Michael. Conklin. J. Robert 1985. Oral mucosal disorders in denture wearers. *J. of Prosthetic Dentistry*, 53: 210-213.
7. Philip, Sap J., Eversole. Lewis R. and Wysocki. George P. 1997. Contemporary oral and maxillofacial pathology. Mosby, 2<sup>nd</sup> edi., China, pp. 241.
8. Myers, ID. 1991. Laser in dentistry. *J. Am. Dent. Assoc.*, Vo. 25, pp. 113.
9. Guha, S.K. 1992. Changes in tissue on laser irradiation and implication in clinical practice. In: Chopra and Chawla (Ed.). *Laser chemical and biological sciences*. Wiely eastern limited, New Delhi, 147-159.
10. Frucht-Perry, J. Mor. M., Evorn. R., Lewis. A. and Zauberman. H. 1993. The effect of the ArF excimer laser on *Candida albicans* in vitro. *Graefes Arch. Clin. Exp. Ophtalmd.*, 231 (7): 413-415.
11. Tolebzadeh, N., Morrison. P.R. and Fried. M.P. 1994. Comparative cell targeting in vitro using the CO<sub>2</sub> laser. *Laser in surgery and medicine*, 14: 164-167.
12. Ward, G.D., Watson. I.A., Stewart. T.D., Wardlaw. A.C. and Chatwin. C.R. 1996. Inactivation of bacteria and yeast on agar surfaces with power Nd:YAG laser light. *Lett. Appl. Microbiol.*, 23 (2): 136-140.
13. Al-Masraf, T. R. 1998. Low power laser (He-Ne) interaction with bacterial cells *Pseudomonas aeruginosa*. M. Sc. Thesis, (In Arabic), College of Education, Univ. Baghdad, pp.77.
14. Rassam, Y. Z. 2005. The influence of 632.8 nm Helium Neon laser on methicillin-resistant *Staphylococcus aureus* (MIRSA) using TBO as photosensitizer. M. Sc. Thesis, Institute of Laser, Univ. Baghdad, pp.80.
15. Gornitsky, M., Paradis. I., Landaverde. G., Marie. Malo A. and Miriam. Velly A. 2002. A

- clinical and microbiological evaluation of denture cleansers for Geriatric patients in long-term care institutions. J. of the Canadian dental association., Vol. 68 (1): 39-45
16. Dills, S.S., Olshan. A.M., Goldner. S. and Brogdon. C. 1988. Comparison of the antimicrobial capability of an abrasive paste and chemical-soak denture cleansers. J. of Prosthetic dentistry, 60 (4): 467-470.
  17. Emmons, C.W., Binford. C.H. and Utz. J.P. 1970. Candidiasis, In: Medical Mycology. Lea and Febiger (ed.), 2<sup>nd</sup> ed. Philadelphia, ch.4, P.167-182.
  18. Silva-Huter, M. and Cooper. B.H. 1974. Medically important yeasts. In: Manual of clinical microbiology. 2<sup>nd</sup> ed., Lennette E.H.; Spaulding E.H., and Truant J.P. (ed.). American Society for Microbiology, Washington D.C., ch. 56, pp. 491-508.
  19. Taschdjian, C.L., Burchall. J.J. and Kozinn. P.J. 1960. Rapid identification of *Candida albicans* by filamentation on serum substitutes. Am. Med. Ass. J. Dis. Child, 99: 212-215.
  20. Al-Aamirry, L. M. 2003. Effect of He-Ne laser (632.8nm) and diode Laser (890nm) on *Streptococcus mutans* isolated from patients of carious teeth. M. Sc. Thesis, Institute of Laser, Univ. Baghdad.
  21. Al-Robaiy, K. N. 2004. Influence of (337nm) and (532 nm) Laser light on the survival of some *Bacillus* species. M. Sc. Thesis, Institute of Laser, Univ. Baghdad.
  22. Tmilade, E., Budiz-Jorgensen. E. and Tmilade. J. 1983. Predominant cultivable microflora of plaque on removable dentures in patients with healthy oral mucosa. Arch. Oral Biol., 28 (8): 675-680.
  23. Wilson, M. and Mia. N. 1993. Sensitization of *Candida albicans* to killing by low power light. J. Oral Pathology Med., 22 (8): 354-357.
  24. Bown, S.G. and Lovate. L.B. 2000. The biology of photodynamic therapy in the gastrointestinal tract. Gastrointestinal Endoscopy Clinics of North America, 10 (3): 533-550.
  25. Fedoseyeva, G.E., Karu. T.I., Letokhov. V.S., Lobko. V.V., Pomoshikova, N.A., Lyopunova. T.S. and Meissd. M.N. 1984. Effect of He-Ne laser irradiation on the reproduction and protein synthesis in yeast. Laser Chemistry, 5: 27-33.
  26. Karu, T.I. and Letokhov. V.S. 1985. Biological action of low intensity monochromatic light in the visible rang. In: Martellucci S., Chester A.N. (Ed.). Laser photobiology and photomedicine. New York, NY: plenumpress, 57-66.
  27. Karu, T.I., Pyatibra. L.V. and Kalendo. G.S. 1978. Biostimulation of Hella cells by low-intensity visible light. Stimulation of cell proliferation in vitro by He-Ne laser irradiation. II Nuovo Cimento D., 9:1485-1494.
  28. Quickenden, T. I. and Daniels. L. L. 1993. Attempted biostimulation of division in *Saccharomyces cerevisiae* using red coherent light. Photochem. Photobiol., 57 (2): 272-278.
  29. Hawar, S.N. 2002. The effect of low power laser (He-Ne) on yeast cell viability of *Candida albicans* isolated from pathogenic cases. M. Sc. Thesis (In Arabic), College of Education, Univ. Baghdad, pp. 93.

## تأثير ليزر الهليوم نيون بالطول الموجي (632.8) على خميرة المبيضات المعزولة من طقم الأسنان العلوي الكامل: دراسة مختبرية في الزجاج

ظلال مهدي المؤذن\*

\*الجامعة المستنصرية-كلية طب الأسنان-فرع العلوم الأساسية-أحياء

كلمات مفتاحية: - ليزر الهليوم-نيون, خميرة المبيضات, دراسة مختبرية في الزجاج.

### الخلاصة: -

هدف هذه الدراسة هو تحديد ما إذا كان استخدام الهليوم نيون ليزر ذو الطول موجي (632.8 نانوميتر) وقدره (0.5 ملي واط) مؤهل و ذو تأثير على قتل او تقليل عدد خلايا خميرة المبيضات المعزولة من طقم الأسنان العلوي الكامل.

تم اخذ (21) مسحة من أشخاص يرتدون طقم أسنان علوي كامل لكن فقط (6) مسحات اظهرت نتيجة زرع موجبة لخميرة المبيضات.

قسمت العزلات إلى مجموعتين، المجموعة الأولى لم يتم تشعيها (سيطرة)، والمجموعة الثانية شععت لفترات زمنية مختلفة (10، 15، 20، و 30 دقيقة).

بعد التشعيع أظهرت النتائج أن هناك نقصان ملحوظ في عدد الخلايا الحية و قطر المستعمرات خاصة عندما كانت فترة التشعيع 10 و 15 دقيقة.

نستنتج من ذلك، على الرغم من أن الهليوم نيون ليزر لم يقتل الخلية نفسها، لكنه أثر على عدد الخلايا الحية و قطر المستعمرات مقارنة مع عزلات السيطرة.