

Topological and Hyper Topological Coindices of Benzenoid Circumcoronene Series

Walid Yousef¹  , Abdu Alameri^{*2}  , Mohammed Alsharafi²  , Noman Al-Naggar² 
, Belal Al-Fuhaid³  

¹Department of Information Technology, Faculty of Computing and IT, University of Science and Technology, Sana'a, Yemen.

²Department of Biomedical Engineering, Faculty of Engineering, University of Science and Technology, Sana'a, Yemen.

³Department of Computer Science, Faculty of Computing and IT, University of Science and Technology, Sana'a, Yemen.

*Corresponding Author.

Received 03/01/2024, Revised 19/02/2024, Accepted 21/02/2024, Published Online First 20/09/2024



© 2022 The Author(s). Published by College of Science for Women, University of Baghdad.

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Numerous applications in chemistry are enabled by chemical graph theory, which is a branch of graph theory. Numerical quantities derived from the chemical graphs of a molecule, known as topological indices and co-indices, are used to model the chemical and physical properties of molecules in quantitative Structure-Property relationships (QSPR) and quantitative structure-activity relationships (QSAR) research. Fortunately, chemical-based experiments have found a strong connection between Topological descriptors (topological indices and co-indices) of molecular structures and their Physicochemical Properties, such as boiling point, and toxicity of drugs. Although several research reports have contributed to the computation of topological indices of the benzenoid circumcoronene series, studies on the calculation of topological co-indices are limited. This paper focuses on some topological co-indices. Several formulas of topological co-indices such as first Zagreb, second Zagreb, forgotten, and Yemen co-indices have been derived for the benzenoid circumcoronene series. In addition, the paper introduced new topological indices and their co-indices such as Gaza, Quds, and Palestine indices and co-indices and their mathematical formulas of the benzenoid circumcoronene series. Moreover, some algorithms have been built using Python programs to implement the mathematical formulas that are generally derived.

Keywords: Benzenoid Circumcoronene Series, Chemical Graph Theory, Molecular Graphs, Topological Co-indices, Topological Indices.

Introduction

Chemical graph theory is a branch of mathematical chemistry that applies graph theory to the study of chemical structure and properties. It is the process of visualizing molecules as graphs, with atoms serving as nodes(vertices) and atom-to-atom bonds as lines(edges). It is possible to describe the geometric structure of chemical compounds and estimate their

physical characteristics, such as melting and boiling point temperatures, by using a variety of topological measures that are calculated using graph theory¹. Chemical graph theory applies mathematical techniques from the field of graph theory and combinatorics to address chemical issues. It is related to several fields of graph theory^{2,3}. It has become an

integrative discipline that has a direct effect on maths and chemistry. It can be especially helpful in fields like structure description and isomer enumeration⁴. Discrete mathematics is used in chemical graph theory to represent the physical and biological characteristics of chemical substances. Compound features like melting and boiling points may be predicted using a variety of topological indices that come from the concept of graphs. These models can be used to anticipate a compound's physical characteristics in the absence of experimental testimony⁵. Topological measures are numerical values that are computed from a molecule's chemical graph and are frequently associated with the molecule's chemical characteristics. These indices allow us to predict the properties of compounds without the need for complicated calculations⁶.

Topological indices and co-indices are powerful mathematical descriptors in chemical graph theory. These descriptors provide a quantitative method for defining the structural characteristics of chemical compounds, and they are built from the underlying molecular graph. Notably, there is a high link between several physicochemical features of molecules and topological indices. Examples of these indices that are often used are the Geometric-Arithmetic (GA) index and the Atom-Bond Connectivity (ABC) index, among others⁷. These topological indices have significance because of their foundational chemical relevance. They are remarkably good at predicting a number of important chemical compound attributes, such as stability, strain energy, boiling point, etc. Other topological indices that have been studied include the Zagreb indices, Forgotten index, Wiener index, Yemen index, Hyper-Zagreb and Hyper Forgotten indices, and Kirchhoff index⁸⁻¹¹. These indices have been used to explore quantitative structure-property relationships and quantitative structure-activity relationships in cheminformatics^{12,13}. They provide simple and easily computed chemical information with explicit mathematical formulas. The correlation between these topological indices and physicochemical properties of molecules has been examined.

A. Ashrafi et al. introduced Zagreb coindices and calculated exact formulae for the relationship between the first and second Zagreb indices and their coindices^{14,15} such as:

$$\overline{M}_1(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma(\mu) + \delta_\Gamma(\nu)] \quad 1$$

$$\overline{M}_2(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma(\mu)\delta_\Gamma(\nu)] \quad 2$$

$$\overline{M}_1(\Gamma) = 2m(n-1) - M_1(\Gamma) \quad 3$$

$$\overline{M}_2(\Gamma) = 2m^2 - \frac{1}{2}M_1(\Gamma) - M_2(\Gamma) \quad 4$$

N. De et al.¹⁶ introduced the F-coindex which is defined as follows.

$$\overline{F}(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^2(\mu) + \delta_\Gamma^2(\nu)] \quad 5$$

They also computed the relationship between F-index and F-coindex as:

$$\overline{F}(\Gamma) = (n-1)M_1(\Gamma) - F(\Gamma) \quad 6$$

Alameri et al.¹⁷ defined a new degree-based descriptor, denoted by the (*Y-coindex*), and it is defined as:

$$\overline{Y}(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^3(\mu) + \delta_\Gamma^3(\nu)], \quad 7$$

In addition, in the same paper, (*Y-coindex*) of a graph Γ was defined as:

$$\overline{Y}(\Gamma) = (n-1)F(\Gamma) - Y(\Gamma), \quad 8$$

Sethumadhavan and Gokulathilagan¹⁸ studied the S-index and S-coindex of some graph operations, and they defined as:

$$S(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^4(\mu) + \delta_\Gamma^4(\nu)], \quad 9$$

$$\overline{S}(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^4(\mu) + \delta_\Gamma^4(\nu)]. \quad 10$$

Veylaki, et al.¹⁹ introduced Hyper-Zagreb coindex and it is defined as:

$$\overline{HM}(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma(\mu) + \delta_\Gamma^2(\nu)]^2, \quad 11$$

Gutman I.²⁰ derived the formula for the Hyper-Zagreb coindex as follows:

$$\overline{HM}(\Gamma) = (n-2)M_1(\Gamma) + 4m^2 - HM(\Gamma), \quad 12$$

Ayache et al.²¹ defined the second Hyper-Zagreb coindex such as:

$$\overline{HM}_2(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} \delta_\Gamma^2(\mu) \delta_\Gamma^2(\nu) \quad 13$$

Such that:

$$\overline{HM}_2(\Gamma) = \frac{1}{2}M_1^2(\Gamma) - \frac{1}{2}Y(\Gamma) - HM_2(\Gamma), 14$$

The benzenoid series includes compounds such as circumcoronene and ovalene. These compounds have been studied for their various properties and applications. For example, the topological indices of the double and strong double graphs of the Circumcoronene series of benzenoids have been calculated²². Additionally, Circumcoronene and Boron Nitride Circumcoronene have been explored as carriers for the anticancer drug floxuridine, showing promise as drug delivery systems²³. The interactions of n-alkanes and n-per fluoroalkanes with Circumcoronene have also been investigated, revealing differences in their adsorption energies and steric repulsion²⁴. Furthermore, the Multiplicative Zagreb Eccentricity index has been computed for the Circumcoronene series of benzenoid H_k ²⁵. The Sanskruti index has also been calculated for the Circumcoronene series of benzenoid²⁶.

Let $(\Gamma = H_k : k \geq 1)$ be the Benzenoid Circumcoronene series (Fig 1), then

$$M_1(\Gamma) = 54k^2 - 30k, \quad 15$$

$$M_2(\Gamma) = 81k^2 - 63k + 6, \quad 16$$

$$F(H_k) = 162k^2 - 114k, \quad 17$$

$$HM(H_k) = 324k^2 - 240k + 12. \quad 18$$




Figure 1. The first three graphs from the Benzenoid Circumcoronene series

Numerous studies have examined the computation of topological indices for Benzenoid Circumcoronene. Chemical descriptors, such as Randic, Zagreb, Harmonic, augmented Zagreb, atom-bond connectivity, and geometric-arithmetic indices, have been computed for Benzenoid networks theoretically²⁷. Although numerous studies have

examined the computation of topological indices for Benzenoid Circumcoronene, investigations into the calculation of topological coindices remain scarce. This paper highlights topological coindices of Benzenoid Circumcoronene. Specifically, mathematical formulas have been derived for the first Zagreb, second Zagreb, forgotten, hyper-first Zagreb, hyper-second Zagreb, and hyper-forgotten coindices of the Benzenoid Circumcoronene series. In addition, new topological coindices are introduced and their mathematical formulas are derived for the benzenoid coronene ocean series. Furthermore, some algorithms have been built using Python programs to calculate the mathematical formulas that are derived.

Preliminary Results

In this section, the topological indices of the benzenoid circumcoronene series H_k are derived by us. $Y(H_k)$, $S(H_k)$ and $HM_2(H_k)$,

Lemma 1: For $k \leq 1$, H_k is the benzene series (see Fig 1). The number of vertices and edges of H_k

$$\bullet n = |V(H_k)| = 6k^2,$$

$$\bullet m = |E(H_k)| = 9k^2 - 3k,$$

Table 1. The edge set partition of the benzenoid circumcoronene series

Edge	E_1	E_2	E_3
Cardinality	6	$12(k - 1)$	$9k^2 - 15k + 6$

- $\bullet E_1(H_k) = \{e = \mu\nu \in E(\Gamma) : \delta(\mu) = \delta(\nu) = 2\}$,
- $\bullet E_2(H_k) = \{e = \mu\nu \in E(\Gamma) : \delta(\mu) = 2, \delta(\nu) = 3\}$,
- $\bullet E_3(H_k) = \{e = \mu\nu \in E(\Gamma) : \delta(\mu) = \delta(\nu) = 3\}$,

Proposition 1: The Y-index of benzenoid circumcoronene is given by:

$$Y(H_k) = 486k^2 - 390k.$$

Proof: By the definition of Y- index and (Lemma 1),

$$\begin{aligned} Y(H_k) &= \sum_{\mu\nu \in E(H_k)} [\delta^3(\mu) + \delta^3(\nu)] \\ &= \sum_{\mu\nu \in E_1} [\delta^3(\mu) + \delta^3(\nu)] \end{aligned}$$

$$\begin{aligned} & + \sum_{\mu\nu \in E_2} [\delta^3(\mu) + \delta^3(\nu)] + \\ \sum_{\mu\nu \in E_3} [\delta^3(\mu) + \delta^3(\nu)] & \\ & = 16|E_1| + 35|E_2| + 54|E_3| \\ & = 16(6) + 35(12(k-1)) + \\ 54(9k^2 - 15k + 6) & \\ & = 486k^2 - 390k. \end{aligned}$$

Proposition 2: The HM_2 -index of benzenoid circumcoronene is given by:

$$HM_2(H_k) = 729k^2 - 783k + 150.$$

Proof: By the definition of HM_2 -index and (Lemma 1),

$$\begin{aligned} HM_2(H_k) &= \\ \sum_{\mu\nu \in E(H_k)} [\delta(\mu)\delta(\nu)]^2 &= \sum_{\mu\nu \in E_1} [\delta(\mu)\delta(\nu)]^2 \\ &+ \sum_{\mu\nu \in E_2} [\delta(\mu)\delta(\nu)]^2 + \\ \sum_{\mu\nu \in E_3} [\delta(\mu)\delta(\nu)]^2 & \\ &= 16|E_1| + 36|E_2| + 81|E_3| \end{aligned}$$

Results and Discussion

A mathematical derivation of some old and new topological and Hyper topological coindices is being studied of the benzenoid circumcoronene series H_k and the mathematical formulas derived using the Python program will be supported.

Well-known Topological Coindices of Benzenoid Circumcoronene Series

In this subsection, a mathematical derivation of topological coindices for the benzenoid circumcoronene series H_k is provided by including $\overline{M}_1(H_k)$, $\overline{M}_2(H_k)$, $\overline{F}(H_k)$, $\overline{Y}(H_k)$ and $\overline{S}(H_k)$.

Theorem 2: The M_1 -coindex of benzenoid circumcoronene is given by:

$$\overline{M}_1(H_k) = 108k^4 - 36k^3 - 72k^2 + 36k$$

Proof: From (Lemma 1), and Eq 15,

$$\begin{aligned} n = |V(H_k)| &= 6k^2, \quad m = |E(H_k)| = 9k^2 - 3k, \\ M_1(\Gamma) &= 54k^2 - 30k, \text{ and } \overline{M}_1(\Gamma) = 2m(n-1) - \\ M_1(\Gamma). & \end{aligned}$$

Thus,

$$\begin{aligned} & = 16(6) + 36(12(k-1)) + \\ 81(9k^2 - 15k + 6) & \\ & = 729k^2 - 783k + 150. \end{aligned}$$

Theorem 1: The S -index of benzenoid circumcoronene is given by:

$$S(H_k) = 1458k^2 - 1266k.$$

Proof: From Eq 9,

$$\begin{aligned} S(H_k) &= \sum_{\mu\nu \in E(H_k)} [\delta^4(\mu) + \\ \delta^4(\nu)] &= \sum_{\mu\nu \in E_1} [\delta^4(\mu) + \delta^4(\nu)] \\ &+ \sum_{\mu\nu \in E_2} [\delta^4(\mu) + \delta^4(\nu)] + \\ \sum_{\mu\nu \in E_3} [\delta^4(\mu) + \delta^4(\nu)] & \\ &= 32|E_1| + 97|E_2| + 162|E_3| \\ &= 32(6) + 97(12(k-1)) + \\ 162(9k^2 - 15k + 6) & \\ &= 1458k^2 - 1266k. \end{aligned}$$

$$\begin{aligned} \overline{M}_1(\Gamma) &= 2m(n-1) - M_1(\Gamma) = \\ 2(9k^2 - 3k)[6k^2 - 1] &+ 54k^2 - 30k \\ &= 108k^4 - 36k^3 - 72k^2 + 36k \end{aligned}$$

Theorem 3: The M_2 -coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{M}_2(H_k) = 162k^4 - 108k^3 - 90k^2 + 78k - 6.$$

Proof: From (Lemma 1) and Eqs 15 & 16,

$$\begin{aligned} n = |V(H_k)| &= 6k^2, \quad m = |E(H_k)| = 9k^2 - 3k, \\ M_1(\Gamma) &= 54k^2 - 30k, \quad M_2(\Gamma) = 81k^2 - 63k + 6, \\ \text{and since } \overline{M}_2(\Gamma) &= 2m^2 - \frac{1}{2}M_1(\Gamma) - M_2(\Gamma). \end{aligned}$$

Thus,

$$\begin{aligned} \overline{M}_2(\Gamma) &= 2m^2 - \frac{1}{2}M_1(\Gamma) - M_2(\Gamma) \\ &= 2(9k^2 - 3k)^2 - \frac{1}{2}[54k^2 - 30k] - (81k^2 \\ &\quad - 63k + 6) \\ &= 162k^4 - 108k^3 - 90k^2 + 78k - 6. \end{aligned}$$

Theorem 4: The F-coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{F}(H_k) = 324k^4 - 180k^3 - 216k^2 + 144k.$$

Proof: From (Lemma 1), and Eq 15, 17

$$n = |V(H_k)| = 6k^2, \quad m = |E(H_k)| = 9k^2 - 3k, \\ M_1(\Gamma) = 54k^2 - 30k, \quad F(H_k) = 162k^2 - 114k, \\ \text{and since } \overline{F}(\Gamma) = (n - 1)M_1(\Gamma) - F(\Gamma).$$

Thus,

$$\overline{F}(\Gamma) = (n - 1)M_1(\Gamma) - F(\Gamma) = (6k^2 - 1)(54k^2 - 30k) - (162k^2 - 114k) \\ = 324k^4 - 180k^3 - 216k^2 + 144k.$$

Theorem 5: The Y-coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{Y}(H_k) = 972k^4 - 684k^3 - 648k^2 + 504k.$$

Proof: From (Lemma 1), Eq 17, and (Proposition 2),

$$n = |V(H_k)| = 6k^2, \quad F(H_k) = 162k^2 - 114k, \\ Y(H_k) = 486k^2 - 390k, \text{ and since}$$

$$\overline{Y}(\Gamma) = (n - 1)F(\Gamma) - Y(\Gamma),$$

Thus,

$$\overline{Y}(\Gamma) = (n - 1)F(\Gamma) - Y(\Gamma) \\ = (6k^2 - 1)(162k^2 - 114k) - (486k^2 - 390k) \\ = 972k^4 - 684k^3 - 648k^2 + 504k.$$

Theorem 6: The S-coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{S}(H_k) = 2916k^4 - 2376k^3 - 1944k^2 + 1656k.$$

Proof: The formula of the general first Zagreb coindex is given by :

$$\overline{M}_1^\alpha(\Gamma) = (n - 1)M_1^{\alpha-1}(\Gamma) - M_1^\alpha(\Gamma), \quad 19$$

$$\text{Since } M_1^\alpha = \sum_{\mu\nu \in E} (\delta^\alpha(\mu) + \delta^\alpha(\nu))$$

$$\text{When } \alpha = 1 \quad M_1 = M_1^1 = \sum_{\mu\nu \in E} (\delta(\mu) + \delta(\nu))$$

$$\alpha = 2 \quad F = M_1^2 = \sum_{\mu\nu \in E} (\delta^2(\mu) + \delta^2(\nu))$$

$$\alpha = 3 \quad Y = M_1^3 = \sum_{\mu\nu \in E} (\delta^3(\mu) + \delta^3(\nu))$$

$$\alpha = 4 \quad S = M_1^4 = \sum_{\mu\nu \in E} (\delta^4(\mu) + \delta^4(\nu)).$$

Then,

$$\overline{M}_1^4(\Gamma) = \overline{S}(\Gamma) = (n - 1)Y(\Gamma) - S(\Gamma), \text{ and}$$

$$n = |V(H_k)| = 6k^2, \quad Y(H_k) = 486k^2 - 390k, \quad S(H_k) = 1458k^2 - 1266k.$$

Thun,

$$\overline{S}(H_k) = (n - 1)Y(H_k) - S(H_k) \\ = (6k^2 - 1)(486k^2 - 390k) - (1458k^2 - 1266k) \\ = 2916k^4 - 2376k^3 - 1944k^2 + 1656k.$$

New Topological Indices and their Coindices of Benzenoid Circumcoronene

In this subsection, a mathematical derivation of new topological indices and their coindices of the benzenoid circumcoronene series H_k is provided, denoted Gaza index and Palestine index (in short G -index " $G(H_k)$ " and P -index " $P(H_k)$ ") respectively. The topological coindices of these indices denoted as $\overline{G}(H_k)$ and $\overline{P}(H_k)$. G -index and P -index and their coindices are special cases of the first general Zagreb index and the first general coindex where $\alpha = 5$ and $\alpha = 6$ which define us:

$$G(\Gamma) = \sum_{v \in V(\Gamma)} [\delta_\Gamma^6 v] = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^5 \mu + \delta_\Gamma^5 \nu], \quad 20$$

$$P(\Gamma) = \sum_{v \in V(\Gamma)} [\delta_\Gamma^7 v] = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^6 \mu + \delta_\Gamma^6 \nu], \quad 21$$

Therefore,

$$\overline{G}(\Gamma) = (n - 1)S(\Gamma) - G(\Gamma), \quad 22$$

and

$$\overline{P}(\Gamma) = (n - 1)G(\Gamma) - P(\Gamma) \quad 23$$

Proposition 3: The G -index of benzenoid circumcoronene is given by:

$$G(H_k) = 4374k^2 - 3990k.$$

Proof: By the definition of G -index and (Lemma 1),

$$\begin{aligned}
 G(H_k) &= \sum_{\mu\nu \in E(H_k)} [\delta^5(\mu) + \delta^5(\nu)] \\
 &= \sum_{\mu\nu \in E_1} [\delta^5(\mu) + \delta^5(\nu)] \\
 &+ \sum_{\mu\nu \in E_2} [\delta^5(\mu) + \delta^5(\nu)] + \sum_{\mu\nu \in E_3} [\delta^5(\mu) + \delta^5(\nu)] \\
 &= 64|E_1| + 275|E_2| + 486|E_3| \\
 &= 64(6) + 275(12(k-1)) + \\
 &486(9k^2 - 15k + 6) \\
 &= 4374k^2 - 3990k.
 \end{aligned}$$

Theorem 7: The G-coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{G}(H_k) = 8748k^4 - 7596k^3 - 5832k^2 + 5256k.$$

Proof: We have $n = |V(H_k)| = 6k^2$, $S(H_k) = 1458k^2 - 1266k$, $G(H_k) = 4374k^2 - 3990k$, and since $\overline{G}(\Gamma) = (n-1)S(\Gamma) - G(\Gamma)$,

Thus,

$$\begin{aligned}
 \overline{G}(H_k) &= (n-1)S(H_k) - \\
 G(H_k) &= (6k^2 - 1)(1458k^2 - 1266k) - \\
 &(486k^2 - 390k) \\
 &= 8748k^4 - 7596k^3 - \\
 &5832k^2 + 5256k.
 \end{aligned}$$

Proposition 4: The P-index of benzenoid circumcoronene is given by:

$$P(H_k) = 13122k^2 - 12354k.$$

Proof: By the definition of P- index and (Lemma 1),

$$\begin{aligned}
 P(H_k) &= \sum_{\mu\nu \in E(H_k)} [\delta^6(\mu) + \\
 \delta^6(\nu)] &= \sum_{\mu\nu \in E_1} [\delta^6(\mu) + \delta^6(\nu)] \\
 &+ \sum_{\mu\nu \in E_2} [\delta^6(\mu) + \delta^6(\nu)] + \\
 \sum_{\mu\nu \in E_3} &[\delta^6(\mu) + \delta^6(\nu)] \\
 &= 64|E_1| + 275|E_2| + 486|E_3| \\
 &= 13122k^2 - 12354k.
 \end{aligned}$$

Theorem 8: The P-coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{P}(H_k) = 26244k^4 - 23940k^3 - 17496k^2 + 16344k.$$

Proof: We have $n = |V(H_k)| = 6k^2$, $G(H_k) = 4374k^2 - 3990k$, $P(H_k) = 13122k^2 - 12354k$, and since $\overline{P}(H_k) = (n-1)G(H_k) - P(H_k)$,

Thus,

$$\begin{aligned}
 \overline{P}(H_k) &= (n-1)G(H_k) - P(H_k) \\
 &= (6k^2 - 1)(4374k^2 - 3990k) \\
 &- (13122k^2 - 12354k) \\
 &= 26244k^4 - 23940k^3 - \\
 &17496k^2 + 16344k.
 \end{aligned}$$

Well-known Hyper Topological Coindices of Benzenoid Circumcoronene

In this subsection, a mathematical derivation of hyper topological coindices of the benzenoid circumcoronene series H_k is provided such as $\overline{HM}_1(H_k)$, $\overline{HM}_2(H_k)$ and $\overline{HF}(H_k)$.

Theorem 9: The HM_1 -coindex of benzenoid circumcoronene is given by:

$$\overline{HM}_1(H_k) = 648k^4 - 396k^3 - 396k^2 + 300k - 12.$$

Proof: From the definition of hyper Zagreb index,

$$\begin{aligned}
 HM_1(\Gamma) &= \sum_{\mu\nu \in E(\Gamma)} [\delta(\mu) + \\
 \delta(\nu)]^2 &= \sum_{\mu\nu \in E(\Gamma)} [\delta^2(\mu) + \delta^2(\nu) + 2\delta(\mu)\delta(\nu)] \\
 &= \sum_{\mu\nu \in E(\Gamma)} [\delta^2(\mu) + \delta^2(\nu)] + \\
 \sum_{\mu\nu \in E(\Gamma)} &[2\delta(\mu)\delta(\nu)] = F(\Gamma) + 2M_2(\Gamma).
 \end{aligned}$$

Therefore,

$$\overline{HM}_1(\Gamma) = \overline{F}(\Gamma) + 2\overline{M}_2(\Gamma), \quad 24$$

Applying Theorems (3 & 4) in Eq 24

$$\overline{HM}_1(H_k) = \overline{F}(H_k) + 2\overline{M}_2(H_k) = 648k^4 - 396k^3 - 396k^2 + 300k - 12.$$

Theorem 10: The HM_2 -coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{HM}_2(H_k) = 1458k^4 - 1620k^3 - 522k^2 + 978k - 150.$$

Proof: By Eq 15, and Propositions 1 & 2,

$$M_1(H_k) = 54k^2 - 30k, \quad Y(H_k) = 486k^2 - 390k, \\ HM_2(H_k) = 3[243k^2 - 261k + 50],$$

and by substitution on $\overline{HM}_2(H_k) = \frac{1}{2}M_1^2(H_k) - \frac{1}{2}Y(H_k) - HM_2(H_k)$, the required was obtained.

Theorem 11: The HF-coindex of benzenoid circumcoronene is given by:

$$\overline{HF}(H_k) = 5832k^4 - 5580k^3 - 2988k^2 + 3612k - 588.$$

Proof: From the definition of hyper F-index,

$$HF(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta(\mu)^2 + \delta^2(\nu)]^2 = \sum_{\mu\nu \in E(\Gamma)} [\delta^4(\mu) + \delta^4(\nu) + 2\delta^2(\mu)\delta^2(\nu)] \\ = \sum_{\mu\nu \in E(\Gamma)} [\delta^4(\mu) + \delta^4(\nu)] + 2\sum_{\mu\nu \in E(\Gamma)} [\delta^2(\mu)\delta^2(\nu)] = S(\Gamma) + 2HM_2(\Gamma).$$

Therefore,

$$\overline{HF}(\Gamma) = \overline{S}(\Gamma) + 2\overline{HM}_2(\Gamma), \quad 25$$

Applying Theorems (6 & 10) in (Eq. 25),

$$\overline{HF}(H_k) = \overline{S}(H_k) + 2\overline{HM}_2(H_k) = 5832k^4 - 5580k^3 - 2988k^2 + 3612k - 588.$$

New Hyper Topological Coindices of Benzenoid Circumcoronene Series

In this subsection, a mathematical derivation of the new hyper topological index and their coindex of the benzenoid circumcoronene series H_k is provided, denoted hyper Yemen index (in short HY -index " $HY(H_k)$ ") which the hyper topological coindex of this index is $\overline{HY}(H_k)$. It is important here to introduce a new topological index for use in deriving the formula of the hyper topological index and their coindex, denoted Quds index (in short Q -index " $Q(H_k)$ "). which the topological coindex of this index is $\overline{Q}(H_k)$.

The definitions of HY -index and Q -index are:

$$HY(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^3 \mu + \delta_\Gamma^3 \nu]^2, \quad 26$$

$$Q(\Gamma) = \sum_{\mu\nu \in E(\Gamma)} [\delta_\Gamma^3 \mu \delta_\Gamma^3 \nu], \quad 27$$

In addition, the definitions of HY -coindex and Q -coindex are as follows.

$$\overline{HY}(\Gamma) = \sum_{\mu\nu \notin E(\Gamma)} [\delta_\Gamma^3 \mu + \delta_\Gamma^3 \nu]^2, \quad 28$$

$$\overline{Q}(\Gamma) = \sum_{\mu\nu \notin E(\Gamma)} [\delta_\Gamma^3 \mu \delta_\Gamma^3 \nu], \quad 29$$

It's easy to see that the M_2 -index, the HM_2 -index and the Q -index are special cases of the second general Zagreb indices where $\alpha = 1$, $\alpha = 2$, and $\alpha = 3$, respectively.

Lemma 2: The Q -coindex of a graph Γ is given by:

$$\overline{Q}(\Gamma) = \frac{1}{2}(F^2(\Gamma) - G(\Gamma)) - Q(\Gamma) \quad 30$$

Proof: From the definitions of Q index and their coindex,

$$Q(\Gamma) + \overline{Q}(\Gamma) = (\sum_{\mu\nu \in E(\Gamma)} + \sum_{\mu\nu \notin E(\Gamma)}) (\delta^3(\mu)\delta^3(\nu)) \\ = \frac{1}{2} [(\sum_{\mu \in V(\Gamma)} \sum_{\nu \in V(\Gamma)}) (\delta^3(\mu)\delta^3(\nu)) - \sum_{(\mu=\nu) \in V(\Gamma)} (\delta^3(\mu)\delta^3(\nu))] \\ = \frac{1}{2} [\sum_{\mu \in V(\Gamma)} (\delta^3(\mu)) \sum_{\nu \in V(\Gamma)} (\delta^3(\nu)) - \sum_{(\mu=\nu) \in V(\Gamma)} (\delta^6(\nu))] \\ = \frac{1}{2} [F(\Gamma)F(\Gamma) - G(\Gamma)] = \frac{1}{2} [F^2(\Gamma) - G(\Gamma)]$$

Proposition 4: The Q -index of benzenoid circumcoronene is given by:

$$Q(H_k) = 6561k^2 - 8343k + 2166$$

Proof: By the definition of Q -index and (Lemma 1),

$$Q(H_k) = \sum_{\mu\nu \in E(H_k)} [\delta(\mu)\delta(\nu)]^3 = \sum_{\mu\nu \in E_1} [\delta(\mu)\delta(\nu)]^3 + \sum_{\mu\nu \in E_2} [\delta(\mu)\delta(\nu)]^3 + \sum_{\mu\nu \in E_3} [\delta(\mu)\delta(\nu)]^3 \\ = 64|E_1| + 216|E_2| + 729|E_3| \\ = 64(6) + 216(12(k-1)) + 729(9k^2 - 15k + 6)$$

$$= 6561k^2 - 8343k + 2166.$$

Theorem 12: The Q-coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{Q}(H_k) = 13122k^4 - 18468k^3 - 2250k^2 + 10338k - 2166.$$

Proof: By Lemma 2,

$$F(H_k) = 162k^2 - 114k, \quad G(H_k) = 4374k^2 - 3990k, \quad Q(H_k) = 6561k^2 - 8343k + 2166,$$

and by substitution on $\overline{Q}(\Gamma) = \frac{1}{2}(F^2(\Gamma) - G(\Gamma)) - Q(\Gamma)$, the required was obtained.

Lemma 3: The HY-index is a linear composition of the P-index and Q-index, in other words, we can write

$$HY(\Gamma) = P(\Gamma) + 2Q(\Gamma) \quad 31$$

Proof: From the definition of $HY(\Gamma)$,

$$\begin{aligned} HY(\Gamma) &= \sum_{\mu\nu \in E(\Gamma)} [\delta_{\Gamma}^3\mu + \delta_{\Gamma}^3\nu]^2 \\ &= \sum_{\mu\nu \in E(\Gamma)} [\delta_{\Gamma}^6\mu + \delta_{\Gamma}^6\nu] + 2 \sum_{\mu\nu \in E(\Gamma)} \delta_{\Gamma}^3(\mu)\delta_{\Gamma}^3(\nu) = P(\Gamma) + 2Q(\Gamma). \end{aligned}$$

Proposition 5: The HY-index of benzenoid circumcoronene is given by:

$$HY(H_k) = 26244k^2 - 29040k + 4332$$

Proof: By Lemma 3,

$$P(H_k) = 13122k^2 - 12354k, \quad Q(H_k) = 6561k^2 - 8343k + 2166,$$

and by substitution on $HY(H_k) = P(H_k) + 2Q(H_k)$, we get the required.

Theorem 13: The HY-coindex of benzenoid circumcoronene (H_k) is given by:

$$\overline{HY}(H_k) = 52488k^4 - 60876k^3 - 21996k^2 + 37020k - 4332.$$

Proof: By Lemma 2 and Lemma 3,

$$\overline{P}(H_k) = 26244k^4 - 23940k^3 - 17496k^2 + 16344k,$$

$$\overline{Q}(H_k) = 13122k^4 - 18468k^3 - 2250k^2 + 10338k - 2166,$$

and by substitution on $\overline{HY}(H_k) = \overline{P}(H_k) + 2\overline{Q}(H_k)$, the required was obtained.

Discuss 1: In the previous part, we derived mathematical formulas for some of the topological descriptors of the benzenoid circumcoronene series by taking advantage of the properties of the compound listed in Table 1. In the next part, we will calculate these formulas in general via the Python program. Also, we will address these formulas as groups in special cases, and each group will be discussed independently.

An Algorithm to Calculate Topological and Hyper-Topological Indices and their Coindices

This section presents a program to implement an algorithm for calculating topological and Hyper-topological indices and their coindices of benzenoid circumcoronene series using Python (see Figs 2,3,4, and 5).

Some special cases of these topological indices will be computed (specifically when the parameter $k = 1, 2, \dots, 8$) and then study the relationship between these indicators by analyzing these data.


```

k = int(input("k = "))

M1=54*k**2-30*k
M2=81*k**2-63*k+6
F =162*k**2 -114*k
Y =486*k**2 - 390*k
S =1458*k**2 - 1266*k
G =4374*k**2 - 3990*k
Q =6561*k**2-8343*k+2166
P = 13122*k**2 - 12354*k

print("M1-index : {}".format(M1))
print("M2-index : {}".format(M2))
print("F-index : {}".format(F))
print("Y-index : {}".format(Y))
print("S-index : {}".format(S))
print("G-index : {}".format(G))
print("Q-index : {}".format(Q))
print("P-index : {}".format(P))
    
```

Figure 2. Topological Indices of the Benzenoid Circumcoronene Series Using PYTHON

Discuss 2: Table 2 presented quantitatively examines eight benzenoid circumcoronene structures of increasing size as denoted by the parameter k , ranging from 1 to 8. It compares the values of several commonly used topological indices in this series to assess their trends concerning molecular size growth. The values of the indices increase with increasing k , indicating that the molecules become more complex as the number of rings in the circumcoronene series

benzenoid H_k increases. Specifically, the First and Second Zagreb indices (M_1 and M_2 , respectively), which are calculated based on vertex degrees, both predictably increase stepwise with each additional k -value. Similarly, the Forgotten, Yemen, S, G, Q, and P indices, which are also based on vertex degrees, in their formulation, also increase more quickly and monotonically from $k = 1$ to $k = 8$, albeit at varying rates due to their distinct mathematical definitions.

Table 2. Topological indices (TI) for benzenoid circumcoronene series H_k , $k = 1, 2, \dots, 8$

<i>TI&k</i>	1	2	3	4	5	6	7	8
<i>M1</i>	24	156	396	744	1200	1764	2436	3216
<i>M2</i>	24	204	546	1050	1716	2544	3534	4686
<i>F</i>	48	420	1116	2136	3480	5148	7140	9456
<i>Y</i>	96	1164	3204	6216	10200	15156	21084	27984
<i>S</i>	192	3300	9324	18264	30120	44892	62580	83184
<i>G</i>	384	9516	27396	54024	89400	133524	186396	248016
<i>Q</i>	384	11724	36186	73770	124476	188304	265254	355326
<i>P</i>	768	27780	81036	160536	266280	398268	556500	740976

```

k = int(input("k = "))

HM1 = 324*k**2 - 240*k + 12
HM2 = 729*k**2 - 783*k + 150
HF = 2916*k**2 - 2832*k + 300
HY = 26244*k**2 - 29040*k + 4332

print("HM1-index : {}".format(HM1))
print("HM2-index : {}".format(HM2))
print("HF-index : {}".format(HF))
print("HY-index : {}".format(HY))
    
```

Figure 3. Hyper Topological Indices of the Benzenoid Circumcoronene Series Using PYTHON

Discuss 3: Table 3 presents the values of four hyper-topological indices (HTIs) for the benzenoid circumcoronene series H_k , $k=1,2,\dots,8$. Specifically, it monitors the Hyper-first Zagreb index (HM_1), Hyper-second Zagreb index (HM_2), Hyper-Forgotten index (HF), and Hyper Yemen index (HY). All indices presented in the table are generalized hyper-topological descriptors that account for higher-order

interactions between vertices, and edges in molecular graphs. As with traditional topological indices, monotonic increasing trends are observed concerning the size parameter k for each index type. For instance, HY exhibits the steepest rises relative to HM_1 , HM_2 , and HF due to its inclusion of longer-range topological features in its definition.

Table 3. Hyper topological indices (HTI) for benzenoid circumcoronene series H_k , $k = 1, 2, \dots, 8$

<i>HTI&k</i>	1	2	3	4	5	6	7	8
<i>HM1</i>	96	828	2208	4236	6912	10236	14208	18828
<i>HM2</i>	96	1500	4362	8682	14460	21696	30390	40542
<i>HF</i>	384	6300	18048	35628	59040	88284	123360	164268
<i>HY</i>	1536	51228	153408	308076	515232	774876	1087008	1451628

```

k = int(input("k = "))

M1c=108*k**4 - 36*k**3 - 72*k**2 + 36*k
M2c=162*k**4 - 108*k**3 - 90*k**2 + 78*k - 6
Fc=324*k**4 - 180*k**3 - 216*k**2 + 144*k
Yc=972*k**4 - 684*k**3 - 648*k**2 + 504*k
Sc=2916*k**4 - 2376*k**3 - 1944*k**2 + 1656*k
Gc=8748*k**4 - 7596*k**3 - 5832*k**2 + 5256*k
Qc=13122*k**4 - 18468*k**3 - 2250*k**2 + 10338*k - 2166
Pc= 26244*k**4 - 23940*k**3 - 17496*k**2 + 16344*k

print("M1-coindex : {}".format(M1c))
print("M2-coindex : {}".format(M2c))
print("F-coindex : {}".format(Fc))
print("Y-coindex : {}".format(Yc))
print("S-coindex : {}".format(Sc))
print("G-coindex : {}".format(Gc))
print("Q-coindex : {}".format(Qc))
print("P-coindex : {}".format(Pc))
    
```

Figure 4. Topological Coindices of the Benzenoid Circumcoronene Series Using PYTHON

Discuss 4: Table 4 presents the values of various topological coindices for an increasingly large series of benzenoid circumcoronene structures, denoted as H_k where $k = 1, 2, \dots, 8$ represents the size of the structure. Specifically, it reports the First Zagreb coindex (M_1^c), Second Zagreb coindex (M_2^c), Forgotten coindex (F^c), Yemen coindex (Y^c), S-coindex (S^c), G-coindex (G^c), Q-coindex (Q^c), and P-coindex (P^c) for each value of k . The coindices generally increase consistently with larger k values,

as would be expected as the size and complexity of the circumcoronene structures grow with each additional ring. M_1^c displays the lowest values while P^c exhibits the highest values across all k . Some coindices like M_1^c , M_2^c , and F^c show relatively linear increases, while others like Y^c , S^c , G^c , Q^c , and P^c rise more sharply suggesting they are more strongly influenced by the rapid topological changes occurring in these structures as they expand in size.

Table 4. Topological coindices (TC) for benzenoid circumcoronene series H_k , $k = 1, 2, \dots, 8$

<i>TC&k</i>	1	2	3	4	5	6	7	8
<i>M1c</i>	36	1224	7236	24336	61380	129816	243684	419616
<i>M2c</i>	36	1518	9624	33426	85884	183846	348048	603114
<i>Fc</i>	72	3168	19872	68544	175320	374112	706608	1222272
<i>Yc</i>	144	8496	55944	196704	508320	1091664	2070936	3593664
<i>Sc</i>	252	23184	159516	569952	1485180	3205872	6102684	10616256
<i>Gc</i>	576	66384	466776	1681056	4398480	9518256	18149544	31611456
<i>Qc</i>	576	71718	572844	2180466	5886024	12995886	25131348	44228634
<i>Pc</i>	1152	191088	1370952	4971744	13054320	28309392	54057528	94249152

```

k = int(input("k = "))

HM1c=648*k**4 -396*k**3 -396*k**2 +300*k -12
HM2c=1458*k**4 -1620*k**3 -522*k**2 +978*k -150
HFc=5832*k**4 -5580*k**3 -2988*k**2 +3612*k -588
HYc=52488*k**4-60876*k**3-21996*k**2+37020*k-4332

print("HM1-coindex :{}".format(HM1c))
print("HM2-coindex :{}".format(HM2c))
print("HF-coindex :{}".format(HFc))
print("HY-coindex :{}".format(HYc))
    
```

Figure 5. Hyper Topological Coindices of the Benzenoid Circumcoronene Series Using PYTHON

Discuss 5: Table 5 presents data on the first Hyper Zagreb coindex, second Hyper Zagreb coindex, Hyper-Forgotten coindex, and Hyper Yemen coindex for benzenoid circumcoronene series H_k of increasing order (k) from 1 to 8. The values of each coindex increase substantially with increasing k , demonstrating a higher topological complexity for benzenoid graphs of larger size according to these metrics. It is noteworthy that the second Hyper

Zagreb coindex yields consistently higher values than the first Hyper Zagreb coindex, while the Hyper-Forgotten coindex and Hyper Yemen coindex present much larger figures overall, indicating greater sensitivity to graph order for these coindices. This analysis of quantitative topological properties aids in characterizing the structural evolution of benzenoid circumcoronene molecular graphs.

Table 5: Hyper topological coindices (HTC) for benzenoid circumcoronene series H_k , $k = 1, 2, \dots, 8$

<i>HTC&k</i>	1	2	3	4	5	6	7	8
<i>HM1c</i>	144	6204	39120	135396	347088	741804	1402704	2428500
<i>HM2c</i>	144	10086	72444	264978	700440	1526574	2926116	5116794
<i>HFc</i>	288	43356	305088	1101924	2890272	6266508	11966976	20867988
<i>HYc</i>	2304	334524	2516640	9332676	24826368	54301164	104320224	182706420

Conclusion

This study contributes to advancing the quantitative structure-topology analysis of benzenoid circumcoronenes by calculating various topological coindices. Although previous work has computed indices for this important class of polycyclic aromatic hydrocarbons, investigations into coindex properties were limited. The present research addresses this gap by deriving mathematical expressions for key coindices describing the circumcoronene topology, including Zagreb, forgotten, Yemen, S, hyper-Zagreb, and hyper-forgotten coindices. The trends in these coindices as a function of molecular size were then systematically analyzed. Moreover, new topological indices and their coindices are introduced, and mathematical derivations of the benzenoid circumcoronene series

are studied. All derived formulas are supported by algorithms using the Python program. The results show that the topological coindices increase steadily with the parameter k , validly encoding and increasing connective complexity. Notably, hyper-topological descriptors more strongly accentuate escalating superstructural characteristics. Overall, this computational analysis enhances understanding of the intricate architectural principles that govern the composition and properties of circumcoronene from a quantitative graph theory perspective. The growth formulas of the topological coindices provide a framework supporting future modeling of the behavior of circumcoronenes and the design of novel structural analogs.

Authors' Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for re-publication, which is attached to the manuscript.
- No animal studies are present in the manuscript.
- No human studies are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee at University of Science and Technology, Sana'a, Yemen

Authors' Contribution Statement

W. S. conceived of the idea, wrote the manuscript, and proofread it. A. A. designed the research, performed the computations, and acquired data. M. A. implemented the research, analyzed, interpreted, and drafted the results. N. A. verified the methods,

reviewed the results, and supervised this work. B. A. reviewed the work, designed the program, and linguistic checking. All authors discussed the results and contributed to the final manuscript.

References

1. Lewis, R. Editorial for the Special Issue on "Algorithms for Graphs and Networks". *Algorithms*. 2020; 13(11): 292. <https://doi.org/10.3390/a13110292>
2. Gutman I, Trinajstić N, Zivković T. Graph Theory and Molecular Orbitals—VI: A Discussion of Non-Alternant Hydrocarbons. *Tetrahedron*. 1973; 29(21): 3449-3454. [http://dx.doi.org/10.1016/S0040-4020\(01\)93501-X](http://dx.doi.org/10.1016/S0040-4020(01)93501-X)
3. Wagner S, Wang H. Introduction to Chemical Graph Theory. 1st edition. New York: Chapman and Hall/CRC; 2018. 269 p. <https://doi.org/10.1201/9780429450532>

4. Estrada E, Bonchev D. Chemical Graph Theory. In Gross JL, Yellen J, Zhang P, editors, Handbook of Graph Theory. Discrete Mathematics and Its Applications Series; 83. 2nd ed. USA: Chapman and Hall/CRC; 2013; 1538-1558. <https://doi.org/10.1201/b16132>.
5. Burch KJ. Chapter 8 - Chemical Applications of Graph Theory. In: Blinder SM, House JE, editors. Mathematical Physics in Theoretical Chemistry [Internet]. Elsevier; 2019: 261–94. (Developments in Physical & Theoretical Chemistry). <https://doi.org/10.1016/B978-0-12-813651-5.00008-5>
6. Diudea MV. Multi-Shell Polyhedral Clusters. Carbon Materials: Chemistry and Physics book series (CMCP). Springer; 2018; 10 Chap 1, Basic Chemical Graph Theory; p.1-21. https://doi.org/10.1007/978-3-319-64123-2_1
7. Rizwan M, Bhatti AA, Javaid M, Jarad F. Some Bounds on Bond Incident Degree Indices with Some Parameters. Math Probl Eng. 2021; 2021: 1-10. <https://doi.org/10.1155/2021/8417486>
8. Abdelgader SM, Wang C, Mohammed SA. Computation of Topological Indices of Some Special Graphs. Mathematics. 2018; 6(3): 33. <https://doi.org/10.3390/math6030033>
9. Roy K, Kar S, Das RN. Understanding the Basics of QSAR for Applications in Pharmaceutical Sciences and Risk Assessment. USA: Academic Press; 2015. Chap4, Topological QSAR; p. 103-149. <https://doi.org/10.1016/B978-0-12-801505-6.00004-1>
10. Mahboob A, Rasheed MW, Bayati JHH, Hanif I. Computation of Several Banhatti and Reven Invariants of Silicon Carbides. Baghdad Sci J. 2023; 20(3(Suppl.)): 1099-1099. <http://dx.doi.org/10.21123/bsj.2023.8212>
11. Mekala A, Murali R. Some K-Banhatti Polynomials of First Dominating David Derived Networks. Baghdad Sci J. 2023; 20(2): 0349-0349. <http://dx.doi.org/10.21123/bsj.2022.6996>
12. Zaman S, Jalani M, Ullah A, Saedi G. Structural Analysis and Topological Characterization of Sudoku Nanosheet. J Math. 2022; 2022: 1-10. <https://doi.org/10.1155/2022/5915740>
13. Wang Y, Yousaf S, Bhatti AA, Aslam A. Analyzing the Expressions for Nanostructures via Topological Indices. Arab J Chem. 2022; 15(1): 1-9. <https://doi.org/10.1016/j.arabjc.2021.103469>
14. Ashrafi A, Doslic T, Hamzeh A. The Zagreb Coindices of Graph Operations. Discrete Appl Math. 2010; 158(15): 1571-1578. <https://doi.org/10.1016/j.dam.2010.05.017>.
15. Malar JS, Meenakshi A. Forgotten Index and Forgotten Coindex of Graphs. Baghdad Sci J. 2023; 21(2): 0504-0510. <http://dx.doi.org/10.21123/bsj.2023.8366>
16. De N, Nayeem S, Pal A. The F-coindex of Some Graph Operations. SpringerPlus. 2016; 5(1): 221. <http://dx.doi.org/10.1186/s40064-016-1864-7>
17. Alameri A, Al-Rumaima M, Almazah M. Y-coindex of Graph Operations and its Applications of Molecular Descriptors. J Mol Struct. 2020; 1221: 128754. <https://doi.org/10.1016/j.molstruc.2020.128754>.
18. Nagarajan S, Gokulathilagan K. S-coindex of Some Graph Operations and its Properties and Conjugated Polymers. Glob J Eng Technol Adv. 2023; 16(02): 140-149. <https://doi.org/10.30574/gjeta.2023.16.2.0151>.
19. Veylaki M, Nikmehr MJ, Tavallaee HA. The Third and Hyper-Zagreb Coindices of Some Graph Operations. J Appl Math Comput. 2016; 50: 315-325. <https://doi.org/10.1007/s12190-015-0872-z>.
20. Gutman I. On Hyper-Zagreb Index and Coindex. Bulletin (Académie serbe des sciences et des arts. Classe des sciences mathématiques et naturelles. Sciences mathématiques). 2017; (42): 1-8.
21. Ayache A, Alameri A, Alsharafi M, Ahmed H. The Second Hyper-Zagreb Coindex of Chemical Graphs and Some Applications. J Chem. 2021; 2021: 1-8. <http://dx.doi.org/10.1155/2021/3687533>
22. Sardar MS, Siddique I, Alrowaili D, Ali MA, Akhtar S. Computation of Topological Indices of Double and Strong Double Graphs of Circumcoronene Series of Benzenoid. J Math. 2022; 2022: 1-11. <https://doi.org/10.1155/2022/5956802>.
23. Khudhair AM, Ben Ahmed A. Utilizing Circumcoronene and BN Circumcoronene for the Delivery and Adsorption of the Anticancer Drug Floxuridine. Comput Theor Chem. 2023; 1222: 114075. <https://doi.org/10.1016/j.comptc.2023.114075>
24. Kikkawa Y, Tsuzuki S. Analysis of Intermolecular Interactions of n-perfluoroalkanes with Circumcoronene Using Dispersion-Corrected DFT Calculations: Comparison with those of n-alkanes. Phys Chem Chem Phys. 2023; 25(16): 11331-11337. <https://doi.org/10.1039/D3CP00790A>
25. Gao Y, Farahani MR, Nazeer W. On Topological Indices of Circumcoronene Series of Benzenoid. Chem Methodol. 2018; 2(1): 39-46. <https://doi.org/10.22631/chemm.2017.99300.1013>
26. Gao Y, Farahani M, Sardar M, Zafar S. On the Sanskruti Index of Circumcoronene Series of Benzenoid. Appl Math. 2017; 8: 520-524. <http://dx.doi.org/10.4236/am.2017.84041>
27. Ali DA, Haidar Ali, Qurat Ul Ain, Kirmani SAK, Ali P, Sesay M. Computation of Benzenoid Planar Octahedron Networks by Using Topological Indices. Math Probl Eng. 2023; 2023: 1-16. <https://doi.org/10.1155/2023/2686873>

التطبيقات التبولوجية والتبولوجية المفرطة لسلسلة بنزينويد سيركوم كورونين

وليد يوسف¹، عبده العامري²، محمد الشرفي²، نعمان النجار²، بلال الفهيد³

¹قسم تقنية معلومات، كلية الحاسبات وتكنولوجيا المعلومات، جامعة العلوم والتكنولوجيا، صنعاء، اليمن.

²قسم الهندسة الطبية الحيوية، كلية الهندسة، جامعة العلوم والتكنولوجيا، صنعاء، اليمن.

³قسم علوم الحاسوب، كلية الحاسبات وتكنولوجيا المعلومات، جامعة العلوم والتكنولوجيا، صنعاء، اليمن.

الخلاصة

يتم تمكين العديد من التطبيقات في الكيمياء من خلال نظرية الرسم البياني الكيميائي، وهو فرع من نظرية الرسم البياني. تُستخدم الكميات العددية المشتقة من الرسوم البيانية الكيميائية للجزيء، والمعروفة باسم المؤشرات التبولوجية والمؤشرات التبولوجية المشتركة، لنمذجة الخواص الكيميائية والفيزيائية للجزيئات في أبحاث العلاقات الكمية بين البنية والملكية (QSPR) والعلاقات الكمية بين البنية والنشاط (QSAR). لحسن الحظ، وجدت التجارب القائمة على المواد الكيميائية علاقة قوية بين الواصفات التبولوجية (المؤشرات التبولوجية والمؤشرات التبولوجية المشتركة) للهياكل الجزيئية وخصائصها الفيزيائية والكيميائية، مثل نقطة الغليان، وسمية الأدوية. على الرغم من أن العديد من التقارير البحثية قد ساهمت في حساب المؤشرات التبولوجية لسلسلة البنزينويد سيركوم كورونين، إلا أن الدراسات حول حساب المؤشرات التبولوجية المشتركة محدودة. تركز هذه الورقة على بعض المؤشرات التبولوجية المشتركة. تم اشتقاق العديد من صيغ المؤشرات التبولوجية المشتركة مثل المؤشر المشترك لزغرب الأول والمؤشر المشترك لزغرب الثاني والمؤشر المشترك المنسي والمؤشر اليمني المشترك لسلسلة البنزينويد سيركوم كورونين. بالإضافة إلى ذلك، قدم البحث مؤشرات تبولوجية ومؤشرات تبولوجية مشتركة جديدة مثل مؤشر غزة ومؤشر القدس ومؤشر فلسطين ومؤشراتها المشتركة وصيغها الرياضية لسلسلة البنزينويد سيركوم كورونين. علاوة على ذلك، تم بناء بعض الخوارزميات باستخدام برامج البايثون لتنفيذ الصيغ الرياضية المشتقة بشكل عام.

الكلمات المفتاحية: سلسلة البنزينويد سيركوم كورونين، نظرية الرسم البياني الكيميائي، الرسوم البيانية الجزيئية، المؤشرات التبولوجية المشتركة، المؤشرات التبولوجية.