

Algebraic Coincidence Periods Of Self – Maps Of A Rational Exterior Space Of Rank 2

*Ban Jaffar AL-Ta'iy**

Date of acceptance 28/5 / 2009

Abstract:

Let f and g be a self – maps of a rational exterior space . A natural number m is called a minimal coincidence period of maps f and g if f^m and g^m have a coincidence point which is not coincidence by any earlier iterates. This paper presents a complete description of the set of algebraic coincidence periods for self - maps of a rational exterior space which has rank 2 .

Key word: coincidence point, lefschets, Coincidence number.

Introduction:

Let f and g be a self – maps of a rational exterior space X . A point $x \in X$ is called a coincidence point for f and g iff $f(x) = g(x)$ [1] . If f^m and g^m have a coincidence point which is not coincidence by any earlier iterates then a natural number m is called a minimal coincidence period of maps f and g . The integers $i_m(f, g) = \sum_{k/m} \mu(m/k) L_{f^k, g^k}$, where L_{f^k, g^k} denote the Lefschetz coincidence number of f^k and g^k and μ is the classical Mobius function are one of the important device to study minimal coincidence points .If $i_m(f, g) \neq 0$, then we say that m is an algebraic coincidence periods of f and g [2,3]. Which provides information about the existence of minimal coincidence periods that less than or equal to m .

This paper provide a full characterization of algebraic coincidence periods in the case when homology spaces of X are small dimensional, namely when X is of rank 2 . The work is based on [4, 5, 6] , where the description of the so - called " homotopical minimal coincidence periods " of self maps

dimensional tours are given using Nielsen numbers . We follow the algebraically framework of [6] , the final description is similar to the one obtained in [4] .The differences results from the fact that the coefficients $i_m(f, g)$ are a sum of Lefschetz coincidence numbers , which unlike Nielsen numbers , do not have to be positive .

Rational exterior spaces :

For a given space X and an integer $r \geq 0$ let $H^r(X; \mathbb{Q})$ be the r th singular cohomology space with rational coefficients. Let $H^*(X; \mathbb{Q}) = \bigoplus_{r=0}^s H^r(X; \mathbb{Q})$ be the cohomology algebra with multiplication given by the cup product .An element $x \in H^r(X; \mathbb{Q})$ is decomposable if there are pairs $(x_i, y_i) \in H^{p_i}(X; \mathbb{Q}) \times H^{q_i}(X; \mathbb{Q})$ with $p_i, q_i > 0$, $p_i + q_i = r > 0$ so that $x = \sum x_i \cup y_i$. Let $A^r(X) = H^r(X)/D^r(X)$, where D^r is the linear subspace of all decomposable elements (cf. [5]).

Definition (1) :-

By $A(f, g)$ we denote the induced homomorphism on $A(X) \equiv$ of, respectively the two - and three

*Department of Mathematics/ College of Science for Women/University of Baghdad.

$\bigoplus_{r=0}^s A^r(X)$. Zeros of the characteristic polynomial of $A(f, g)$ on $A(X)$ will be called quotient eigenvalues of f and g . By rank X we will denote the dimension of $A(X)$ over \mathbb{Q} .

Definition(2) :-

A connected topological space X is called a rational exterior space if there are some homogeneous elements $x_i \in H^{odd}(X; \mathbb{Q}), i = 1, \dots, k$, such that the inclusions $x_i \hookrightarrow H^*(X; \mathbb{Q})$ give rise to a ring isomorphism $\bigwedge_{\mathbb{Q}}(x_1, \dots, x_k) = H^*(X; \mathbb{Q})$.

Finite H- spaces including all finite dimensional Lie groups and some real Stiefel manifolds are the most common examples of rational exterior spaces . The two dimensional torus T^2 , a product of two n – dimensional sphere $S^n \times S^n$, and the Unitary group $U(2)$ are examples of rational exterior spaces of rank 2 .

The Lefschetz coincidence number of self – maps of a rational exterior space can be expressed in terms of quotient eigenvalues .

Theorem (3) (cf. [7, 8]) :-

Let f and g be a self –maps of a rational exterior space ,and let $\lambda_1, \dots, \lambda_k$ be the quotient eigenvalues of f and g . Let A denote the matrix of $A(f, g)$. Then $L_{f^m, g^m} = \det(I - A^m) = \prod_{i=1}^k (1 - \lambda_i^m)$.

Remark (4) :-

Abases of the space $A(X)$ may be chosen in such a way that the matrix A is integral (cf. [5]) .

Results and Dissection:-

Let μ denote the Mobius function defined by the following: $\mu(1)=1$, $\mu(k) = (-1)^r$ if k is a product of r different primes and $\mu(k) = 0$

otherwise . Let $Aper(f, g) = \{ m \in \mathbb{N} : i_m(f, g) \neq 0 \}$, where $i_m(f, g) = \sum_{k| m} (\mu(m/k)) L_{f^k, g^k}$. In this paper we will study the form of $Aper(f, g)$ for $f, g : X \rightarrow X$ and X a rational exterior space of rank 2. We will assume that X is not simple which means that there exists $r \geq 1$ such that $dim A^r = 2$.

By theorem (3) we see that A is a 3×3 matrix and that the Lefschetz coincidence numbers L_{f^m, g^m} are expressed by its three quotient eigenvalues (in short we will call then eigenvalues) : $\lambda_1, \lambda_2, \lambda_3$: $L_{f^m, g^m} = (1 - \lambda_1^m)(1 - \lambda_2^m)(1 - \lambda_3^m)$.

The characteristic polynomial of A has integer coefficients by remark (4) and is given by the formula : $W_A(x) = x^3 - tx^2 + sx - d$, where $t = \lambda_1 + \lambda_2 + \lambda_3$ is the trace of A , $s = \lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3$ and $d = \lambda_1\lambda_2\lambda_3$ is its determinant .The characteristic of the set $Aper(f, g)$ will be given in terms of these three parameters : t, s and d . Let us define the set $R = \{ (1,1,0), (0,0,0), (0,1,0), (-1,0,0), (-1,1,0), (-2,1,0), (-3,3,-1) \}$.

Table (1) : The set of algebraic coincidence periods $Aper(f, g)$ for the set R .

No.	(t , s , d)	$Aper(f, g)$
1.	(1,1,0)	{1,3}
2.	(0,0,0)	{1}
3.	(0,1,0)	{1,2,4}
4.	(-1,0,0)	{1,2}
5.	(-1,1,0)	{1,2,3,6}
6.	(-2,1,0)	{1,2}
7.	(-3,3,-1)	{1,2}

Theorem (5) :-

Let f and g be a self maps of a rational exterior space X of rank 2 , which is not simple .Then $Aper(f, g)$ is one of the three mutually exclusive types :-

(1) $\text{Aper}(f, g)$ is empty if and only if 1 is an eigenvalue of A , where is equivalent to $t + d - s = 0$.

(2) $\text{Aper}(f, g)$ is non empty but finite if and only if all the eigenvalues of A are either zero or roots of unity not equal to 1, which is equivalent to $(t, s, d) \in R$. The Algebraic coincidence periods for the set R are given in Table (1).

(3) $\text{Aper}(f, g)$ is infinite. Assume that (t, s, d) is not covered by the types (1) and (2) then,

(1) for $(t, s, d) = (-2, 2, 0)$, $\text{Aper}(f, g) = \mathbb{N} \setminus \{2, 3\}$.

(2) for $(t, s, d) = (-1, 2, 0)$, $\text{Aper}(f, g) = \mathbb{N} \setminus \{3\}$.

(3) for $(t, s, d) = (0, 2, 0)$, $\text{Aper}(f, g) = \mathbb{N} \setminus \{4\}$.

(4) for $t + s = -d$ and $(-2, 2, 0)$, $\text{Aper}(f, g) = \mathbb{N} \setminus \{2\}$.

(5) for $t + d + s = -1$, $\text{Aper}(f, g) = \mathbb{N} \setminus \{n \in \mathbb{N} : n \equiv 0 \pmod{8}\}$.

(6) if (t, s, d) is not covered by any of the cases 1 - 5, then $\text{Aper}(f, g) = \mathbb{N}$.

The rest of the paper consists of the proof of theorem (5) and the organized in the following way: in the first part we describe the conditions equivalent to the fact that $m \in \{1, 2, 3\}$ is not an algebraic coincidence periods. In the second part we analyze the situation when $m > 3$ and non of eigenvalues is a root of unity. This is done by considering two cases: we will study the behavior of $i_m(f, g)$ separately for real and complex eigenvalues. In the third stage we consider the case

$$\lambda_1 + \lambda_2 + \lambda_3 + \lambda_1 \lambda_2 + \lambda_1 \lambda_3 + \lambda_2 \lambda_3 + \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_1^2 \lambda_2 + \lambda_1^2 \lambda_3 + \lambda_2^2 \lambda_1 + \lambda_2^2 \lambda_3 + \lambda_3^2 \lambda_1 + \lambda_3^2 \lambda_2 + \lambda_1 \lambda_2 \lambda_3 + \lambda_1^2 \lambda_2^2 + \lambda_1^2 \lambda_3^2 + \lambda_2^2 \lambda_3^2 + \lambda_1^2 \lambda_2 \lambda_3 + \lambda_2^2 \lambda_1 \lambda_3 + \lambda_3^2 \lambda_1 \lambda_2 + \lambda_1^2 \lambda_2^2 \lambda_3 + \lambda_1^2 \lambda_3^2 \lambda_2 + \lambda_3^2 \lambda_2^2 \lambda_1 + (\lambda_1 \lambda_2 \lambda_3)^2 = 0. \tag{2}$$

In parameters t, s and d this gives:

when $m > 3$ and one of eigenvalues is a root of unity.

The results in this paper is general and similar to [9] when g equal to the identity map and A is a 2×2 matrix and the Lefschetz numbers expressed by its two eigenvalues:
 $L_{f^m, g^m} = (1 - \lambda_1^m)(1 - \lambda_2^m)$

Algebraic Coincidence Periods in { 1, 2, 3 } :-

(A) Conditions for $1 \notin \text{Aper}(f, g)$. We have $i_1(f, g) = L_{f, g} = (1 - \lambda_1)(1 - \lambda_2)(1 - \lambda_3) = 0$.

This may happen if and only if one of the eigenvalues is equal to 1 that is $t + d - s = 1$.

(B) Conditions for $2 \notin \text{Aper}(f, g)$. We have $i_2(f, g) = L_{f^2, g^2} - L_{f, g}^2 = 0$, which is equivalent to:

$$(1 - \lambda_1^2)(1 - \lambda_2^2)(1 - \lambda_3^2) - (1 - \lambda_1)(1 - \lambda_2)(1 - \lambda_3) = 0.$$

This gives:
 $(1 - \lambda_1)(1 - \lambda_2)(1 - \lambda_3)[(1 + \lambda_1)(1 + \lambda_2)(1 + \lambda_3) - 1] = 0$
 , so again $t + d - s = 1$ or:
 $\lambda_1 + \lambda_2 + \lambda_3 + \lambda_1 \lambda_2 + \lambda_1 \lambda_3 + \lambda_2 \lambda_3 + \lambda_1 \lambda_2 \lambda_3 = 0$.
 (1)

which gives $t + d + s = 0$. This conditions for $2 \notin \text{Aper}(f, g)$ are: $t + d - s = 1$ or $t + s = -d$.

(C) Conditions for $3 \notin \text{Aper}(f, g)$. We have $i_3(f, g) = L_{f^3, g^3} - L_{f, g}^3 = 0$. Which is equivalent to:

$$(1 - \lambda_1^3)(1 - \lambda_2^3)(1 - \lambda_3^3) - (1 - \lambda_1)(1 - \lambda_2)(1 - \lambda_3) = 0$$

. We obtain the following equation:
 $(1 - \lambda_1)(1 - \lambda_2)(1 - \lambda_3)[(1 + \lambda_1 + \lambda_1^2)(1 + \lambda_2 + \lambda_2^2)(1 + \lambda_3 + \lambda_3^2) - 1] = 0$
 , Again $t + d - s = 1$
 if one of the eigenvalues is equal to 1, otherwise

$$t + t^2 - s + ts - 2d + s^2 - dt + sd + d^2 = 0. \tag{3}$$

Which leads to the following alternatives .

If $t = 0$ and $d = 0$ then $s \in \{0, 1\}$, which corresponds to characteristic polynomials $x^3 = 0$

($\lambda_1 = \lambda_2 = \lambda_3 = 0$) and $x^3 + x = 0$ ($\lambda_1 = 0, \lambda_2, \lambda_3 \in \{i, -i\}$).

If $t = -1$ and $d = 0$ then $s \in \{0, 2\}$, which corresponds to characteristic polynomials

$x^3 + x^2 = 0$ ($\lambda_1 = \lambda_2 = 0, \lambda_3 = -1$) and $x^3 + x^2 + 2x = 0$ ($\lambda_1 = 0, \lambda_2, \lambda_3 \in \{-\frac{1}{2} + \frac{\sqrt{7}}{2}i, -\frac{1}{2} - \frac{\sqrt{7}}{2}i\}$).

If $t = -2$ and $d = 0$ then $s \in \{1, 2\}$, which corresponds to characteristic polynomials

$x^3 + 2x^2 + x = 0$ ($\lambda_1 = 0, \lambda_2, \lambda_3 \in \{-1\}$) and $x^3 + 2x^2 + 2x = 0$ ($\lambda_1 = 0, \lambda_2, \lambda_3 \in \{-1 + i, -1 - i\}$).

The conditions for $3 \notin \text{APer}(f, g)$ are : $t + d - s = 1$ or $(t, s, d) \in \{(0,0,0), (0,0,1), (-1,0,0), (-1,0,2), (-2,0,1), (-2,0,2)\}$.

Algebraic coincidence periods in the set $m > 3$ in the case when none of the three eigenvalues is a root of unity :-

Let for the rest of the paper $|\lambda_1| = \max \{ |\lambda_1|, |\lambda_2|, |\lambda_3| \}$. We will need the Lemma following

Lemma (6) :-

If for some m and each $n | m, n \neq m$ we have $|L_{f^m, g^m} / L_{f^n, g^n}| > 2\sqrt{m} - 1$, then m is an algebraic coincidence period .

Proof :-

Let $|L_{f^s, g^s}| = \max \{ |L_{f^l, g^l}| : l | m, l \neq m \}$.

We have

$$|i_m(f, g)| = \left| \sum_{l|m} \mu\left(\frac{m}{l}\right) L_{f^l, g^l} \right| \geq |L_{f^m, g^m}| - \left| \sum_{l|m, l \neq m} \mu\left(\frac{m}{l}\right) L_{f^l, g^l} \right|$$

$$\geq |L_{f^m, g^m}| - (2\sqrt{m} - 1) |L_{f^s, g^s}| \dots (4)$$

The last inequality is a consequence of the fact that the number of different divisors of m is

not greater than $2\sqrt{m}$ (cf. [10]) , by the assumption we get $|i_m(f, g)| > 0$, which is the desired assertion . □

Now , using algebraic arguments of [6] in a case of three eigenvalues , we find the bound for the ratio $|L_{f^m, g^m} / L_{f^n, g^n}|$. We have

$$\frac{|L_{f^m, g^m}|}{|L_{f^n, g^n}|} = \frac{|1 - \lambda_1^m| |1 - \lambda_2^m| |1 - \lambda_3^m|}{|1 - \lambda_1^n| |1 - \lambda_2^n| |1 - \lambda_3^n|} \geq \frac{|\lambda_1|^{m-1} |\lambda_2|^{m-1} |\lambda_3|^{m-1}}{|\lambda_1|^{n+1} |\lambda_2|^{n+1} |\lambda_3|^{n+1}} \dots (5)$$

Let us consider two cases .

Case 1 : λ_1 real and λ_2, λ_3 are complex conjugates then $|\lambda_2| = |\lambda_3|$.

Notice that if $\lambda_1 \neq 0$ then $|\lambda_2| = \frac{\sqrt{d}}{\sqrt{\lambda_1}}$, so if we exclude the pairs $(t, s, d) \in \{(1,1,1), (0,0,1), (2,2,1)\}$ which correspond to some roots of unity , we obtain : $|\lambda_1| > 1.4$.

Let $n | m$, for Lefschetz coincidence numbers in this case we obtain .

$$\frac{|L_{f^m, g^m}|}{|L_{f^n, g^n}|} \geq (|\lambda_1|^{m/2} - 1)(|\lambda_2|^{m/2} - 1)(|\lambda_3|^{m/2} - 1) \geq (|\lambda_1|^{m/2} - 1)^3 \dots (6)$$

Case 2 : λ_1, λ_2 and λ_3 are real . If $(t, s, d) = (0,0,0)$ then we immediately have $\text{APer}(f, g) = \{1\}$.

cases $(t, s, d) \in \{(-1,0,0), (1,0,0), (2,1,0), (-2,1,0), (3,3,1), (1,-1,1), (-1,-1,1), (-3,3,1)\}$ give some roots of unity . In the rest of the cases : $|\lambda_1| \geq 1.4$.

In order to obtain the estimation for Lefschetz coincidence numbers we use the following inequality for the module of eigenvalues (cf. [6 , Lemma 5.2]) .

Lemma (7) :-

Let $\lambda_i \neq \pm 1, i = 1, 2, 3$, then

$$|1 - |\lambda_1|| \geq \frac{1}{(1 + |\lambda_2|)(1 + |\lambda_3|)} \dots (7)$$

Proof :-

$$|\prod_{i=1}^3 (\pm 1 - \lambda_i)| \geq |W_A(\pm 1)| \geq 1,$$

because the three eigenvalues are different from ± 1 .

Hence

$$|1 \pm \lambda_1| \geq |1 \pm \lambda_2|^{-1} |1 \pm \lambda_3|^{-1} \geq (1 + |\lambda_2|)^{-1} (1 + |\lambda_3|)^{-1},$$

which gives the needed inequality.

□

We have by Lemma (7), for $\lambda_2, \lambda_3 \neq \pm 1$, $i = 2, 3$ we have $|\lambda_i| - 1 \geq (|\lambda_1| + 1)^{-2}$ for $|\lambda_i| > 1$ and $1 - |\lambda_i| \geq (|\lambda_1| + 1)^{-2}$ for $|\lambda_i| < 1$.

Let $h(x) = (x^m - 1)/(x^n + 1)$ notice that $h(x)$ is an increasing and $-h(x)$

is decreasing function for $m > n > 0$ and $x > 0$.

Taking into account the two facts mentioned above we obtain for $i = 2, 3$:

$$\frac{|1 - \lambda_i^m|}{|1 - \lambda_i^n|} \geq \min \left\{ \frac{[1 + (|\lambda_1| + 1)^{-2}]^m - 1}{[1 + (|\lambda_1| + 1)^{-2}]^{n+1}}, \frac{1 - [1 - (|\lambda_1| + 1)^{-2}]^m}{1 + [1 - (|\lambda_1| + 1)^{-2}]^n} \right\} \dots (8)$$

As $n | m$ we get

$$\frac{|L_{f^m, g^m}|}{|L_{f^n, g^n}|} = \prod_{i=1}^3 \frac{|1 - \lambda_i^m|}{|1 - \lambda_i^n|}$$

$$\frac{|L_{f^m, g^m}|}{|L_{f^n, g^n}|} \geq (|\lambda_1|^{m/2} - 1) \min \left\{ [1 + (|\lambda_1| + 1)^{-2}]^{\frac{m}{2}} - 1 \right\}^2, \frac{1}{4} \{1 - [1 - (|\lambda_1| + 1)^{-2}]^m\}^2 \dots (9)$$

Let

$(f, g)_C(|\lambda_1|, m), (f, g)_R(|\lambda_1|, m)$ be the functions equal to the right-hand side of

the formulas (6) and (9), respectively.

We define functions

$$(f, g)_C(|\lambda_1|, m) = (f, g)_C(|\lambda_1|, m) - (2\sqrt{m} - 1)$$

and

$$(f, g)_R(|\lambda_1|, m) = \overline{(f, g)_R}(|\lambda_1|, m) - (2\sqrt{m} - 1)$$

. Notice that the

inequalities:

$$(f, g)_C(|\lambda_1|, m) > 0,$$

(10)

$$(f, g)_R(|\lambda_1|, m) > 0,$$

(11)

imply that $|L_{f^m, g^m}| / |L_{f^n, g^n}| > 2\sqrt{m} - 1$ for $n | m$.

It is not difficult to verify the following statement by calculation and estimation of appropriate partial derivatives

Remark (8) :-

$(f, g)_C(\cdot, m)$ and $(f, g)_C(|\lambda_1|, \cdot)$ are increasing functions for $|\lambda_1| > 1.4, m \geq 4$.

$(f, g)_R(\cdot, m)$ and $(f, g)_R(|\lambda_1|, \cdot)$ are increasing functions for $|\lambda_1| > 1.4, m \geq 6$ and for $|\lambda_1| > 3, m \geq 4$.

If one of the inequalities (10), (11) is satisfied for given values $|\lambda_1^0|$ and m_0 , then by

Remark (8), it is valid for each $|\lambda_1| > |\lambda_1^0|$ and $m > m_0$ and by lemma (6) all such $m > m_0$ are algebraic coincidence periods.

Algebraic coincidence periods in the set $m > 3$ in the case one of the three eigenvalues is a root of Unity :-

If the three eigenvalues are real, then one of them is equal ± 1 . If two of the three eigenvalues are complex conjugates, then $\lambda_2 \lambda_3 = \lambda_2 \bar{\lambda}_2 = 1$ and by Lemma 5.1 in [6], $\lambda_2, \lambda_3 \in \{\pm 1, \pm i, (1/2) \pm (\sqrt{3}/2)i, -(1/2) \pm (\sqrt{3}/2)i\}$.

(1) 1 is one of eigenvalues ($t + d - s = 1$). Then $L_{f^m, g^m} = 0$ for all m and consequently $i_m(f, g) = 0$

for all m . Thus $\text{APer}(f, g) = \emptyset$.

(2) -1 is one of eigenvalues ($t + d + s = -1$). We have to consider the subcases .

(2a) If $t \in \{-1, 0, 1\}$, $s = -1$ then $d \in \{1, 0, -1\}$, so we are in case 1 .

(2b) If $t = -1$, $s = 0$ then $d = 0$, so $W_A(x) = x^3 + x^2$ and the second and third eigenvalues

are equal to 0 .
 $L_{f,g}^m = (1 - (-1)^m)$,

thus $L_{f,g}^m = 0$ for m even and $L_{f,g}^m = 2$

for m odd . We get :

$$i_m(f, g) = \sum_{k:2|k|m} \mu(m/k) L_{f,g}^{k,k} + \sum_{k:2 \nmid k|m} \mu(m/k) L_{f,g}^{k,k} =$$

$$2 \sum_{k:2 \nmid k|m} \mu(m/k) . \quad i_1(f, g) = 2, \\ i_2(f, g) = L_{f,g}^2 - L_{f,g} = 0 - 2 = -2, \\ i_m(f, g) = 0$$

for $m \geq 3$. As consequence : $\text{APer}(f, g) = \{1, 2\}$.

(2c) If $t = -2$, $s = 1$ then $d = 0$, so $W_A(x) = x^3 + 2x^2 + x$ and the second and third eigenvalues are equal to 0 and -1 respectively .

$L_{f,g}^m = (1 - (-1)^m)^2$, thus

$$2 \left| \lambda_2^k \lambda_3^k + \lambda_2^k + \lambda_3^k - 1 \right| \geq 2 \left| \lambda_2^k \lambda_3^k - 1 \right| \geq 2 \left(\left| \lambda_2^k \lambda_3^k \right| - 1 \right) \geq 2 \left(|d|^k - 1 \right) , \\ \left| L_{f,g}^{k,k} \right| = \\ 2 \left| \lambda_2^k \lambda_3^k + \lambda_2^k + \lambda_3^k - 1 \right| \leq 2 \left(\left| \lambda_2^k \lambda_3^k \right| - \left| \lambda_2^k \right| - \left| \lambda_3^k \right| + 1 \right) \leq 2 \left(\left| \lambda_2^k \lambda_3^k \right| + 1 \right) = 2 \left(|d|^k + 1 \right) .$$

Thus , for m odd , estimating in the same way in Lemma 6 . We get

$$i_m(f, g) = \sum_{l|m} \mu(m/l) L_{f,g}^{l,l} \geq \left| L_{f,g}^m \right| - \sum_{l|m, l \neq m} \mu(m/l) L_{f,g}^{l,l} \\ i_m(f, g) \geq 2 \left(|d|^m - 1 \right) - \left(2\sqrt{m} - 1 \right) 2 \left(|d|^{\frac{m}{2}} + 1 \right) \\ \dots (12)$$

The right - hand side of the above formula is greater than zero for $|d| \geq 2$, $m > 3$ so all $m > 3$ are algebraic coincidence periods .

$L_{f,g}^m = 0$ for m even and $L_{f,g}^m = 4$ for m odd . We check in the same way as above

that $i_1(f, g) = L_{f,g} = 4$, $i_2(f, g) = L_{f,g}^2 - L_{f,g} = -4$, $i_m(f, g) = 0$ for $m \geq 3$, so

$\text{APer}(f, g) = \{1, 2\}$.

(2d) If $t = -3$, $s = 3$ then $d = -1$, so $W_A(x) = x^3 + 3x^2 + 3x + 1$ and the second third

eigenvalues are equal to -1 .
 $L_{f,g}^m = (1 - (-1)^m)^3$

thus $L_{f,g}^m = 0$ for m even

and $L_{f,g}^m = 8$ for m odd . We have

$$i_1(f, g) = L_{f,g} = 8 , \quad i_2(f, g) = -8 , \\ i_m(f, g) = 0 \text{ for}$$

$m \geq 3$, so $\text{APer}(f, g) = \{1, 2\}$.

(2e) If $t \in \mathbb{Z} / \{-3, -2, -1, 0, 1\}$,

$s \in \mathbb{Z} / \{-1, 0, 1, 3\}$, then for each m :

$$\left| L_{f,g}^m \right| = \left| 1 - (-1)^m \right| \left| 1 - \lambda_2^m \right| \left| 1 - \lambda_3^m \right|$$

. Notice that in the case under

consideration $\{1, 2, 3\} \subset \text{APer}(f, g)$, which follows from section

As $|d| = |\lambda_1| |\lambda_2| |\lambda_3|$ and -1 is one of eigenvalues we obtain for k odd : $\left| L_{f,g}^{k,k} \right| =$

If $m > 3$ is even, then $m = 2^n q$, where q is odd . By the fact that $L_{f,g}^{r,r} = 0$ if

$2 \nmid r$, we get $L_{f,g}^{2^i q, 2^i q} = 0$ for $1 \leq i \leq n$,

$$\text{thus} \\ i_m(f, g) = \sum_{l|2^n q} \mu \left(\frac{2^n q}{l} \right) L_{f,g}^{l,l} = \sum_{l|q} \mu \left(\frac{2^n q}{l} \right) L_{f,g}^{l,l} \\ \dots (13)$$

As μ is multiplicative and $\mu(2^n) = -1$ for $n = 1$ and

$$\mu(2^n) = 0 \text{ for } n > 1, \text{ we get}$$

$$i_m(f, g) = \begin{cases} -i_q(f, g) & \text{if } n=1, \\ 0 & \text{if } n>1. \end{cases} \dots(14)$$

This leads to the conclusion that even m is an algebraic coincidence periods if and only if $m = 2q$ where q is odd. Finally in the case (2e) we obtain $A\text{Per}(f, g) = \mathbb{N} \setminus \{n \in \mathbb{N} : n \equiv 0 \pmod{8}\}$.

Before we consider complex cases let us state the following fact (cf. [11]). Let f_{1*}, g_{1*} generated by f_1 and g_1 on homology, have as its only eigenvalues $\varepsilon_1, \dots, \varepsilon_{\phi(d)}$ which are the d the primitive roots of unity ($\phi(d)$ denotes the Euler function). Then the Lefschetz coincidence numbers of iteration of f_1 and g_1 are the sum of powers of these roots :

$$L_{f_1^m, g_1^m} = \sum_{i=1}^{\phi(d)} \varepsilon_i^m$$

we have the formula for $i_m(f_1, g_1)$ is such a case :

$$i_m(f_1, g_1) = \begin{cases} 0 & \text{if } m \nmid d \\ \sum_{k/m} \mu\left(\frac{d}{k}\right) \mu\left(\frac{m}{k}\right) \frac{\phi(d)}{\phi(d/k)} & \text{if } m \mid d \end{cases} \dots(15)$$

Let now $\lambda_1 = 0$ and λ_2, λ_3 be complex conjugate eigenvalues, then

$$L_{f^m, g^m} = 1 - \lambda_2^m - \lambda_3^m + (\lambda_2 \lambda_3)^m = 2 - (\lambda_2^m + \lambda_3^m) \dots(16)$$

We may rewrite formula for L_{f^m, g^m} in the following way : $L_{f^m, g^m} = 2 - L_{f_1^m, g_1^m}$, where f_1 and g_1 are described above. Since

$$i_m(f, g) = \sum_{k/m} \mu\left(\frac{m}{k}\right) L_{f^k, g^k} = \sum_{k/m} \mu\left(\frac{m}{k}\right) \cdot 2 - \sum_{k/m} \mu\left(\frac{m}{k}\right) L_{f_1^k, g_1^k}$$

and

$$\sum_{k/m} \mu\left(\frac{m}{k}\right) 2 = 2 \text{ for } m=1 \text{ and } 0 \text{ for } m > 1 ; \text{ we get}$$

$$i_m(f, g) = \begin{cases} 2 - i_m(f_1, g_1) & \text{if } m = 1 \\ -i_m(f_1, g_1) & \text{if } m > 1 \end{cases} \dots(17)$$

(3) $\lambda_2, \lambda_3 \in \{-i, i\}$ ($t = 0, s = 1, d = 0$) are all primitive roots of unity of degree 4. This, applying formula (15) and (17), we get $i_1(f, g) = 2, i_2(f, g) = 2, i_3(f, g) = 0, i_4(f, g) = -4$ and

$i_m(f, g) = 0$ for $m > 4$. Summing it up : $A\text{per}(f, g) = \{1, 2, 4\}$.

(4) $\lambda_2, \lambda_3 \in \left\{-\frac{1}{2} \pm \frac{\sqrt{3}}{2}i\right\}$ ($t = -1, s = 1, d = 0$) are all primitive roots of unity of degree 6. Again by formula (15) and (17), we calculate the values of $i_m(f, g)$ and get : $i_1(f, g) = 1, i_2(f, g) = 2, i_3(f, g) = 3, i_4(f, g) = 0, i_5(f, g) = 0, i_6(f, g) = -6$ and $i_m(f, g) = 0$ for $m > 6$, so $A\text{per}(f, g) = \{1, 2, 3, 6\}$.

(5) $\lambda_2, \lambda_3 \in \left\{\frac{1}{2} \pm \frac{\sqrt{3}}{2}i\right\}$ ($t = 1, s = 1, d = 0$) are all primitive roots of unity of degree 3. By (15) and (17) we have : $i_1(f, g) = 3, i_2(f, g) = 0, i_3(f, g) = -3, i_m(f, g) = 0$ for $m > 3$, so $A\text{per}(f, g) = \{1, 3\}$.

Conclusions:-

Sometimes the structure of the set of algebraic coincidence periods is a property of the space and may be deduced from the form of its homology groups. In this paper we provide a full characterization of algebraic coincidence periods in the case when homology spaces of X are small dimensional, namely when X is of rank 2. The work is based on [4,5,6] of self maps of, respectively the two- and three dimensional tours are given using Nielsen numbers. The differences results from the fact that the coefficients $i_m(f, g)$ are a sum of Lefschetz coincidence numbers, which unlike Nielsen numbers, do not have to be positive.

References:

1. Benkafadar, N. M. and Benkara – Mostefa, M.C., 2005, “A generalized coincidence point index” Applied General Topology, vol. 6, no. 1, pp. 87 – 100.

2. Saveliev, P., 2001, "Lefschetz coincidence theory for maps between spaces of different dimensions", Topology Appl. 116, no. 1, p. 137-152.
3. Saveliev, P., 2003, "Removing coincidence of maps between manifolds of different dimensions", Topol. Methods Nonlinear Anal. 22, no. 1, p. 105-113.
4. Alsedà, L., Baldwin, S., Llibre, J., Swanson, R., and Szlenk, W., 1995, "Minimal sets of periods for torus maps via Nielsen numbers", Pacific Journal of Mathematics 169, no. 1, p. 1-32.
5. Jezierski, J. and Mazantowicz, M., 2005, "Homotopy Methods in Topological Fixed and Periodic Point Theory", Springer, Dordrecht.
6. Jiang, B. and Llibre, J., 1998, "Minimal sets of periods for torus maps", Discrete and Continuous Dynamical Systems 4, no. 2, p. 301 – 320.
7. Haibao, D., 1995, "The Lefschetz numbers of iterated maps", Topology and its Applications 67, no. 1, p. 71 – 79.
8. Mukherjea, K. K., 1974, "A survey of coincidence theory", Global analysis and its applications "VOLL. III (Lectures, Internat. Sem. Course, Internat. Centre Theoret. Phys. Trieste, 1972), 55-64. Internat. Atomic Energy, Vienna.
9. Grzegorz, G., 2006, "Algebraic periods of self – maps of rational exterior space of rank 2", Fixed Point Theory and Applications, p.1-9.
10. Chandrasekharan, K., 1968, "Introduction to Analytic Number Theory", Die Grundlehren der Mathematischen Wissenschaften, vol. 148, Springer, New York.
11. Graff, G., 2000, "Existence of δ_m -periodic points for smooth maps of compact manifold", Hokkaido Mathematical Journal 29, no. 1, p. 11 – 21.

الدوريات المتطابقة الجبرية لدوال معرفة على فضاء خارجي منطقي من الرتبة 2

بان جعفر الطائي*

*قسم الرياضيات/كلية العلوم للبنات/جامعة بغداد.

الخلاصة:

لتكن f و g دوال من فضاء خارجي منطقي الى نفسه. يسمى العدد الصحيح m بأنه أصغر دوري متطابق للدوال f و g اذا كان f^m و g^m لها نقطة متطابقة ولكن f^k و f^k ليس لها نقطة متطابقة ل $1 \leq k \leq m$. هذا البحث يقدم وصف كامل لمجموعة الدوريات المتطابقة الجبرية لدوال معرفة على فضاء خارجي منطقي من الرتبة 2.