

Spexin is a Reliable Predictor of Diabetes Postmenopausal Iraqi Women: A Case-Control Study

Mustafa.s. shareef¹  , Ali B. Roomi^{*2}  , Bassem Charfeddine³  

¹ Department of Biochemistry, College of Health and Medical Techniques, AL-Bayan University, Baghdad, Iraq.

² Department of Chemistry, College of Science, University of Thi-Qar, Thi-Qar, 64001, Iraq.

³ Department of Biochemistry, Faculty of Medicine, University of Sousse, Tunisia.

*Corresponding Author.

Received 27/03/2024, Revised 27/07/2024, Accepted 29/07/2024, Published Online First 20/12/2024



© 2022 The Author(s). Published by College of Science for Women, University of Baghdad.

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

A peptide called spexin (SPX) regulates blood glucose and lipid metabolism. The study primarily focused on the pathogenesis of type 2 diabetes mellitus (T2DM) in postmenopausal women, examining its relationship with serum SPX levels. A case-control study involved 180 postmenopausal women from Thi-Qar, Iraq were enrolled in the current study. Ninety women were recently diagnosed with diabetic postmenopausal women, while the other 90 were non-diabetic postmenopausal women. Blood samples were collected to test Glycated hemoglobin (HbA1c), SPX, Malondialdehyde (MDA), Superoxide dismutase (SOD), C-reactive protein (CRP), fasting blood glucose (FSG), insulin, lipid profile and interleukin (IL-6). Results: FSG, HbA1c, insulin, and Homeostatic Model Assessment for Insulin Resistance (HOMA-IR), values raised significantly ($P < 0.001$) in T2DM postmenopausal women. Also, levels of MDA, CRP, and IL-6 dramatically elevated, and the levels of SPX and SOD significantly decreased ($P < 0.001$). Significant lipid profile changes were observed in postmenopausal T2DM subjects. Additionally, the connection in T2DM postmenopausal women showed a negative relationship between SPX with waist-to-height ratio, SPX with FSG, SPX with HOMA-IR, MDA with SPX, as well as a positive relationship between SPX and MDA, SPX and age, CRP and IL-6, CRP and age, IL-6 and FSG, SOD and insulin. It was demonstrated that SPX is crucial to the pathogenesis of T2DM in postmenopausal women. In diabetic individuals, targeting SPX may offer a novel strategy for preserving adequate glucose control.

Keywords: CRP, Malondialdehyde, Spexin, Superoxide dismutase, T2DM.

Introduction

Diabetes mellitus (DM) is a metabolic disease expressed as high blood sugar resulting from a dysfunction in insulin production, activity, or both. The eyes, kidneys, nerves, heart, and blood vessels are particularly vulnerable to long-term damage, dysfunction, and failure caused by chronic hyperglycemia, which is related to diabetes¹.

The onset of diabetes is caused by multiple pathological mechanisms. These can vary from autoimmune damage to the pancreatic β -cells leading

to insulin shortages to anomalies that cause insulin resistance². DM is characterized by dysfunctional glucose, lipid, and protein metabolism, caused by insulin not acting on the tissues that need it. Inadequate secretion of insulin and/or reduced tissue responses to insulin at one or more points in the intricate hormone action pathways cause the insulin deficiency impact³.

Many patients have problems with both insulin secretion and action simultaneously, and it is

frequently unclear which defect is the primary cause of hyperglycemia if one is alone. There are two types of diabetes symptoms: the first is marked, and the other is long-term⁴. Signs of severe hyperglycemia include the following: increased thirst, lack of appetite, hazy vision, and more frequent urine. Chronic hyperglycemia can lead to stunted growth and an increased risk of contracting certain illnesses⁵. Hyperglycemia with ketoacidosis, or ketoacidosis hyperosmolar syndrome, is one of the immediate and potentially fatal complications of uncontrolled diabetes⁶. Diabetes can cause retinopathy, which can lead to vision loss, and nephropathy, which can cause kidney failure⁷. Foot ulcers, amputations, and Charcot joints are all possible outcomes of peripheral neuropathy, while gastrointestinal, genitourinary, cardiovascular, and sexual dysfunction symptoms can be caused by autonomic neuropathy. Atherosclerosis of the coronary and peripheral

arteries, as well as cerebrovascular disorders, are more common in diabetic patients. Diabetic individuals frequently exhibit hypertension and lipoprotein metabolism problems⁷.

Type 2 diabetes, termed noninsulin-dependent or adult-onset diabetes mellitus, is often associated with factors such as obesity, a sedentary lifestyle, a family history of diabetes, and increasing age. Management of Type 2 diabetes can often involve dietary interventions or the use of oral hypoglycemic agents. This form of diabetes accounts for a variable prevalence, encompassing 90-95% of all diabetes cases⁶. Research findings have indicated that reduced expression of the spexin gene and lower serum spexin concentrations are associated with obesity, T2DM, hypertension, and cardiovascular disease, particularly in the context of metabolic syndrome conditions⁷.

Materials and Methods

Study population and samples collection

The study was approved by the Al-Hussein Teaching Hospital's Ethical Committee (IRB number: 01/2023).

The blood samples were collected from 180 postmenopausal women from Thi-Qar, Iraq; 90 of them visited the Thi-Qar Governorate's diabetes and endocrinology centers with T2DM disease, while the other 90 samples were collected from non-diabetic postmenopausal women who visited the governorate's Al-Hussein Teaching Hospital. The samples were collected between February 28 and April 17, 2023. The name, age, weight, height, waist circumference, and family history at the time of study of each woman were recorded.

Samples preparation

Two blood samples were taken; the first was taken in test tubes containing EDTA, and the second was taken in test tubes without EDTA and allowed to coagulate for at least fifteen minutes at room temperature. After that, the samples were centrifuged for ten minutes at 3600 rpm to separate the serum, which was then stored at -20 °C until it was needed for the biochemical analysis.

The whole blood was used to analyze glycated hemoglobin (HbA1c) (Catalogue number; 06378676190), while the serum was used for the analyses of SPX (Catalogue number; orb1140635), Malondialdehyde (MDA) (Catalogue number;

EEA015) Superoxide dismutase (SOD) (Catalogue number; 15,665,069), C-reactive protein (CRP) (Catalogue number C0128), fasting blood glucose (FS (Catalogue number; 11351-05051), insulin (Catalogue number; 15140122) , lipid profile, and interleukin 6 (IL-6) (Catalogue number; 341-36). The levels of FSG and insulin were used to evaluate the Homeostatic Model Assessment for Insulin Resistance (HOMA-IR) level.

Included criteria

Women with newly diagnosed diabetes were diagnosed by the doctors of the Thi-Qar Governorate's diabetes and endocrinology centers. This criterion was based on the causes of T2DM in the menopausal stage, which were frequent urination, obesity, autoimmunity, and insulin resistance.

Statistical analysis

The findings were expressed as mean \pm SD by using BM's SPSS software version 24, and the difference between more than two groups was ascertained T-test. Bonferroni's analysis was employed. The parameters were correlated using the Pearson correlation (r) coefficient. SPX parameter's diagnostic viability was determined using ROC curve analysis or receiver operating characteristic curve analysis. Statistics were significant at $p < 0.001$.

Results and Discussion

The anthropometric indications of healthy and postmenopausal women with T2DM are shown in Table 1. The mean age of the postmenopausal participants was in the mid-fifties, showing no significant difference ($P < 0.001$) between the control group (54.97 ± 4.31 years) and those with T2DM (56.21 ± 4.37 years). The waist circumferences of healthy postmenopausal women (78.98 ± 5.21 cm) and postmenopausal women with T2DM (80.70 ± 5.77 cm) were statistically significant ($P < 0.001$), and the patients showed a higher waist circumference than the control. The differences in BMI were non-significant ($P < 0.001$) between postmenopausal healthy women (25.70 ± 1.63 kg/m²) and postmenopausal women with T2DM (26.11 ± 1.64 kg/m²). Postmenopausal women with T2DM (0.49 ± 0.03) showed significant ($P < 0.001$)

differences in the waist-to-height ratio (WHtR) compared to postmenopausal healthy women (0.48 ± 0.03). Age is a crucial risk factor for many health disorders, including T2DM disease. Since this study was conducted on postmenopausal women, the age of the participants was above 45, and it was restricted upon the selection of the study participants to maintain the differences between control and patients in the non-significant region. So, the age was eliminated from the fluctuations of the studied parameters. Moreover, the BMI was treated similarly to avoid any influence of obesity or overweight on the levels of the studied parameters. Yet, there was a significant difference in the values of waist circumference and WHtR between the control and patients^{8,9}.

Table 1. The basic anthropometric criteria of the study subgroups expressed as means \pm SD.

Parameter	Postmenopausal women (N=90)	healthy	Postmenopausal women with newly T2DM diagnosis (N=90)	P-value
Age (year)	54.97 \pm 4.31		56.21 \pm 4.37	0.056
Waist circumference (cm)	78.98 \pm 5.21		80.70 \pm 5.77	0.037
BMI (kg/m ²)	25.70 \pm 1.63		26.11 \pm 1.64	0.096
WHtR	0.48 \pm 0.03		0.49 \pm 0.03	0.030

The level of IL-6 was elevated significantly ($P < 0.01$) in postmenopausal women with newly diagnosed T2DM (13.45 ± 4.37 pg/mL) compared to postmenopausal healthy women (6.21 ± 2.13 pg/mL), as shown in Table 2. Stanimirovic et al. reported a significant increase in CRP levels in T2DM patients. They attributed this increase to the elevation of the overall inflammation occurring due to insulin resistance in T2DM patients⁹. Previous studies showed that even slight increases in the inflammatory marker CRP are indicative of a higher likelihood of experiencing cardiovascular events in both diabetic and non-diabetic individuals¹⁰.

Furthermore, in individuals without prior health issues, T2DM risk raise in those with elevated C-reactive protein levels. Previous research suggested that CRP not only serves as a predictor for cardiovascular risk but may also play an active role in atherogenesis^{10,11}. Sadeghabadi et al. observed a notable elevation in IL-6 levels in the serum of individuals with T2DM. They found a positive correlation between IL-6 levels and factors such as BMI, FSG, and HOMA-IR in these patients. This increase in IL-6 was attributed to the presence of insulin resistance among individuals with T2DM^{12,13}.

Table 2. The levels of inflammatory indicators in the study participants.

Parameter	Postmenopausal women (N=90)	healthy	Postmenopausal women with newly T2DM diagnosis (N=90)	P-value
CRP (mg/L)	7.07 \pm 2.30		14.12 \pm 6.71	<0.001*
IL-6 (pg/dL)	6.21 \pm 2.13		13.45 \pm 4.37	<0.001*

The level of MDA elevated significantly ($P < 0.01$) in postmenopausal women with newly diagnosed T2DM (6.29 ± 2.02 μ mol/L) compared to postmenopausal healthy women (1.72 ± 0.84 μ mol/L). The level of SOD reduced significantly ($P < 0.05$) in

postmenopausal women with newly diagnosed T2DM (4.08 ± 1.90 IU/mL) compared to postmenopausal healthy women (6.47 ± 1.96 IU/mL), as shown in Table 3.

Table 3. The levels of oxidative stress indicators in the study participants.

Parameter	Postmenopausal women (N=90)	healthy	Postmenopausal women with newly T2DM diagnosis (N=90)	P-value
MDA ($\mu\text{mol/L}$)	1.72 \pm 0.84		6.29 \pm 2.02	<0.001*
SOD (IU/mL)	6.47 \pm 1.96		4.08 \pm 1.90	<0.001*

The results of lipid profile parameters, including TG, TC, HDL, LDL, and VLDL, are shown in Table 4. The level of TG significantly increased ($P<0.001$) in postmenopausal women with newly diagnosed T2DM (137.75 \pm 27.63 mg/dL) compared to postmenopausal healthy women (120.15 \pm 20.83 mg/dL). The level of TC non-significantly increased ($P<0.001$) in postmenopausal women with newly diagnosed T2DM (167.17 \pm 27.11 mg/dL) compared to postmenopausal healthy women (159.82 \pm 23.80 mg/dL). The level of HDL reduced significantly ($P<0.001$) in postmenopausal women with newly diagnosed T2DM (39.00 \pm 6.01 mg/dL) compared to postmenopausal healthy women (40.92 \pm 5.66 mg/dL). The level of LDL increased non-significantly ($P<0.001$) in postmenopausal women with newly diagnosed T2DM (100.62 \pm 25.90 mg/dL)

compared to postmenopausal healthy women (94.87 \pm 23.77 mg/dL). The level of VLDL significantly increased ($P<0.001$) in postmenopausal women with newly diagnosed T2DM (27.55 \pm 5.53 mg/dL) compared with postmenopausal healthy women (24.03 \pm 4.17 mg/dL). The outcomes of lipid profile parameters, including TG, TC, HDL, LDL, and VLDL, are shown in Table 4.

The level of TG significantly increased ($P<0.001$) in postmenopausal women with newly diagnosed T2DM (137.75 \pm 27.63 mg/dL) compared with postmenopausal healthy women (120.15 \pm 20.83 mg/dL). Tejaswi et al. reported significant alteration in the levels of lipid profile parameters, where the levels of TG, TC, LDL, and VLDL elevated while the level of HDL decreased in T2DM patients^{14, 15}.

Table 4. The level of lipid profile parameters in the study participants.

Parameter	Postmenopausal women (N=90)	healthy	Postmenopausal women with newly T2DM diagnosis (N=90)	P-value
TG (mg/dL)	120.15 \pm 20.83		137.75 \pm 27.63	<0.001*
TC (mg/dL)	159.82 \pm 23.80		167.17 \pm 27.11	0.055
HDL (mg/dL)	40.92 \pm 5.66		39.00 \pm 6.01	0.029
LDL (mg/dL)	94.87 \pm 23.77		100.62 \pm 25.90	0.123
VLDL (mg/dL)	24.03 \pm 4.17		27.55 \pm 5.53	<0.001*

Postmenopausal women with newly diagnosed T2DM disease showed a significant positive correlation between CRP and IL-6 ($r= 0.330$, $p= 0.001$), CRP and age ($r= 0.304$, $p= 0.004$), IL-6 and FSG ($r= 0.229$, $p= 0.030$), as shown in Table 5. Postmenopausal women with newly diagnosed T2DM disease showed a significant positive

correlation between SPX and MDA ($r = 0.214$, $p= 0.042$), SPX and age ($r= 0.324$, $p= 0.002$), SPX and FSG ($r= -0.250$, $p= 0.018$), as shown in Table 5. Tejaswi et al. reported a significant reduction of SPX levels in T2DM patients. They indicated that SPX levels correlate with T2DM in patients¹⁴.

Table 5. Correlation among parameters in T2DM postmenopausal women.

Parameters	SPX		CRP		IL-6	
	R	P	R	P	R	p
CRP	0.173	0.103	-	-	0.330*	0.001
IL-6	-0.024	0.825	0.330*	0.001	-	-
MDA	0.214*	0.042	-0.054	0.613	-0.097	0.363
Age	0.324*	0.002	0.304*	0.004	0.075	0.485
FSG	-0.250*	0.018	-0.074	0.488	0.229*	0.030

The present study on postmenopausal women with Type 2 Diabetes Mellitus (T2DM) highlights the potential of interleukin-6 (IL-6) as a valuable biomarker for disease prognosis, with an excellent sensitivity of AUC=0.941 and a cut-off value of 8.45 pg/dL, providing 91.1% sensitivity and 80.1% specificity. Additionally, malondialdehyde (MDA) and superoxide dismutase (SOD) were also evaluated as biomarkers for T2DM prognosis in postmenopausal women, showing promising results. MDA exhibited excellent sensitivity with an AUC of 0.990, a cut-off value of 2.90 $\mu\text{mol/L}$, 97.8% sensitivity, and 90% specificity. On the other hand, SOD demonstrated good sensitivity with an AUC of 0.816, a cut-off value of 4.99 IU/mL, 80% sensitivity, and 72.2% specificity. These findings suggest the potential of IL-6, MDA, and SOD as prognostic biomarkers for T2DM in postmenopausal

women, with IL-6 showing particularly promising sensitivity and specificity levels, as shown in Table 6. and Figs. 1, 2 and 3. Chang et al. reported a significant alteration in the levels of IL-6, MDA and SOD in T2DM¹⁶. Both MDA and SOD are indicators of the redox state and oxidative stress, which arises from high levels of ROS. There are many pathways in which ROS levels are elevated in T2DM patients, most importantly high blood glucose levels and insulin resistance¹⁷. The electron transport system in the mitochondria is another route that contributes to oxidative stress. Adenosine triphosphate is a crucial organic energy source created in the mitochondrial inner membrane, where this system is found. Water molecules are formed in this electron transfer system by deoxidizing four oxygen molecule electrons; ROS is created as an intermediate product during this process, and some ROS escape from the system¹⁸.

Table 6. ROC analysis outcomes.

Parameters	AUC	Standard error	<i>p</i> -value	Cut-off value	Sensitivity	Specificity
IL-6 (pg/dL)	0.941	0.018	<0.001	8.45	91.1%	80.1%
MDA ($\mu\text{mol/L}$)	0.990	0.005	<0.001	2.90	97.8%	90%
SOD (IU/mL)	0.816	0.032	<0.001	4.99	80%	72.2%




Figure 1. The ROC curve of IL-6 in the prognosis of postmenopausal women with T2DM disease.




Figure 2. The ROC curve of MDA in the prognosis of postmenopausal women with T2DM disease.




Figure 3. The ROC curve of SOD in the prognosis of postmenopausal women with T2DM disease.

Conclusion

Spexin levels are reduced significantly in postmenopausal women with T2DM disease, which reflects a state of dysregulation of glucose homeostasis and lipid metabolism. Targeting SPX in T2DM patients may improve the glycemic state of

postmenopausal patients. Postmenopausal women with T2DM disease have shown a significant increase in MDA levels and a significant decrease in SOD, indicating a condition of oxidative stress.

Acknowledgment

The study was conducted at the Al-Hussein Teaching Hospital and diabetes and endocrinology centers in

Thi-Qar Governorate, the authors express their thanks to all who contributed to its construction.

Authors' Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for re-publication, which is attached to the manuscript.

- No animal studies are present in the manuscript.
- Authors sign on ethical consideration's approval.
- Ethical Clearance: The project was approved by the local ethical committee at University of Thi-Qar.

Authors' Contribution Statement

M. S. S, A. B. R and B. C. designed and conducted the research; A. B. R. and M .S. S performed the data management and statistical analyses; M .S. S and A. B. R wrote the manuscript. B. C., A. B. R and M. S.

S reviewed/edited the manuscript for important intellectual content. A. B. R, M. S. S and B. C. read and approved the final manuscript.

References

1. Iser BPM, Stein C, Alves LF, Carvalho MLS, Espinoza SAR, Schmidt MI. A portrait of gestational diabetes mellitus in Brazil: A systematic review and meta-analysis. *Arch Endocrinol Metab.* 2023 Oct 18; 67(6): e220521. <https://doi.org/10.20945/2359-4292-2022-0521>.
2. Pakyurek H, Aykota MR, Kilic-Erkek O, Ozban M, Senol H, Bor-Kucukatay M. Investigation of time-dependent alterations in adipokine levels and endoplasmic reticulum stress markers in obese patients with laparoscopic sleeve gastrectomy. *Life Sci.* 2023 Oct 1; 330: 121987. <https://doi.org/10.1016/j.lfs.2023.121987>
3. Kengne AP, Ramachandran A. Feasibility of prevention of type 2 diabetes in low- and middle-income countries. *Diabetologia.* 2024 May; 67(5): 763-772. <https://doi.org/10.1007/s00125-023-06085-1>

4. Sacks DB, Arnold M, Bakris GL, Bruns DE, Horvath AR, Kirkman MS, et al., Guidelines and recommendations for laboratory analysis in the diagnosis and management of diabetes mellitus. *Clin Chem.* 2011 Jun; 57(6): e1-e47. <https://doi.org/10.1373/clinchem.2010.161596>.
5. Çiftçi Y, Gurger M, Gul E, Yilmaz M, Telo S, Atescelik M, et al., Spexin level in acute myocardial infarction in the emergency department. *J Med Biochem.* 2023 Aug 25; 42(3): 407-411. <https://doi.org/10.5937/jomb0-39485>.
6. Seshiah V, Balaji V, Chawla R, Gupta S, Jaggi S, Anjalakshi C, et al., Diagnosis and management of gestational diabetes mellitus guidelines by DIPSI (Revised). *Int J Diabetes Dev Ctries.* 2023 Aug; 43(4): 485-501. <https://doi.org/10.1007/s13410-023-01222-3>.
7. Balke S, Weid P, Fangmann L, Rostin P, Henrich W, Koenigbauer JT. Glucose Levels of the Oral Glucose Tolerance Test (oGTT) Can Predict Adverse Pregnancy Outcomes in Women with Gestational Diabetes (GDM). *J Clin Med.* 2023 May 27; 12(11): 3709-3722. <https://doi.org/10.3390/jcm12113709>.
8. Zhang K, Zhang W, Xia F, Wang N, Lu Y, Sui C, et al. Obesity Patterns, Metabolic Abnormality, and Diabetic Kidney Disease in Patients with Type 2 Diabetes. *Diabetes Metab Syndr Obes.* 2023 Dec 31: 3999-4011. <https://doi.org/10.2147/DMSO.S436046>.
9. Chittezhath M, Gunaseelan D, Zheng X, Hasan R, Tay VSY, Lim ST, et al. Islet macrophages are associated with islet vascular remodeling and compensatory hyperinsulinemia during diabetes. *Am J Physiol Endocrinol Metab.* 2019 Dec 1; 317(6): E1108-E1120. <https://doi.org/10.1152/ajpendo.00248.2019>.
10. Stanimirovic J, Radovanovic J, Banjac K, Obradovic M, Essack M, Zafirovic S, et al. Role of C- Reactive Protein in Diabetic Inflammation. *Mediators Inflamm.* 2022 May 17; 2022: 3706508. <https://doi.org/10.1155/2022/3706508>.
11. Cardoso CR, Leite NC, Salles GF. Prognostic Importance of C- Reactive Protein in High Cardiovascular Risk Patients With Type 2 Diabetes Mellitus: The Rio de Janeiro Type 2 Diabetes Cohort Study. *J Am Heart Assoc.* 2016 Oct 26; 5(11): e004554. <https://doi.org/10.1161/JAHA.116.004554>.
12. Arab Sadeghabadi Z, Abbasalipourkabir R, Mohseni R, Ziamajidi N. Investigation of oxidative stress markers and antioxidant enzymes activity in newly diagnosed type 2 diabetes patients and healthy subjects, association with IL-6 level. *J Diabetes Metab Disord.* 2019 Sep 9; 18(2): 437-443. <https://doi.org/10.1007/s40200-019-00437-8>.
13. Shareef MS, Charfeddine B, Roomi AB. The Association between Spexin Levels, Lipid Profiles, Glucose Metabolism, and Immune Function in Newly Diagnosed Diabetes in Postmenopausal Iraqi Women. *Rev Diabet Stud.* 2023; 19(3): 166-170. <https://doi.org/10.1900/RDS.2029.19.166>.
14. Tejaswi G, Dayanand CD, Prabhakar K. Insulin resistance and decreased spexin in Indian Patients with Type 2 Diabetes Mellitus. *Bioinformation.* 2021 Sep 30; 17(9): 790-797. <https://doi.org/10.6026/97320630017790>.
15. Wilson PW, D'Agostino RB, Parise H, Sullivan L, Meigs JB. Metabolic syndrome as a precursor of cardiovascular disease and type 2 diabetes mellitus. *Circulation.* 2005 Nov 15; 112(20):3066-72. <https://doi.org/10.1161/circulationaha.105.539528>.
16. Chang CM, Hsieh CJ, Huang JC, Huang IC. Acute and chronic fluctuations in blood glucose levels can increase oxidative stress in type 2 diabetes mellitus. *Acta diabetologica.* 2012 Dec; 49: 171-177. <https://doi.org/10.1007/s00592-012-0398-x>.
17. Roomi AB, AL-Salih RM, Ali SA. Impact metformin and insulin therapy on parathyroid hormone and 25 (OH) vitamin D in diabetic post-menopausal Iraqi women. *J Phys Conf Ser* 2019 Jul 1 (Vol. 1279, No. 1, p. 012008). <https://doi.org/10.1088/1742-6596/1279/1/012008>.
18. Kawahito S, Kitahata H, Oshita S. Problems associated with glucose toxicity: role of hyperglycemia-induced oxidative stress. *World J Gastroenterol: WJG.* 2009 Sep 9; 15(33): 41374142-. <https://doi.org/10.3748/wjg.15.4137>.

سببسين كمؤشر لدى العراقيات المصابات بالسكري في سن اليأس

مصطفى سليم شريف¹، علي بدر رومي²، باسم شرف الدين³

¹ قسم الكيمياء الحيوية، كلية التقنيات الصحية والطبية، جامعة البیان، بغداد، العراق.

² قسم الكيمياء، كلية العلوم، جامعة ذي قار، ذي قار، 64001، العراق.

³ قسم الكيمياء الحيوية، كلية الطب، جامعة سوسة، تونس.

الخلاصة

يبتدئ سببسين (SPX) ينظم نسبة الجلوكوز في الدم واستقلاب الدهون. ركزت على مسببات مرض السكري من النوع 2 (T2DM) لدى النساء بعد انقطاع الطمث، ودراسة علاقتها بمستويات SPX في الدم. الطريقة: دراسة الحالات والشواهد تطوعت 180 امرأة بعد انقطاع الطمث من ذي قار. تم تسجيلها في الدراسة الحالية. تم تشخيص إصابة 90 امرأة بالسكري في مرحلة ما بعد انقطاع الطمث، في حين أن الـ 90 الأخرى غير مصابات بالسكري بعد انقطاع الطمث. تم جمع عينات الدم لقياس HbA1c و SPX و MDA و SOD و CRP، جلوكوز الدم الصائم (FSG)، الأنسولين، lipid Profile، والإنترلوكين (IL-6). النتائج: ارتفاع بقيم FSG، HbA1c، والأنسولين، مقاومة الأنسولين (HOMA-IR) بشكل ملحوظ ($P < 0.001$) في النساء بعد انقطاع الطمث من النوع T2DM. كما أن مستويات MDA و CRP و IL-6 كانت مرتفعة بشكل كبير، كما انخفضت مستويات SPX و SOD بشكل ملحوظ ($P < 0.001$). وقد لوحظت تغييرات كبيرة في الدهون في النساء بعد انقطاع الطمث و T2DM. أظهر الارتباط في T2DM لدى النساء بعد انقطاع الطمث علاقة سلبية بين SPX مع نسبة الخصر إلى الطول، SPX مع FSG، SPX مع HOMA-IR، MDA مع SPX، وجود علاقة إيجابية بين SPX و MDA، SPX والعمر، CRP و IL-6، CRP والعمر، IL-6 و FSG، SOD والأنسولين. الاستنتاج: لقد ثبت أن انخفاض SPX مهم في التسبب في مرض T2DM لدى النساء بعد انقطاع الطمث. ، قد يوفر استهداف SPX استراتيجية جديدة للحفاظ على التحكم في الجلوكوز.

الكلمات المفتاحية: بروتين سي التفاعلي، مالونديالدهيد، سببسين، سوبروكسيد ديسموتاز ومرض السكري من النوع الثاني.