

Effect of Pregnancy on Selenium, Copper ,Zinc and Others Biochemical Feacture

*Layla O.Farhan**

*Seenaa A.Mustafa**

*Noora S.Mubder**

Received 6, June, 2012

Accepted 26, November, 2012

Abstract:

Pregnancy is a stressful condition in which many physiological and metabolic functions are altered to a considerable extent . Pregnancy is a physiological state accompanied by a high-energy demand and an increased oxygen requirement. The present study aim to study selenium ,zinc copper in the first trimester of pregnancy. The study group comprised of Fourty five pregnant women and twenty six non pregnant women as control . The samples were taken from pregnant women who come to several heath center in Baghdad city to cheak up. Laboratory investigations including Cupper, Ceruloplasmin, Total Antioxidant (TAA), malondialdehyde (MDA), glutathione (GSH), Zinc, Uric acid, and Selenium had been measured in pregnant women and control . There were significant difference in Cupper, Selenium, Ceruloplasmin, TAA, Zinc, GSH, and MDA, in the pregnant women whene compared to control group. In this study, a significantly positive association was observed between zinc [mg/dl] with ceruloplasmin [mg/dl] ($R = 0.47$ $p < 0.01$) and TAA [mmol/l] ($R = 0.42$ $p < 0.05$) , and negative association was observed between zinc [mg/dl]and cupper [mg/dl] ($R = 0.602$, $p < 0.01$) , a significantly negative association between ceruloplasmin [mg/dl] with cupper [mg/dl] ($R = 0.754$, $p < 0.01$) ,and positive with TAA [mmol/l]($R = 0.562$ $p < 0.01$). The present study observed that pregnant women in first trimester were more susceptible to oxidative damage than in non pregnant as indicated by increased lipid peroxidation (MDA) and decreased other antioxidanta

Key words: Pregnancy, Cupper ,Zinc , Ceruloplasmin , MDA .

Introduction:

Pregnancy is associated with increased demand of all the micronutrients like Iron, copper, zinc, vitamin B12, folic acid and ascorbic acid [1]. The deficiency of these nutrients could affect pregnancy, delivery and outcome of pregnancy. Vitamins and minerals are collectively re-ferred to as micronutrients and have important influence on the health of pregnant women and growing fetus [2]. The trace elements namely zinc and copper are necessary during

pregnancy and these elements should be supplemented as a daily requirement to the pregnant women [3, 4]. Zinc is essential for the growth and development of human life and has an active role in body function. During pregnancy, zinc is also used to assist the fetus to develop the brain [5]. The trace mineral selenium is an essential component of the selenoproteins, especially glutathione peroxidase, required for normal health and reproduction. Selenium deficiency in humans can result in an increased

* Department of Chemistry, College of Science for women, Baghdad University, Baghdad, Iraq.

susceptibility to viral infections, cancer, cardiovascular disease, miscarriage [6,7]. Various studies have demonstrated that during pregnancy, the whole blood and plasma selenium concentrations, and the activity of glutathione peroxidase in red cells and plasma decline in a linear fashion from the first trimester to parturition, with the lowest levels at delivery [8]. It has also been demonstrated that the requirement of selenium is increased during pregnancy as a result of transport to the growing fetus [9]. Glutathione, an intracellular tripeptide of the amino acids cysteine, glycine and glutamic acid is the most abundant thiol-based antioxidant compound found in living animal and plant tissues. Glutathione participate directly in the neutralization of free radicals, reactive oxygen compounds and maintains exogenous antioxidants such as vitamins C and E in their active forms [10]. Glutathione also plays a role in the detoxification of many xenobiotics, it helps in the storage and transport of cysteine and maintain the reduced state of proteins and thiols [11]. The present study aim to study selenium ,zinc copper in first trimester pregnancy.

Materials and Methods:

A Five ml of blood serum had been collected from each subject by vein puncture, centrifuged at 3000 rpm for 5 min after clotting at room temperature. Fourty five serum sample obtained from pregnant women age (19-35) years ($M \pm SD$: 27.4 ± 3.44). Twenty six healthy subjects age (19-35) years ($M \pm SD$: 26.5 ± 3.82), copper

and zinc was estimated by atomic absorption spectrometry (AAS) method [12]. Serum ceruloplasmin was estimated by kinetic method [13]. Glutathione concentration was estimated by the method described by Tietz 1986 [14]. Selenium concentration of serum was measured using a spectrofluorometer by the modified method of Kho and Benson [15]. The operating conditions for the spectrofluorometer equipped with a UV lamp were an excitation wavelength of 369 nanometers and an emission wavelength of 525 nanometers. MDA was measured by Thiobarbituric acid reactive substances assay (TBRAS) method [16]. The uric acid was measured by uricase method using Human diagnostics kit [17-19]. Serum total antioxidants activity estimated by the method described by [20]. Statistical analyses of this study were performed using SPSS version 15.0 for Windows (Statistical Package for Social Science, Inc., Chicago, IL, USA). Descriptive analysis was used to show the mean and standard deviation of variables. The significance of the difference between mean values was estimated by Student T-Test. The probability $P < 0.05$ = significant, $P > 0.05$ = non-significant. Correlation analysis was used to test the linear relationship between parameters. ANOVA test was used to show the differences between variables of differentiated groups.

Results and Discussion:

There is no significant different in age between pregnant women and control group as shown in table 1.

Table(1): The mean and standard deviation of Age and Gastation in 1st pregnancy in trimester women group and control group.

Characteristic	Pregnant women mean±SD	Control mean±SD	P Vale
Age (year)	27.4 ±3.44	26.5±3.82	N.S
Gastation Age (week)	10.51±1.01	-----	

In pregnant women the mean cupper, ceruloplasmin ,and MDA had significantly increased ($p<0.01$), ($p<0.05$) and ($p<0.001$) respectivity . TAA, zinc, and GSH were found to be significantly decrease with P value <

0.05 ,selenium was significantly decreased($p<0.01$) . There were no significant difference in level of uric acid in pregnancy women group when compared to control group as shown in Table 2.

Table (2): The mean and standard deviation of cupper, Ceruloplasmin,TAA, MDA ,GSH, zinc , uric acid and selenium in pregnancy in 1st trimester women group and control group.

Characteristic	Pregnant women mean±SD	Control mean±SD	P Vale
Cupper (mg/dl)	157±12.36	130.8±17.87	<0.01
Selenium(mg/ml)	59.66±3.6	68.22±3.76	<0.01
Ceruloplasmin (mg/dl)	41.28±5.7	34.9±7.07	<0.05
TAA (mmol/l)	1.6±0.13	1.8±0.13	<0.05
Zinc (mg/dl)	76.42	79.53±9.52	<0.05
GSH(mg/dl)	41.8±2.8	46.94±3.55	<0.05
MDA (m mol/l)	4.30±8.81	1.51±0.24	<0.001
Uric acid (mg/dl)	3.87±0.74	3.6±0.64	N.S




Fig (1): Correlation between zinc with ceruloplasmin ,TAA and copper in pregnancy and non pregnant women.




Fig (2) : Correlation between ceruloplasmin with ,TAA and copper in pregnancy and non pregnant women.

The glutathione level showed a significant decreased ($P<0.05$) pregnant women as in agreement with previous reports [21]. Depletion in serum level of glutathione has also been linked with lipid peroxidation [22], which occurs as a result of hypertension and thus the production of free radicals that damage the endothelial tissues [23]. In the present study, the serum concentration of malondialdehyde (MDA) is higher in 1st pregnant women than control. This is in agreement with earlier study by Patil *et al* [22, 24]. This may be due to increased generation of ROS because of increased oxygen demand during pregnancy [25], reduction in the activities of antioxidant enzymes

such as superoxide dismutase and glutathione peroxidase, and reduction in plasma vitamin C and vitamin E concentration during pregnancy [26].

The present study observed a significant decrease in all the trace elements in Pregnant women when compared to the control group. This suggests a possible involvement of zinc, copper, selenium in the development and pathogenesis of Pregnancy, current data agrees with that of Mahomed *et al.* [27] who reported reduced zinc concentration in women with Pregnancy. Decrease in the level of selenium is known to enhance Pregnancy by its resultant effect on glutathione peroxidase. This is important as increased oxidative

stress as a result of low density lipoproteins has been confirmed in Pregnancy and this complication is strictly regulated by antioxidants of which selenium forms a part (Witztum, 2001)[28]. The present study showed that the concentration of ceruloplasmin was high compared to the non-pregnant controls ($p < 0.05$). Elevated concentrations of ceruloplasmin in pregnancy represent perhaps a response to an oxidative injury [29].

In this study, a significantly positive association was observed between zinc [mg/dl] with ceruloplasmin [mg/dl] ($R = 0.470$, $p < 0.01$), TAA [mmol/l] ($R = 0.42$ $p < 0.05$), and negative association was observed with copper [mg/dl] ($R = 0.602$, $p < 0.01$), in pregnant women while there was no significant correlation observed in the non pregnant women group as shown in figure 1. a significantly negative association was observed between ceruloplasmin [mg/dl] with copper [mg/dL] ($R = 0.754$, $p < 0.01$), and positive association was observed with TAA [mmol/l] ($R = 0.562$, $p < 0.01$) in pregnant women while there was no significant correlation was observed in the non pregnancy women group as shown in figure 2.

Conclusions :

Increased in lipid peroxidation MDA and decreased in TAA, Selenium, Copper, zinc and other Antioxidants in the present study may be due to oxidative damage in pregnant women and counteract the cellular changes mediated by free radicals.

References:

1. Rayman MP. 2000 The importance of selenium to human health. , Lancet; Pub Med , 15: 35, (9225) 233-41.
2. Savita R., Ashish G. and Hitender S Batra, 2011 " Comparative study of

trace elements and serum ceruloplasmin level in normal and pre-eclamptic pregnancies with their cord blood" JBR 2011; 22 (2): 207-210

3. Barrington JW, Lindsay P, James D et al. 1996 " Selenium deficiency and miscarriage : A possible link ? Br J Obst Gyn ;103:130-2.
4. AMA Department of Foods and Nutrition. Guideline for Essential Trace Elements Preparation for Parinatal Use: A Statement by Expert Panel JAEN J parenter Entral Nutr. 3(4):263-7.
5. Moutquin J. M., Rainville C. and Giroux L. A 1985 Prospective Study of Blood Pressure in Pregnancy: Prediction of Pre-Eclampsia. Br J Obst Gyn; 151: 191-195.
- 6- Batra HS, Singh P, Somani BL, Gupta A, Sampath S, and Ambade V. 2007 Serum ferroxidase albumin ratio as a marker in pulmonary tuberculosis. Ind J Clin Biochem; 22: 106-108.
- 7- Cheesman KH and Slater TF. 1993 An introduction to free radical biochemistry. Br Med Bull;49:481-93.
8. Pankaj D , Purvi P and Shailaja M . 2006" Selenium levels and glutathione peroxidase activity in spontaneous inevitable abortion." Br J Obst Gyn Vol. 56, No. 4P: 311-314
- 9- Kumru S, Aydin S, Simsek M, Sahin K, Yaman M. Comparison of Serum Copper, Zinc, Calcium, and Magnesium Levels in Pre-Eclamptic and Healthy Pregnant Women. Biol Trace Elem Res 2003; 94: 105-112.
- 10- Witschi A, Reddy S and Stofer B. 1992 Lauterburg; Eur. Jour. Clin. Pharmacol., 43(6): 667-669.
- 11- Raijmakers M , Roes E, Poston L and Steegers W . 2009 Peters; Eur. Jour. Br J Obst Gyn., 138(1): 39-44

- 12-Wildman REC and Medeiros DM. 2000 Advanced human nutri-tion. Boca Raton, CRC Press,.
13. Somani BL and Ambade V. 2005 Novel composition for kinetic assay of ceruloplasmin. Clin Chem; 51: Suppl A90.
- 14- Teitz NW 1986 : - Determination of Cobalamine and Folic acid. In. Textbook of Clinical Chemistry W. B. Saunders and Company. Philadelphia, London 3rd edition pp.1571–1573.
- 15 -Koh T and Benson TH. (1982)Critical re-appraisal of fluorometric method for determination of selenium in biological materials. J Assoc Off Anal Chem 66:918–926.
- 16- Stocks, J and Dormandy ,T. 1971 "The autoxidation of human red cell. Lipid induced by hydrogen peroxide." Br. J. Haematol. 20(1) : 95-111 .
17. Barham D and Trinder P. 1972 An improved colour reagent for the determination of blood glucose by the oxidase system. Analyst; 97(151) : 142-145
18. Fossati LP and Berti G. 1980 Use of 3,5-dichloro-2-hydroxybenzenesulfonic acid/4-aminophenazone chromogenic system in direct enzymic assay of uric acid in serum and urine. Clin Chem;26(2):227-231.
- 19 .Thefeld L. Dtsch. 1973 Untersuchungen zur Frage der Hämodialyse bei der Knollenblätterpilzvergiftung.Med. Wschr; 98: 300 -309
- 20- Arinola, O.G., Adedapo K.S., Kehinde A.O., Olaniyi J.A. and Akiibinu, M.O. 2004 . Acute phase proteins, trace elements in asymptomatic human immune deficiency virus infection in Nigerians. Afr. J. Med. med. Sci.33: pp: 317-322.
- 21- Modupe F. and Samuel S. 2010 . Plasma thiol levels in pregnancy induced hypertension in Nigerians. Scholars Research Library J. 2 (2):443-447.
- 22- Luqman A. O. , Salihu M. A. , Gafar A.A. and Ayodele O.S. 2008 " Effect of vitamin c on malondialdehyde (MDA) in pregnant Nigerian women." Basic and Applied Sciences J. Vol. 4, No. 2, 105-108.
- 23- Zachara BA, Dobrzanski W, Trafikowska U et al. 2001 Blood selenium and glutathione peroxidases in miscarriage. BJOG;108: pp 244-247.
- 24-Patil SB, Kodliwadmth MV and Kodliwadmth SM. 2006 . Lipid peroxidation and nonenzymatic antioxidants in normal pregnancy. Br J Obst Gyn, 56(6): 399-401.
- 25-Halliwell B. and Gutteridge JM. 1990 . Role of free radicals and catalytic metal ions in human disease: an overview method in Enzymology (B), 186: 1-85.
- 26-Walsh SW and Wang Y. 1993 . Secretion of lipid peroxides by the human placenta American Journal of Obst Gyn, 169(6) : 1462-1466.
- 27- Mahomed K, Williams MA, Woelk GB, Mudzamiri S, Madzime S, King IB and Bankson DD 2004 . Leckocyte selenium, zinc and copper concentrations in preeclampsia and normotensive pregnant women. Biol. Trace Elem. Res. 75 (1-3) : 107-118.
- 28-Witztum J 2001 The oxidation hypothesis of atherosclerosis. Lancet, 344 (8925): 793-795.
- 29-Marana M and Otila,M .2001 ."Correlation between reactive oxygen species and homocysteine levels in normal pregnancy " .Farmcia ,59(2),P(179-190).

تأثير الحمل على السلينيوم والنحاس والزنك وبقية العوامل الكيميائية الحياتية

ليلى عثمان فرحان * سينا عيود مصطفى * نورا سعد مبدن *

* قسم الكيمياء، كلية العلوم للبنات، جامعة بغداد.

الخلاصة :

الحمل هو حالة من التوتر في العديد من الوظائف الفسيولوجية والتمثيل الغذائي وهو حالة الفسيولوجية يرافقتها طلب متزايد للاوكسجين والطاقة. هدفت الدراسة الحالية هو دراسة تأثير السلينيوم والزنك والنحاس في فترة الثلث الاول من الحمل. شملت الدراسة 45 من النساء الحوامل و 26 من النساء غير الحوامل (كمجموعة ضابطة) من النساء اللواتي يراجعن المركز الصحي في بغداد للاطمئنان على الصحة. شملت الفحوصات المختبرية قياس النحاس والسلينيوم والسيروبلازمين والزنك و MDA و TAA و حامض اليوريك و GSH في النساء الحوامل ومجموعة السيطرة. لوحظ وجود زيادة مقبولة احصائيا في النحاس والسلينيوم والسيروبلازمين والزنك و TAA و حامض اليوريك و GSH في النساء الحوامل فقط عند مقارنتها بالمجموعة القياسية. وقد اشارت الدراسة الى وجود علاقة خطية مقبولة احصائيا بين الزنك والسيروبلازمين ($r = 0.42$, $p < 0.05$) و TAA ($R=0.42$, $p < 0.05$) و النحاس ($R=0.602$, $p < 0.01$) وبين السيروبلازمين و النحاس ($P < 0.01$) و TAA ($R=0.754$, $p < 0.001$) و النحاس ($R= 0.562$, $p < 0.001$) للنساء الحوامل، بينما لاوجود لهذة العلاقة عند النساء غير الحوامل. اوضحت الدراسة الحالية ان النساء الحوامل في المرحلة الاولى من الحمل اكثر عرضة لضرر الاكسدة من النساء غير الحوامل كما مشار الية بزيادة MDA وانخفاض مضادات الاكسدة الاخرى.