

Evaluating Coll2-1NO2 Level in Pre- and Post-menopausal Iraqi Women with Osteoarthritis

Noor Basil Ghanim* 🔍 🖄, Bushra Faris Hasan

Department of Chemistry, College of Science for Women, University of Baghdad, Baghdad, Iraq. *Corresponding Author.

Received 28/03/2023, Revised 18/06/2023, Accepted 20/06/2023, Published Online First 20/01/2024

© 2022 The Author(s). Published by College of Science for Women, University of Baghdad. This is an Open Access article distributed under the terms of the <u>Creative Commons Attribution 4.0 International</u> <u>License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Osteoarthritis (OA) is not an autoimmune but a chronic inflammatory disease affecting the joints of humans due to mechanical stress mostly in the knees and the hip. In OA a modification of the cartilage is encountered causing a pain in the joints upon movement. Coll2-1NO2 is a specific sequence of amino acids produced from the degradation of type II collagen, which is abundantly found in the cartilage, followed by a nitration in the tyrosine residue as a consequence of stress-inflammatory events. In this study, Coll2-1NO2 is used as a biomarker of OA to predict the inflammatory stress condition as well as the cartilage degradation in pre- and post-menopausal women. The results showed a significant increase of serum Coll2-1NO2 in pre- and post-menopausal women with OA compared to pre- and post-menopausal health women, at similar body mass index category (overweight). Pearson's correlation analysis showed non-significant association between Coll2-1NO2 and other studied parameters. Moreover, ROC analysis showed a very excellent sensitivity of Coll2-1NO2 as a prognostic biomarker for OA disease in pre- and post-menopausal women.

Keywords: Coll2-1NO2, C-reactive protein, FSH, lipid profile, osteoarthritis.

Introduction

Osteoarthritis (OA) is not an autoimmune but a chronic inflammatory disease affecting the joints of humans due to mechanical stress mostly in the knees and the hip ¹. OA is also known as degenerative arthritis and considered as a major health problems that faces the elder people ². The global prevalence rate of OA has reached a serious point, where the major incidence rates were among women, elders, people with overweight and obesity, and certain ethnics ³. OA was once thought to be only a "wear and tear" disease. Prolonged stress and poor biomechanics on the joints were thought to be the root causes of the breakdown of cartilage in the

joint and the ensuing inflammation. Stiffness, edema, and impaired movement were the results. It is now understood that OA is a much more complex condition made up of inflammatory and metabolic factors ⁴⁻⁶. Women at postmenopausal stage exhibit wide spectrum of fluctuations in their gonadal hormones, and signal transduction system ⁷ that can increase the mechanical stress ⁸ and inflammatory events ⁹. Therefore, it is important to determine the differences in the pathophysiology of OA between pre- and post-menopausal women.

The complex structure of the cartilage enables a smooth, free of pain movement for humans. In OA

condition, this structure is disrupted and its function declined, causing a pain in the joints during the movement ¹⁰⁻¹². It has been reported that age is a crucial factor for the modification of the cartilage leading to the development of OA, through several mechanisms including oxidative stress and oxidative damage ¹³. The extracellular matrix of the cartilage contains high amount of type II collagen which gives a unique and specific sequence of amino acid residues (His-Arg-Gly-Tyr-Pro-Leu-Asp-Gly) upon denaturation called Coll2-1^{14, 15}. Hence, Coll2-1 can be used as specific biomarker for the degradation of collagen type II, and consequently for OA since it degraded during this degenerative arthritis ¹⁶. More recently, a modified form of Coll2-1 has been introduced, known as Coll2-1NO₂, which has a similar sequence but with

Materials and Methods

One hundred and fifty female subjects were enrolled in this work, 80 of which were diagnosed with OA by the specialized medical doctors at Orthopedic Consultancy Department of Baghdad Hospital, at Baghdad Medical City. The rest of the female subjects were healthy and used as control in the study. Moreover, the OA patients were divided into two subgroups based on the age of the participants, namely, OA I and OA II. The first

Methods

The analyses were involved the assessment of ESR by using the whole blood sample at the moment of blood sample collecting. Serum samples were used to analyze CRP, LH, FSH, and testosterone by using cobas commercial kits on cobas b101 of CRP and cobas E411 autoanalyzer for the rest (Roche, Germany). Coll2-1NO₂ was analyzed in the serum by using Sandwich based ELISA MyBioSource research kit in BioTech (USA) microplate reader. At last, lipid profile parameters were detected by

Results and discussion

In Table 1 comparisons among the subgroups are demonstrated by using the mean value and the standard deviation, encountered by the *P*-value that obtained from ANOVA analysis. According to the design of the work the age of control I and OA I

a nitrated tyrosine residue at the position 4 of the sequence. Coll2-1NO₂ can be used as an indicator for inflammatory and oxidative/nitrative damage of the cartilage ¹⁷. Since the recent researches on OA have shown a link the inflammatory events in the development and/or the progressiveness of the disease, Coll2-1NO₂ can give a better indication on the condition of OA. We have aimed to assess Coll2-1NO₂ in the sera of pre- and post-menopausal women with OA disease to predict the age-related effect on this biomarker as well as its correlation Follicle-Stimulating Hormone with (FSH), Luteinizing Hormone (LH), Testosterone, and inflammatory biomarker including Erythrocyte Sedimentation Rate (ESR), and C-Reactive Protein (CRP).

subgroup contained 40 pre-menopausal women with OA disease while the second subgroup contained 40 post-menopausal women with OA disease. Also, control group was divided into similar groups based on their age (Control I and Control II) where each group contained 35 females. From each female, the veinous blood was collected and the serum was obtained from the blood sample and stored at -20 °C to be analyzed later.

spectrophotometric kits (Biolabo, France) for triglycerides (TGs), total cholesterol (TC), and high-density lipoprotein cholesterol (HDL-C). Finally, the data were analyzed statistically for mean comparisons (ANOVA, and LSD post-Hoc test) where the results were expressed as mean \pm standard deviation (SD), correlation (Pearson's coefficient), and receiver operating characteristics (ROC) in SPSS software program (version 26.0).

were under 45 years old and significantly differed from control II and OA II who were above 45 years old. Body mass index (BMI) of the participants lies at the overweight category for all subgroups. The serum levels of Coll2-1NO₂ were at their highest value in OA II patients (14.44 ± 2.40 ng/mL), where OA I patients exhibited a high level of the biomarker as well (10.96 ± 2.32 ng/mL). This increase of Coll2-1NO₂ was significant compared to the corresponding control groups (OA I higher than control I, and OA II higher than control II). Moreover, OA II showed significant higher levels of Coll2-1NO₂ compared to OA I. The obtained results of CRP and ESR were similar to that of $Coll2-1NO_2$ in which OA II patients showed the highest values. Calcium on the other hand decreased in OA I and II subgroups compared to the corresponding control groups. TGs was the only parameter in the lipid profile which was significantly higher in OA I and OA II subgroups compared to the corresponding control groups.

Parameter	Control I	Control II	OA I	OA II	<i>P</i> -value
Age (year)	34.77±5.00	55.20±4.35	35.93±4.36 ^c	56.05 ± 4.64	< 0.001
BMI (Kg/m ²)	27.74±3.73	28.01±2.06	28.92±2.97	29.22±2.77	0.094
Coll2-1NO ₂ (ng/mL)	5.00±1.01 ^a	4.56 ± 0.86^{b}	10.96±2.32 ^c	14.44 ± 2.40	< 0.001
FSH (mIU/mL)	5.10±2.75	34.12±8.19	$6.77 \pm 2.60^{\circ}$	35.11±8.58	< 0.001
LH (IU/L)	6.80±3.85	31.07±8.72	$5.84 \pm 3.56^{\circ}$	30.00±7.65	< 0.001
Testosterone (nmol/L)	0.88±0.25	0.85±0.32	0.80±0.29	0.89±0.32	0.601
ESR (mm/h)	18.37 ± 8.35^{a}	$22.77 {\pm} 6.52^{b}$	$39.15 \pm 5.18^{\circ}$	43.25±6.03	< 0.001
CRP (mg/L)	$3.72{\pm}1.49^{a}$	6.67 ± 2.20^{b}	$13.18 \pm 5.26^{\circ}$	16.90±6.22	< 0.001
Calcium (mg/dL)	$9.27{\pm}0.70^{a}$	9.01 ± 0.60^{b}	8.68±0.63	8.61±0.69	< 0.001
TGs (mg/dL)	$125.03{\pm}26.78^{a}$	126.58±23.88 ^b	139.23±31.68	143.30±27.56	0.009
TC (mg/dL)	163.69±26.34	171.60 ± 24.08	168.98 ± 26.50	174.38 ± 24.80	0.321
HDL-C (mg/dL)	41.04±6.58	38.09±5.42	38.80±6.92	39.30±5.32	0.216

Table 1. Outcomes of controls and OA patients

Letters on values indicate significant differences (a: Control I vs OA I, b: Control II vs OA II, c: OA I vs OA II).

The relationship between $Coll2-1NO_2$ and the other studied parameters is illustrated in Table 2, which showed that there was no significant association.

Table 2. Correlation of Coll2-1NO2 with the other variables in OA patients

Variables	Pre-mer	opausal	Post-mei	Post-menopausal	
variables	r	Р	r	Р	
Age	0.200	0.216	0.077	0.637	
BMI	-0.141	0.384	-0.042	0.798	
ESR	0.153	0.346	0.137	0.401	
CRP	0.092	0.574	-0.196	0.225	
FSH	0.132	0.418	-0.131	0.421	
LH	0.101	0.534	-0.039	0.810	
Testosterone	-0.074	0.648	0.006	0.970	
Calcium	-0.194	0.231	0.306	0.055	
TGs	-0.009	0.956	-0.075	0.646	
TC	-0.144	0.375	-0.156	0.336	
HDL-C	0.279	0.082	-0.033	0.840	



The ROC analysis indicated very excellent sensitivity (the area under the curve is 1.000) for Coll2-1NO2 in the prognosis of OA disease in women at both pre- and post-menopausal age. In pre-menopausal women with OA disease, a cut-off value of 7.29 ng/mL can be used with 100%

sensitivity and 100% specificity as showed in Fig. 1A, while in post-menopausal women with OA disease, a cut-off value of 8.38 ng/mL ,can be used with 100% sensitivity and 100% specificity as shown in Fig. 1B.



Figure 1. ROC curve of Coll2-1NO2 in the prognosis of OA disease for (A) pre-menopausal women, (B) post-menopausal women.

Pre- and post-menopausal women with OA disease were submitted to Coll2-1NO₂ analysis in their serum. As shown in Table 1, all of Coll2-1NO₂, CRP, and ESR were elevated significantly in OA patients at both pre- and post-menopausal age scale. Nevertheless, CRP and ESR cannot be used as specific biomarkers for OA disease considering the wide spectrum affecters of these two inflammatory biomarkers, in which they are increased in other conditions such as rheumatoid arthritis ¹⁸. Yet, Coll2-1NO₂ can be used as specific biomarkers for OA considering its production from the degradation of type II collagen that was found predominantly in the cartilage. Also, the nitration of its sequence that arises from inflammatory gives an impression of the stress-induced inflammation that accompanied OA disease. Hick et al. reported that both Coll2-1 and Coll2-1NO₂ were linked to knee OA and can be used to predict the knee cartilage features in patients. Also, they showed quite engagement between Coll2-1NO₂ serum level and pain in OA patients ¹⁴. Mutar et al. showed that serum Coll2-1NO₂ level increased significantly in both males and females with OA disease. Additionally, their reports agreed with our findings regarding the use of Coll2-1NO₂ as predicative biomarker of OA disease ¹⁹.

The levels of FSH, LH, and Testosterone nonsignificantly changed between OA patients and the age corresponding control groups. Yet, it was reported that FSH in post-menopausal women with OA disease can enhance the pain worsening in these patients through decreasing the type II collagen concentration in the cartilage of the knee 20 . Nevertheless, we did not find a correlation between Coll2-1NO₂ and FSH in post-menopausal women with OA disease. But it can explain the significant high level of OA II compared to OA II (Table 1). Like FSH, high levels of LH in post-menopausal women with OA were reported to enhance the progression of OA disease²¹. Furthermore, TGs level in OA patients was clearly increased compared to healthy women at both pre- and postmenopausal age. Hypertriglyceridemia is linked to the lifestyle of individuals predominantly ²². Since all of the enrolled women were overweight, high TGs is expected, as it was reported by previous studies $^{23-25}$. Increased TGs level in OA patients enhances the inflammatory condition 26 of these

Conclusion

In conclusion, high level of $Coll2-1NO_2$ in serum of OA patients can predict high cartilage damage with the presence of stress-inflammatory condition in these patients. It is safe to presume that $Coll2-1NO_2$

Acknowledgment

We would like to express our gratitude for Department of Chemistry at College of Science for Women, University of Baghdad for their approval

Author's Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for

Authors' Contributions

N. B. G. collected the specimens, carried out the research experiment, analyzed, and discussed the

References

- Mutar H S, Hasan B F, Muhi S A. Study the Level of Cartilage Acidic protein (CRTAC1) in Serum of Iraqi Patients with osteoarthritis, Neuro Quant. 2022; 20(7): 153-158. https://doi.org/10.14704/nq.2022.20.7.NQ33018
- Kraus V B, Blanco F J, Englund M, Karsdal M A, Lohmander L S. Call for standardized definitions of osteoarthritis and risk stratification for clinical trials and clinical use. Osteoarthr Cartil. 2015; 23(8): 1233-1241. <u>https://doi.org/10.1016/j.joca.2015.03.036</u>
- 3. Allen K D, Thoma L M, Golightly Y M. Epidemiology of osteoarthritis. Osteoarthr Cartil. 2022; 30(2): 184-195. https://doi.org/10.1016/j.joca.2021.04.020
- Abramoff B, Caldera F E. Osteoarthritis: pathology, diagnosis, and treatment options. Med Clin. 2020; 104(2): 293-311. https://doi.org/10.1016/j.mcna.2019.10.007
- 5. Musumeci G, Aiello F C, Szychlinska M A, Di Rosa M, Castrogiovanni P, Mobasheri, A, et al. Osteoarthritis in the XXIst century: risk factors and behaviours that influence disease onset and

patients and can lead to increase the pain accompanying this disease.

can be used as a specific and sensitive biomarker for OA disease based on the collected data in this study and previously mentioned studies.

and scientific and administrative help in this work, and the staff of Baghdad Hospital for their help in specimens' collection.

re-publication, which is attached to the manuscript.

- Ethical Clearance: The project was approved by the local ethical committee in University of Baghdad.

results. B F. H. put the design of the work and supervised the project.

progression. Int J Mol Sci. 2015; 16(3): 6093-6112. https://doi.org/10.3390/ijms16036093

- Mobasheri A, Batt M. An update on the pathophysiology of osteoarthritis. Ann Phys Rehabil. 2016; 59(5-6): 333-339. https://doi.org/10.1016/j.rehab.2016.07.004
- Slavich G M, Sacher J. Stress, sex hormones, inflammation, and major depressive disorder: Extending Social Signal Transduction Theory of Depression to account for sex differences in mood disorders. Psychopharmacol. 2019; 236(10): 3063-3079. https://doi.org/10.1007/s00213-019-05326-9
- Hughes-Fulford, M. Signal transduction and mechanical stress. Science's STKE. 2004; 2004(249): 1-8. <u>https://doi.org/10.1126/stke.2492004re12</u>
- Taleb-Belkadi O, Chaib H, Zemour L, Fatah A, Chafi B, Mekki, K. Lipid profile, inflammation, and oxidative status in peri-and postmenopausal women. Gynecol Endocrinol. 2016; 32(12): 982-985. <u>https://doi.org/10.1080/09513590.2016.1214257</u>
- 10. Buckwalter J A, Mankin H J, Grodzinsky A J. Articular cartilage and osteoarthritis. Instructional

Baghdad Science Journal

Course Lectures-American Academy of Orthopaedic Surgeons. 2005; 54: 465-480.

- 11. Malfait A M, Miller R E, Miller R J. Basic mechanisms of pain in osteoarthritis: experimental observations and new perspectives. Rheum Dis Clin. 2021; 47(2): 165-180. https://doi.org/10.1016/j.rdc.2020.12.002
- 12. Khajehsaeid H, Abdollahpour Z. Progressive deformation-induced degradation of knee articular cartilage and osteoarthritis. J Biomech. 2020; 111(109995): 1-8. https://doi.org/10.1016/j.jbiomech.2020.109995
- Li Y, Wei X, Zhou J, Wei L. The age-related changes in cartilage and osteoarthritis. Biomed Res Int. 2013; 2013: 1-13. <u>https://doi.org/10.1155/2013/916530</u>
- 14. Hick A C, Malaise M, Loeuille D, Conrozier T, Maugars Y, Pelousse F, et al. Cartilage Biomarkers Coll2-1 and Coll2-1NO2 Are Associated with Knee OA MRI Features and Are Helpful in Identifying Patients at Risk of Disease Worsening. Cartilage. 2021; 13(1_suppl): 1637s-1647s. https://doi.org/10.1177%2F19476035211021892
- 15. Lambert C, Borderie D, Dubuc J-E, Rannou F, Henrotin Y. Type II collagen peptide Coll2-1 is an actor of synovitis. Osteoarth cartil. 2019; 27(11): 1680-1691.

https://doi.org/10.1016/j.joca.2019.07.009

- 16. Hick A-C, Fonck M, Costes B, Cobraiville E, Pirson, S, Garcia L, et al. Serum levels of Coll2-1, a specific biomarker of cartilage degradation, are not affected by sampling conditions, circadian rhythm, and seasonality. Cartilage. 2021; 13(1_suppl): 540S-549S. <u>https://doi.org/10.1177/1947603519878489</u>
- 17. Henrotin Y, Costes B, Malaise M, Loeuille D, Conrozier T, Maugars Y, et al. S-Coll2-1 and s-Coll2-1NO2 sont des marqueurs de la sévérité et de la progression de l'arthrose du genou : résultats de l'étude PRODIGE. Revue du Rhumatisme. 2020; 87(supplement 1): A53. https://doi.org/10.1016/j.rhum.2020.10.083
- 18. Mohammed H A, Ahmed Z A, Alrawi A A O. DC-SIGN Receptor Level in Rheumatoid Arthritis Patients in Baghdad; Serological study. Baghdad Sci

J. 2022; 19(6): 1212. https://doi.org/10.21123/bsj.2022.6559

- Mutar H S, Hasan B F, Muhi S A. Study the Level of Collagen2-1NO2 (Coll2-1NO2) in Serum of Iraqi Patients with Osteoarthritis. Pak J Med. 2022; 16(06): 464-464. <u>https://doi.org/10.53350/pjmhs22166464</u>
- 20. Liu Y, Zhang M, Kong D, Wang Y, Li J, Liu W, et al. High follicle-stimulating hormone levels accelerate cartilage damage of knee osteoarthritis in postmenopausal women through the PI3K/AKT/NFκB pathway. FEBS Open Bio. 2020; 10(10): 2235-2245. https://doi.org/10.1002/2211-5463.12975
- 21. Xu J, Xiao J, Shi Z J. Correlation between age-related serum follicle stimulating hormone levels and osteoarthritis in postmenopausal women. Biomed Res. 2017; 28(13): 5772-5775.
- 22. Santos-Baez L S, Ginsberg H N. Hypertriglyceridemia—causes, significance, and approaches to therapy. Front Endocrinol. 2020; 11(616): 1-7. https://doi.org/10.3389/fendo.2020.00616
- 23. Taay Y, Mohammed M, Abbas R, Ayad A, Mahdi M. Determination of some biochemical parameters in sera of normotensive and hypertensive obese female in Baghdad. J Phys: Conf Ser. 2021; 1853: 012037, IOP Publishing. https://doi.org/10.1088/1742-6596/1853/1/012037

24. Hamza M A, Al Tamer Y Y, Al habib O A. Modification of Irisin Level in Overweight/Obese Women during Pregnancy and Its Association with Some Metabolic Risk Factors. Baghdad Sci J. 2020; 17(3(Suppl.)): 1124. https://doi.org/10.21123/bsj.2020.17.3

- 25. Kadium, T E, Alrubaie, A, Ghanim, S A M. The Link between Serum Omentin Level and Insulin Resistance Biomarkers, Lipid Profile, and Atherogenic Indices in Iraqi Obese Patients, Baghdad Sci J. 2023; 20(1): 0074. https://doi.org/10.21123/bsj.2022.6535
- 26. Peng X, Wu H. Inflammatory links between hypertriglyceridemia and atherogenesis. Curr Atheroscler Rep. 2022; 24(5): 297-306. https://doi.org/10.1007/s11883-022-01006-w



تقييم مستوى Coll2-1NO2 لدى النساء العراقيات المصابات بمرض الفصال العظمي قبل وبعد انقطاع الطمث

نور باسل غانم، بشری فارس حسن

قسم الكيمياء، كلية العلوم للبنات، جامعة بغداد، بغداد، العراق.

الخلاصة

يعد مرض الفصال العظمي (OA) ليس بمرض مناعة ذاتية ولكنه مرض التهابي مزمن يؤثر على مفاصل الأنسان بسبب جهد ميكانيكي وخصوصاً مفاصل الركبة و الحوض. وفي هذا المرض تحدث تغييرات في الغضروف مسببةً ألام في المفاصل خلال الحركة. Coll2-1NO2 هو مصطلح يشير الى سلسلة من الأحماض الأمينية تتكون خلال تحطيم النوع الثاني من الكولاجين والذي يتواجد بشكل وافر في الغضروف، حيث يتم إضافة مجموعة نترات الى الحامض الاميني التايروسين نتيجة لحالة التهابية ناتجة عن الجهد. في هذه الدراسة تم استخدام Coll2-1NO2 كمؤشر حيوي لمرض هشاشة العظام للتنبؤ بحالة الاجهاد الألتهابي وكذلك تحطم الغضروف في النساء قبل وبعد انقطاع الطمث. أظهرت النتائج زيادة ملحوظة في مستوى 2001-2002 في امصال النساء المصابات بمرض هشاشة العظام قبل وبعد انقطاع الطمث بالمقارنة مع النساء السليمات بمستوى العمر ذاته واللاتي يمتلكن مؤشر كتلة المصابات بمرض هشاشة العظام قبل وبعد انقطاع الطمث بالمقارنة مع النساء السليمات بمستوى العمر ذاته واللاتي يمتلكن مؤشر كتلة وسم مماثل (زيادة في الوزن). وأظهر تحليل ارتباط بيرسون عدم وجود علاقة بين 2001-2012 وراقي متغيرات الدراسة. إضافة لذلك، أظهر تحليل عمر المثارة الحاك مؤشر حيوي تم مع النساء السليمات بمستوى العمر ذاته واللاتي يمتلكن مؤشر كلي قبل وبعد انقطاع الطمث المون عدم وجود علاقة بين 2001-2012 في المواس النساء المصابات بمرض هشاشة العظام قبل وبعد انقطاع الطمث بالمقار مع م وجود علاقة بين 2002-2011 وراقي متغيرات الدراسة. إضافة وبد عمر معالي الذلك، أظهر تحليل ارتباط بيرسون عدم وجود علاقة بين 2002-2011 وراقي متغيرات الدراسة. إضافة لذلك، أظهر تحليل ROC حساسية ممتازة جداً لـ2012-2011 كمؤشر حيوي تشخيصي لمرض هشاشة العظام في كل من النساء

الكلمات المفتاحية: نايتر وكو لاجين2-1، البروتين الفعال C، الهرمون المنشط للحوصلة، ملف الدهون، الفصال العظمي.