

## Spectrophotometric Determination of Some Adrenergic Drugs Using H<sub>2</sub>O<sub>2</sub> and Dichlorophenolindophenol

Mohamed Y. Dhamra\*<sup>1</sup>, Mohammed salim Al-Enizzi<sup>2</sup>, Reem A. Al-Luhaiby<sup>3</sup>

<sup>1</sup>Department of Chemistry, College of Education for Pure Science, University of Mosul, Mosul, Iraq.

<sup>2</sup>Department of Chemistry, College of Education for Girls, University of Mosul, Mosul, Iraq.

<sup>3</sup>Department of Medical Laboratory Techniques, Al-Hadba University College, Mosul, Iraq.

\*Corresponding Author.

Received 12/04/2023, Revised 08/09/2023, Accepted 10/09/2023, Published Online First 20/04/2024



© 2022 The Author(s). Published by College of Science for Women, University of Baghdad. This is an Open Access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract


Based on the catalytic response between hydrogen peroxide and dichlorophenolindophenol, a spectrophotometric method was developed to estimate adrenaline hydrochloride (ADR), Salbutamol sulfate (sal), and dopamine hydrochloride (DOPA) in pharmaceutical preparations. A calculated value of the oxidizing agent (hydrogen peroxide) was added to the basic medium in the presence of dichlorophenolindophenol to form a blue color, which suffers from a bleaching process due to the oxidation process of the drugs mentioned above by the oxidizing agent at  $\lambda_{\max}$  552 nm. The estimated amounts of adrenaline hydrochloride and Salbutamol sulfate were between 0.25 - 5  $\mu\text{g/ml}$  and 0.5 - 5  $\mu\text{g/ml}$ , respectively. The amount of dopamine hydrochloride was between 1 - 7  $\mu\text{g/ml}$  with good sensitivity, as the molar absorptance was  $3.8023 \times 10^4$ ,  $8.3506 \times 10^4$  and  $2.7213 \times 10^4$ , respectively. All other statistical values were thoroughly examined. Good agreement with certified value was observed.

**Keywords:** adrenaline hydrochloride, catalytic response, dopamine hydrochloride, salbutamol sulfate, Spectrophotometric.

### Introduction


Adrenaline hydrochloride is 4-(1-hydroxy-2-(methylamino) ethyl) benzene-1,2-diol hydrochloride, Fig. 1. It's a hormone that the adrenal glands secrete that helps the sympathetic nervous system fight or flight<sup>1</sup>. In addition, it speeds up the body's response to stress or threat. It also increases blood flow to the muscles<sup>2</sup> releasing sugar into the bloodstream, along with a series of other effects that make the body alert and better able to fight off the attacker. It is used as a drug in acute asthma by relieving the contraction of bronchial smooth muscles by stimulating beta receptors. Therefore, it reduces the spasm of the pulmonary bronchial tubes and combats bronchospasm, anaphylaxis and anaphylactic shock: thus, preventing the risk of fluid

accumulation in the arteries, and cases of cardiac arrest<sup>3</sup>.



**Figure 1.** The chemical formula of adrenaline hydrochloride


Salbutamol sulfate is bis(4-[2-(tert-butylamino)-1-hydroxyethyl]-2- (hydroxymethyl)phenol); sulfuric acid, Fig. 2, is a medication used to treat asthma and other respiratory conditions such as chronic obstructive pulmonary disease (COPD)<sup>4</sup>. It belongs to a class of drugs called bronchodilators, which work by relaxing the muscles in the airways and allowing more air to flow into the lungs. It is usually taken through an inhaler or nebulizer and can provide quick relief of symptoms such as wheezing, coughing, and shortness of breath. It may also be used to prevent exercise-induced asthma or other triggers that can cause breathing difficulties<sup>5</sup>.



**Figure 2. The chemical formula of Salbutamol sulfate**

Dopamine hydrochloride, or 4-(2-aminoethyl) benzene-1,2-diol; hydrochloride, Fig. 3, is a chemical or hormone found in the human body. It can increase feelings of happiness and acts as a neurotransmitter, sending signals to the body and

brain. Maintaining a proper balance of dopamine hydrochloride in the body is very important as it plays a role in controlling motor skills and emotional responses, which are essential for physical and mental health. Its effects can be found in the major areas of the brain, impacting mood, sleep, focus, learning, and memory. Low dopamine hydrochloride levels can lead to illnesses such as Parkinson's and depression, which can be caused by either a low production of dopamine hydrochloride in the body or a problem with the nerve receptors in the brain<sup>6</sup>.



**Figure 3. The chemical formula of dopamine hydrochloride**

In the chemical literature, there are numerous techniques for measuring adrenaline hydrochloride, Salbutamol sulfate, and dopamine hydrochloride, including spectrophotometric<sup>7-10</sup>, fluorometric<sup>11,12</sup>, electric<sup>13-15</sup> and chromatographic methods<sup>16-18</sup>. The article talks about a straightforward spectrophotometric technique that can be used in water to measure the drugs mentioned previously.

## Materials and Methods

### Apparatus

Absorbance and spectral measurements were done using a PG Instruments-England T92+ UV spectrophotometer using a 1.0 cm thick quartz cell.

### Reagents

Drug solutions were prepared by dissolving 0.01 g of these pure substances in a small amount of distilled water, then completing the volume in a 100ml volumetric flask up to the mark.

### Hydrogen peroxide Solution 0.1 M

Prepared by dissolving 0.9406g of urea-H<sub>2</sub>O<sub>2</sub> compound in 100 ml water, the other oxidizing agents were prepared at a concentration of 0.01 M by dissolving the appropriate amount in a volumetric flask of 50 ml capacity, then filling the volume up to the mark with distilled water. This ought to be freshly prepared each day. Bases stock solutions were prepared with a 1 M concentration, and the

correct amount of each was dissolved in a 100 mL volumetric flask of distilled water for a perfect result and then diluted as needed. All surfactants were prepared at a concentration of 1% by dissolving the optimal amount in a suitable volume and as needed.

### 2,6-Dichloroindophenol Sodium 0.1%

It is prepared by dissolving 0.1g of 2,6-Dichloroindophenol sodium in 100ml distilled water.

### Procedure

Aliquots of solutions containing 0.25–5 µg/ml adrenaline hydrochloride, 0.5-5 µg/ml Salbutamol sulfate, and 1–7 µg/ml dopamine hydrochloride, respectively, were added to three separate groups of 10 ml volumetric flasks, followed by the addition of 1 ml 0.1 M H<sub>2</sub>O<sub>2</sub> and 1.25 ml of 1M NaOH to each group. This was allowed to stand for 10 minutes. Then the optimal amount of dichlorophenolindophenol was added 40 µg/ml and

supplemented with distilled water to the mark. The absorbance was measured versus blank at  $\lambda$  max 552 nm room temperature.


## The Procedure of Pharmaceutical Preparations

### Adrenaline Hydrochloride Injections

The two adrenaline hydrochloride injections each contained 1.0 mg of adrenaline hydrochloride in 1.0 mL. They were diluted in 20 mL of distilled water to create the 100 g/mL solution, which was then withdrawn in various volumes as needed and administered under the method of action.

## Results and Discussion

It is known that a solution of 2,6-Dichlorophenolindophenol produces a blue colour with maximum absorption in the visible region between 500 and 600 nm in the presence of some oxidizing agents. This depends on the nature of the oxidizing agent used and the estimated material. As a result, after oxidizing drugs (adrenaline hydrochloride, salbutamol sulfate, and dopamine hydrochloride) with the same oxidizing agent, this property can be used to evaluate them. The proposed method is considered an indirect method based on quantitatively reducing or bleaching the colour of the 2,6-dichlorophenolindophenol due to the oxidation of the added pharmaceutical compounds<sup>19,20</sup>, scheme 1.



### Scheme 1. The redox reaction between 2,6-Dichlorophenolindophenol

#### Optimum Conditions:

Because the estimation of the drugs under study depends on measuring the optimal amount of dye (2,6-dichlorophenolindophenol) according to Beer's law, the effect of the oxidizing agent, the medium,

### Butadin Tablets

Five butadin tablets, each containing 2 mg, were used in total. They were thoroughly ground before being dissolved in 100 ml of distilled water to produce 100 ppm. The suggested approach to doing the job was followed.

### Dopamine Hydrochloride Injection Analysis

In a 200 ml volumetric flask, distilled water was used to dilute a pharmaceutical injection containing 200 mg/5 ml of dopamine hydrochloride to produce 1000 g/ml. Subsequent samples were then taken from it and handled according to the method.

and stability time on dye formation must be investigated. All preliminary tests were performed in a 10 ml volumetric flask.

### The Effect of Oxidant Amount

Suitable oxidizing agents were chosen to obtain a stable color for the dye. The preliminary study found that hydrogen peroxide is the best. The reason is that the resulting dye was stable, so it was studying the effect of the amount of oxidizing agent 0.1 M on the color of (5  $\mu$ g/ml) dichlorophenolindophenol in the presence of 1ml of each HCl and NaOH (1M). Fig. 4 shows that 1ml of 0.1 M H<sub>2</sub>O<sub>2</sub> was the best volume in the basic medium and was used in the subsequent study.




Figure 4. The effect of oxidant amount.

### The Effect of Type Base

To find the best base that increases the intensity of the colour of the dye, some bases were added in a volume of 1 ml with a concentration of 1 M., Fig. 5. This shows that sodium hydroxide gave the highest intensity and was used in the subsequent study.




Figure 5. The effect of type base.

### The Effect of the Base Volume Used

Since the reaction takes place in an alkaline medium, taking different volumes of the base is appropriate. Figure 6 shows that 1.25 is the most suitable. The pH of the final solution was measured, and its value was 10. Then several buffer solutions with the same pH10 were used, which found that using the base sodium hydroxide was better, so it was used in subsequent studies.




Figure 6. The impact of base volume

### The Effect of Time

To find the best stability time for colour development of the dye (5 µg/ml) after adding the optimal amount of oxidizer (1.0 ml) and 1.25ml of 1M NaOH, the absorbance was measured at contrasting times, and the graph shows that the maximum absorbance was reached after 10 minutes and remained stable for four hours, Fig.7. So, an increasing amount of 2,6-dichlorophenolindophenol was added to obtain the highest value that obeys Beer's law for use in determining the drugs under study. Figs. 8 and 9 show the peak and that 40 µg/mL is the highest.




Figure 7. The effect of time.




Figure 8. The calibration graph of 2,6-dichlorophenolindophenol





Figure 9. The peak of 10 µg/ml of 2,6-dichlorophenolindophenol Vs blank

### The Effect of Oxidation Time on the Drugs

It is expected that when adding the oxidizing agent in the basic medium to adrenaline hydrochloride, salbutamol sulfate, and dopamine hydrochloride

(2µg/ml), an oxidation process will occur, and the remaining oxidizing agent works to show the colour of the dye quantitatively. Therefore, different times for the oxidation process were studied. Fig. 10 indicates that after 10 minutes, the oxidation process was completed, as is evident from ΔA values.




**Figure 10. The effect of oxidation time on the drugs**

#### Effect of Surfactant

Both negative and positive surfactants were used in this study and the result was negative, so they were not used.

#### Calibration Plot

After following the optimal conditions above, it was found that each of the adrenaline hydrochloride, salbutamol sulfate, and dopamine hydrochloride can be estimated Fig. 11. Table 1 shows the limits of the estimate in addition to the values of LOD, LOQ and the rest of the other statistical values.



**Figure 11. Calibration plot of adrenaline hydrochloride (A), salbutamol sulfate (B) and dopamine hydrochloride(C)**

**Table 1. Analytical and statistical values of the drugs under study**

Parameters	Adrenaline hydrochloride HCl	salbutamol sulfate	Dopamine hydrochloride hydrochloride
Linearity range(µg/ml)	0.25-5	0.5-5	1-7
Molar absorptivity (L.mol <sup>-1</sup> . cm <sup>-1</sup> )	3.8023×10 <sup>4</sup>	8.3506×10 <sup>4</sup>	2.7213×10 <sup>4</sup>
Intercept	0.0039	-0.011	-0.0496
Slope	0.1731	0.1448	0.1435
R <sup>2</sup>	0.9990	0.9991	0.9991
LOD* (µg/ml)	0.0421	0.0393	0.0715
LOQ* (µg/ml)	0.1390	0.1297	0.2360

### Method accuracy and Precision

The method was verified by evaluating the accuracy and precision of the method by calculating the relative standard deviation (RSD) of the three

different concentrations. The results presented in Table 2 show that the method has good precision, as the recovery rate reaches 99.69-101.83 and the good agreement (RSD) is less than 3%.

**Table 2. Accuracy and precision of the method**



Drug	Amount taken $\mu\text{g/ml}$	Recovery%	Average recovery %	RSD	Average RSD%
Adrenaline hydrochloride HCl	2	98.45	100.78	2.11	1.88
	3	103.11		1.66	
Salbutamol sulfate s	2	103.35	101.83	2.78	2.31
	3	100.32		1.85	
Dopamine hydrochloride HCl	3	98.16	99.69	1.94	2.2
	5	101.22		2.46	


### Analytical Application

The standard addition methods were applied to pharmaceutical preparations of adrenaline hydrochloride, Salbutamol sulfate, and dopamine hydrochloride as there were no requirements for the three standard methods adopted in the British Pharmacopoeia to demonstrate the lack of spectral overlap in the proposed method. Figs. 12 and Table 3,4 show that the results obtained for adrenaline hydrochloride and dopamine hydrochloride agree well with the original levels of these drugs. This means that the process has a satisfactory selectivity.

**Table 3. The analytical application of the method**

Pharmaceutical preparation	Certified value	Amount present ( $\mu\text{g/ml}$ )	Recovery %
Adrenaline hydrochloride ampoule Egypt	1 mg	2	101.375
		4	96.925
Butadin tablets Iraq	2 mg	2	102.500
		4	98.750
Dopan ampoule Mumbai	200 mg/5ml	4	102.600
		6	98.550





**Figure 12. Standard addition method of adrenaline hydrochloride(A), Salbutamol sulfate (B), and dopamine hydrochloride(C).**

**Table 4. Standard addition method of adrenaline hydrochloride, Salbutamol sulfate, and dopamine hydrochloride summary**

Pharmaceuticals	Certificate value	Amount present ( $\mu\text{g/ml}$ )	Drug content ( $\mu\text{g/ml}$ )	Recovery %
Adrenaline hydrochloride ampoule	1mg/1ml	1	1.010	101.02
Butadin tablets	2 mg	1	0.9566	99.566
Dopan ampoule	200 mg/5ml	2	1.949	97.491

## Conclusion

The current method implements the evaluation of the above-mentioned three drugs with high accuracy and efficiency in the water with short steps in addition to the successful application of the method to a pharmaceutical preparation. It is also possible to

estimate many drugs with a condition and compounds that can suffer from the oxidation process, and the reason is that it depends on the standard curve.

## Acknowledgment

The authors express their appreciation for the staff in the University of Mosul for their support and encouragement throughout this project.

## Authors' Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for republication, which is attached to the manuscript.
- Authors sign on ethical consideration's approval.

- Ethical Clearance: The project was approved by the local ethical committee in Mosul University.
- Ethics statement: No animal studies are present in the manuscript. No human studies are present in the manuscript. No potentially identified images or data are present in the manuscript.

## Authors' Contribution Statement

Each of the individuals listed in the study made an equal contribution to the work and writing.

## References

- Greene BH, Lalonde DH, Seal SK. Incidence of the "adrenaline rush" and vasovagal response with local anesthetic injection. *Plast Reconstr Surg Glob Open* 2021; 9(6):1-5. <https://doi.org/10.1097/GOX.0000000000003659>.
- Filogonio R, Crossley DA. Long term effects of chronic prenatal exposure to hypercarbia on organ

- growth and cardiovascular responses to adrenaline and hypoxia in common snapping turtles. *Comp Biochem Physiol A Mol Integr Physiol* 2019; 234: 10-7. <https://doi.org/10.1016/j.cbpa.2019.04.009>
3. Tschopp C, Tramèr MR, Schneider A, Zaarour M, Elia N. Benefit and harm of adding epinephrine to a local anesthetic for neuraxial and locoregional anesthesia: a meta-analysis of randomized controlled trials with trial sequential analyses. *Anesth Analg*. 2018; 127(1): 228-39. <https://doi.org/10.1213/ANE.0000000000003417>.
  4. Douros K, Moriki D, Sardeli O, Boutopoulou B, Galani A, Papaevangelou V, et al. Assessment and management of asthma exacerbations in an emergency department unit. *Allergol. Immunopathol*. 2023; 51(1): 74-6. <https://doi.org/10.15586/aei.v51i1.720>.
  5. Kruizinga MD, Birkhoff WA, van Esdonk MJ, Klarenbeek NB, Cholewinski T, Nelemans T, et al. Pharmacokinetics of intravenous and inhaled salbutamol and tobramycin: An exploratory study to investigate the potential of exhaled breath condensate as a matrix for pharmacokinetic analysis. *Br J Clin Pharmacol*. 2020; 86(1): 175-81. <https://doi.org/10.1111/bcp.14156>.
  6. Cassidy CM, Zucca FA, Girgis RR, Baker SC, Weinstein JJ, Sharp ME, et al. Neuromelanin-sensitive MRI as a noninvasive proxy measure of dopamine function in the human brain. *Proc Natl Acad Sci*. 2019; 116(11): 5108-5117. <https://doi.org/10.1073/pnas.1807983111>.
  7. Demkiv O, Stasyuk NY, Gayda G, Grynychshyn N, Novikevuch O, Demchuk O, et al. A new spectrophotometric method analysis of adrenaline in pharmaceuticals based on laccase-like nanozymes. *LNUVMB Vet Sci* 2022; 24(106): 142-8. <https://doi.org/10.32718/nvlvet10622>.
  8. Jabar FM, Al-Sabha TaN, Ismael SO. Spectrophotometric Determination of Salbutamol and Terbutaline using 9-Chloroacridine Reagent. *Egypt J Chem*. 2022; 65(2): 61-70. <https://doi.org/10.21608/EJCHEM.2021.80175.3967>.
  9. Al Abdali ZZ, Habeeb NN, Salih ES. Spectrophotometric Determination of Salbutamol Sulphate and Isoxsuprine Hydrochloride in Pharmaceutical Formulations. *Baghdad Sci J*. 2023; 20(2): 0262-. <https://doi.org/10.21123/bsj.2022.6902>.
  10. Hamoudi A T, Spectrophotometric assay of salbutamol sulphate in pharmaceutical preparations by coupling with diazotized p-bromoaniline. *Baghdad Sci J*. 2019; 16(1): 610-615. <https://doi.org/10.21123/bsj.2019.16.3.0610>.
  11. Dhamra MY, Theia'a N, salim Al-Enizzi M. Spectrofluorimetric determination of adrenaline and dopamine. *J Sci Educ*. 2022; 31(3): 17-26. <https://doi.org/10.33899/edusj.2022.133845.1238>.
  12. Pandya HN, Berawala HH, Khatri DM, Mehta PJ. Spectrofluorimetric estimation of salbutamol sulphate in different dosage forms by formation of inclusion complex with  $\beta$ -cyclodextrin. *Pharm methods*. 2010; 1(1): 49-53. [https://doi.org/10.1016/S2229-4708\(10\)11007-3](https://doi.org/10.1016/S2229-4708(10)11007-3).
  13. Khot G, Kaboli M, Celikel T, Shirtcliffe N. Electrochemical detection of adrenaline and hydrogen peroxide on carbon nanotubes. *Surf Innov*. 2022; 10(6): 379-86. <https://doi.org/10.1680/jsuin.21.00038>.
  14. Rajaji U, Chinnapaiyan S, Chen S-M, Govindasamy M, Alothman AA, Alshgari RA. Bismuth telluride decorated on graphitic carbon nitrides based binary nanosheets: Its application in electrochemical determination of salbutamol (feed additive) in meat samples. *J Hazard Mater* 2021; 413: 125265. <https://doi.org/10.1016/j.jhazmat.2021.125265>.
  15. Tran TTT, Doan MD, Dinh QK. Differential pulse voltammetry determination of salbutamol using disulfite tungsten/activated carbon modified glassy carbon electrode. *Chemosphere*. 2022; 303: 135202. <https://doi.org/10.1016/j.chemosphere.2022.135202>.
  16. Britto-Júnior J, Antunes NJ, Campos R, Sucupira M, Mendes GD, Fernandes F, et al. Determination of dopamine, noradrenaline, and adrenaline in Krebs-Henseleit solution by liquid chromatography coupled with tandem mass spectrometry and measurement of their basal release from *Chelonoidis carbonaria* aortae in vitro. *Biomed Chromatogr*. 2021; 35(2): e4978. <https://doi.org/10.1002/bmc.4978>.
  17. Jouyban A, Farajzadeh MA, Khoubnasabjafari M, Jouyban-Gharamaleki V, Mogaddam MRA. Derivatization and deep eutectic solvent-based air-assisted liquid-liquid microextraction of salbutamol in exhaled breath condensate samples followed by gas chromatography-mass spectrometry. *J Pharm Biomed Anal*. 2020; 191: 113572. <https://doi.org/10.1016/j.jpba.2020.113572>.
  18. Chan W. Investigation of the chemical structure and formation mechanism of polydopamine from self-assembly of dopamine by liquid chromatography/mass spectrometry coupled with isotope-labelling techniques. *Rapid Commun Mass Spectrom*. 2019; 33(5): 429-36. <https://doi.org/10.1002/rcm.8373>.
  19. Chang S, Ismail A, Daud Z. Ascorbic acid: properties, determination and uses. In: *Encyclopedia of Food and Health*. Academic Press, London, United Kingdom, 2016 ; 16(1): 275-284. <https://doi.org/10.3390/cosmetics3040038>.
  20. Prodromidis M, Stalikas C, Veltsistas PT, Karayannis M. Spectrophotometric kinetic determination of copper (II) trace amounts based on its catalytic effect on the reaction of the reduced 2, 6-dichlorophenolindophenol and hydrogen peroxide. *Talanta*. 1994; 41(10): 1645-9. [https://doi.org/10.1016/0039-9140\(94\)E0090-E](https://doi.org/10.1016/0039-9140(94)E0090-E).

## التقدير الطيفي لبعض الأدوية الأدرينالية باستخدام $H_2O_2$ وثنائي كلوروفينوليندوفينول

محمد يحيى ضمرة<sup>1</sup>، محمد سالم العنزي<sup>2</sup>، ريم عبد احمد اللهيبي<sup>3</sup>

<sup>1</sup> قسم الكيمياء، كلية التربية للعلوم الصرفة، جامعة الموصل، الموصل، العراق.

<sup>2</sup> قسم الكيمياء، كلية التربية للبنات، جامعة الموصل، الموصل، العراق.

<sup>3</sup> قسم تقنيات المختبرات الطبية، كلية الحداثة الجامعية، الموصل، العراق.

### الخلاصة

بناءً على الاستجابة التحفيزية بين بيروكسيد الهيدروجين وثنائي كلوروفينول اندوفينول، تم تطوير طريقة قياس طيفية لتقدير هيدروكلوريد الأدرينالين (ADR) وكبريتات السالبيتامول (sal) وهيدروكلوريد الدوبامين (DOPA) في المستحضرات الصيدلانية. إذ تمت إضافة كمية محسوبة من العامل المؤكسد (بيروكسيد الهيدروجين) في الوسط القاعدي وبوجود ثنائي كلوروفينول اندوفينول لتكوين صبغة ذات لون أزرق، والتي تعاني من عملية القصر نتيجة لعملية أكسدة الأدوية المذكورة في أعلاه بواسطة العامل المؤكسد (بيروكسيد الهيدروجين) عند طول موجي مقداره 552 نانوميتر. كانت الكميات المقدره من هيدروكلوريد الأدرينالين وكبريتات السالبيتامول بين 0.25 - 5 مايكروغرام / مل و 0.5 - 5 مايكروغرام / مل على التوالي. كانت كمية الدوبامين هيدروكلوريد ما بين 1 - 7 مايكروغرام / مل مع حساسية جيدة حيث أن قيمة الامتصاص المولاري كانت  $10^4 \times 3.8023$  و  $10^4 \times 8.3506$  و  $10^4 \times 2.7213$  على التوالي. تم فحص جميع القيم الإحصائية الأخرى بدقة وتوافق جيدين ومقارنتها مع القيمة الحقيقية.

**الكلمات المفتاحية:** ادرينالين هيدروكلوريد، الاستجابة التحفيزية، القياس الطيفي، دوبامين هيدروكلوريد، كبريتات السالبيتامول.