

Third-Order Differential Subordination for Generalized Struve Function Associated with Meromorphic Functions

Suha J. Hammad^{*1}  , Abdul Rahman S. Juma²  , Hassan H. Ebrahim³  

¹Department of Mathematics, College of Education for Pure Sciences, University of Tikrit, Tikrit, Iraq.

²Department of Mathematics, College of Education for Pure Sciences, University of Anbar, Anbar, Iraq.

³Department of Mathematics, College of Computer Science and Mathematics, University of Tikrit, Tikrit, Iraq.

*Corresponding Author.

Received 23/09/2023, Revised 22/12/2023, Accepted 24/12/2023, Published Online First 20/03/2024,
Published 01/10/2024



© 2022 The Author(s). Published by College of Science for Women, University of Baghdad.

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Previously, many works dealt with the study of the 1st – order differential subordination and shortly after that other studies dealt with the 2nd –order differential subordination in the unit disc. Recently the 3rd – order differential subordination was presented by Antonino and Miller (2011). This paper looks at a considerably broader class of 3rd –order differential inequalities and subordination. The authors define the criteria on an admissible class of operators, implying that 3rd –order differentiale subordination exists. Meromorphic in D is a function that is holomorphic in domain D except for poles. If $D = \mathbb{C}$, it simply states the function is meromorphic. Meromorphic functions in \mathbb{C} are those that may be represented as a quotient of two entire functions. Struve functions have applications in surface –wave and water –wave issues, unstable aerodynamics, optical direction and resistive MHD instability theory. Struve functions have lately appeared in a number of particle systems. The idea of differential subordination in \mathbb{C} is a generalization of differential inequality in \mathbb{R} , and it was initiated in 1981 by the works of Miller, Mocanu and Reade. In this artical, appropriate classes of admissible functions are examined and the properties of 3rd –order differential subordination are established by using the operator $\mathfrak{S}_{c,p}^{\xi,\alpha}$ of meromorphic multivalent functions connected with generalized Struve function. In this study, there is a need to present many concepts including subordination, superordination, the dominant, the best dominant, convolution (or Hadamard product), meromorphic multivalent function, the Struve function addition to the concept of shifted factorial (or Pochhammer symbol) and admissible functions.

Keywords: Admissible Functions, Analytic Function, Convolution (or Hadamard product), Meromorphic Functions, Struve Function, 3rd – Order Differential Subordination.

Introduction

Let $\mathfrak{H}(UD)$ represent the class of analytic functions in $UD = \{z: z \in \mathbb{C} \text{ and } |z| < 1\}$ and let $\mathfrak{H}[a, i]$ be the subclass of analytic functions defined by:

$$\mathfrak{H}[a, i] = \left\{ f \in \mathfrak{H}(UD): f(z) = a + \sum_{\zeta=i}^{\infty} a_{\zeta} z^{\zeta} + \dots \right\}$$

Assume that $\mathfrak{H}[1, i] = \mathfrak{H}_i$.

For two analytic function \check{f} and \check{g} in UD , the function \check{f} is said to be subordinate to the function \check{g} in UD , or the function \check{g} is said to be superordinate to the function \check{f} in UD , and write

$$\check{f}(z) < \check{g}(z) \quad (z \in UD),$$

if there exists a Schwarz function $\varpi(z)$ analytic in UD with $|\varpi(z)| < 1$ and $\varpi(0) = 0$ ($z \in UD$) such that $\check{f}(z) = \check{g}(\varpi(z))$ ($z \in UD$)

It is well known that

$$\check{f}(z) < \check{g}(z) \quad (z \in UD) \Rightarrow \check{f}(0) = \check{g}(0)$$

and $\check{f}(UD) \subseteq \check{g}(UD)$

Furthermore, if the function \check{g} is univalent in UD , the following equivalence hold.

$$\check{f}(z) < \check{g}(z) \quad (z \in UD) \Leftrightarrow \check{f}(0) = \check{g}(0)$$

and $\check{f}(UD) \subseteq \check{g}(UD)$.¹⁻³

Let $\Upsilon(t, u, v, w; z): \mathbb{C}^4 \times UD \rightarrow \mathbb{C}$ and $\check{h}(z)$ be univalent in unit disk UD . If $\mathcal{M}(z)$ is analytic in UD satisfies:

$$\Upsilon(\mathcal{M}(z), z\mathcal{M}'(z), z^2\mathcal{M}''(z), z^3\mathcal{M}'''(z); z) < \check{h}(z). \quad 1$$

Then, $\mathcal{M}(z)$ is the solution to the differential subordination Eq 1 described above. If $\mathcal{M}(z)$ is subordinate to $\mathcal{Q}(z)$ for all $\mathcal{M}(z)$ satisfying Eq 1, the univalent function $\mathcal{Q}(z)$ is said to be a dominant solution of Eq.1. A univalent dominant q satisfying $q < \mathcal{Q}$ for all dominants of Eq 1 is referred to as the best dominant.^{4,5}

Let $\Sigma(p, i)$ be the class of all meromorphic functions of the form:

$$f(z) = z^{-p} + \sum_{\zeta=i}^{\infty} a_{\zeta} z^{\zeta-p} \quad (p, i \in N = \{1, 2, 3, \dots\}), \quad 2$$

which are analytic and p -valent in the punctured disk $(UD)^* = UD \setminus \{0\} = \{z: z \in \mathbb{C} \text{ and } 0 < |z| < 1\}$.

Let $f, l \in \Sigma(p, i)$ where f is given by Eq 2 and l is defined by

$$l(z) = z^{-p} + \sum_{\zeta=i}^{\infty} b_{\zeta} z^{\zeta-p} \quad (z \in (UD)^*), \quad 3$$

the convolution (or Hadamard product) of two functions f and l is defined by

$$(f * l)(z) = z^{-p} + \sum_{\zeta=i}^{\infty} a_{\zeta} b_{\zeta} z^{\zeta-p} = (l * f)(z)$$

In this study, it was considered one of these functions, the series solution of an inhomogeneous second-order Bessel differential equation, which was presented and explored by Struve.⁴ Struve functions and their generalization are used in a variety of areas of applied mathematics and physics.

After a series of mathematical operations, the Struve function of order has the following form⁶:

$$\begin{aligned} \check{U}_{r,b,c}(z) &= 2^r \sqrt{\pi} \Gamma(r + \\ &\frac{b+2}{2}) z^{\frac{-r-1}{2}} \sum_{\zeta=0}^{\infty} \frac{(-c)^{\zeta} (\sqrt{z}/2)^{2\zeta+r+1}}{\Gamma(\zeta+3/2) \Gamma(r+\zeta+(b+2)/2)} = \\ &\sum_{\zeta=0}^{\infty} \frac{(-c/4)^{\zeta}}{(3/2)_{\zeta} (a)_{\zeta}} z^{\zeta}, \quad (z \in UD) \end{aligned}$$

and

$$\check{U}_{a,c}(z) = z \check{U}_{r,b,c}(z) = z + \sum_{\zeta=1}^{\infty} \frac{(-c/4)^{\zeta}}{(3/2)_{\zeta} (a)_{\zeta}} z^{\zeta+1},$$

where $r, b, c \in \mathbb{C}, a = r + (b+2)/2 \neq 0, -1, -2, \dots$ and $(a)_{\zeta}$ is the shifted factorial (or Pochhammer symbol)⁶ expressed terms of the gamma function, by

$$(a)_{\zeta} = \frac{\Gamma(a + \zeta)}{\Gamma(a)} = \begin{cases} 1 & ; (\zeta = 0), \\ a(a+1)(a+2) \dots (a+\zeta-1) & ; (\zeta \in N = \{1, 2, 3, \dots\}) \end{cases}$$

Numerous authors have used the theory of 1st and 2nd order differential subordination and superordination to address a variety of issues in geometric function theory. It is difficult to take the dual issues for higher-order instances into consideration. The idea of 3rd - order differential subordination was first introduced in the works of Ponnusamy and Juneje⁷, but more recently, work by Antonino and Miller⁷ has wekindled interest in this field among researchers. Tang et al⁸. presented the concept of third-order differential subordination

in 2014 as a generalization of the second-order case.

Many studies on the results of the 3rd – order differential subordination and superordination in various contexts have been conducted in recent years.

By setting

$$\delta_{a,c,p}(z) = z^{-p} + \sum_{\zeta=i}^{\infty} \frac{(-c/4)^\zeta}{(3/2)_\zeta (a)_\zeta} z^{\zeta-p}$$

By using the convolution (or Hadamard product), in this work, it is defined a new operator

$$\check{S}_{c,p}^{\zeta,a}: \sum (p, i) \rightarrow \sum (p, i)$$

Which is defined as follows

$$\begin{aligned} \check{S}_{c,p}^{\zeta,a} f(z) &= \delta_{a,c,p}(z) * f(z) \\ &= z^{-p} \\ &\quad + \sum_{\zeta=i}^{\infty} \frac{(-c/4)^\zeta}{(3/2)_\zeta (a)_\zeta} a_\zeta z^{\zeta-p} \quad (z \\ &\in UD), \quad 4 \end{aligned}$$

From Eq 4 it can be easily concluded that

$$\begin{aligned} z \left(\check{S}_{c,p}^{\zeta,a} f(z) \right)' &= (a + \zeta - p) \check{S}_{c,p}^{\zeta,a} f(z) - \\ (a + \zeta) \check{S}_{c,p}^{\zeta,a+1} f(z) \quad 5 \end{aligned}$$

To derive the results of this paper the following lemmas and definitions are needed.

Definition 1:^{5,9} Let \mathcal{Q} be the collection of analytic and univalent functions \mathcal{Q} on $\overline{UD} \setminus E(\mathcal{Q})$, where $E(\mathcal{Q}) = \left\{ \gamma \in \partial UD : \lim_{z \rightarrow \gamma} \mathcal{Q}(z) = \infty \right\}$ and $\min |\mathcal{Q}'(\gamma)| = \partial > 0$ for $\gamma \in \partial UD \setminus E(\mathcal{Q})$. Further, let $\mathcal{Q}(a)$ denote the subclass of \mathcal{Q} consisting of functions \mathcal{Q} for which $\mathcal{Q}(0) = a$ and $\mathcal{Q}(1) \equiv \mathcal{Q}_1$.

Definition 2:⁵ If $\mathcal{N} \subseteq \mathbb{C}, \mathcal{Q} \in \mathcal{Q}$ and $i \geq 2$. Let $\Psi_i[\mathcal{N}, \mathcal{Q}]$ be the family of admissible functions that include functions $\Upsilon: \mathbb{C}^4 \times UD \rightarrow \mathbb{C}$ that satisfy the requirement of acceptability as:

$$\Upsilon(t, u, v, w, ; z) \notin \mathcal{N},$$

when

$$\begin{aligned} t &= \mathcal{Q}(\gamma), u = n\gamma \mathcal{Q}'(\gamma), \\ \operatorname{Re} \left\{ \frac{v}{u} + 1 \right\} &> n \operatorname{Re} \left\{ 1 + \frac{\gamma \mathcal{Q}''(\gamma)}{\mathcal{Q}'(\gamma)} \right\} \end{aligned}$$

and

$$\operatorname{Re} \left\{ \frac{w}{u} \right\} \geq n^2 \operatorname{Re} \left\{ \frac{\gamma^2 \mathcal{Q}'''(\gamma)}{\mathcal{Q}'(\gamma)} \right\},$$

where $z \in UD, \gamma \in \partial UD \setminus E(\mathcal{Q})$ and $n \geq i$.

Lemma 1:⁵ Let $\mathcal{M} \in \mathbb{H}[a, i]$ with $i \geq 2$. Furthermore, let $\mathcal{Q} \in \mathcal{Q}(a)$ and achieve the following requirements:

$$\operatorname{Re} \left\{ \frac{\gamma \mathcal{Q}''(\gamma)}{\mathcal{Q}'(\gamma)} \right\} \geq 0 \text{ and } \left| \frac{z \mathcal{M}'(z)}{\mathcal{Q}'(\gamma)} \right| \leq n,$$

where $z \in UD, \gamma \in \partial UD \setminus E(\mathcal{Q})$ and $n \geq i$. If \mathcal{N} is a set in $\mathbb{C}, \Upsilon \in \Psi_i[\mathcal{N}, \mathcal{Q}]$ and

$$\Upsilon(\mathcal{M}(z), z \mathcal{M}'(z), z^2 \mathcal{M}''(z), z^3 \mathcal{M}'''(z); z) \in \mathcal{N},$$

then $\mathcal{M}(z) \prec \mathcal{Q}(z)$.

Several writers have derived many important conclusions involving numerous operators connected by differential superordination and differential subordination for example^{2,3,10}.

Definition 3: If $\mathcal{N} \subseteq \mathbb{C}$ and $\mathcal{Q} \in \mathcal{Q}_1 \cap \mathbb{H}_i$. Let $\delta_1[\mathcal{N}, \mathcal{Q}]$ be the family of admissible functions, which comprises the functions $\Psi: \mathbb{C}^4 \times UD \rightarrow \mathbb{C}$ that meet the admissibility requirement:

$$\Psi(a', b', c', d'; z) \notin \mathcal{N},$$

when

$$\begin{aligned} a' &= \mathcal{Q}(\gamma), b' = \mathcal{Q}(\gamma) - \frac{1}{a + \zeta} n\gamma \mathcal{Q}'(\gamma), \\ \operatorname{Re} \left\{ \frac{c'(a + \zeta + 1) + a'(a + \zeta) - b'(2a + 2\zeta + 1)}{a' - b'} \right\} \\ &> n \operatorname{Re} \left\{ 1 + \frac{\gamma \mathcal{Q}''(\gamma)}{\mathcal{Q}'(\gamma)} \right\} \end{aligned}$$

and

$$\operatorname{Re} \left\{ \frac{3c'(a + \zeta + 1)(a + \zeta) - a'(5a^2 + 5\zeta^2 + 10a\zeta - 3a - 3\zeta) - d'(a + \zeta + 2)(a + \zeta + 1) - 3b'(a + \zeta)(a + \zeta - 2)}{a' - b'} \right\} \geq n^2 \operatorname{Re} \left\{ 1 + \frac{\zeta^2 \mathcal{Q}'''(\zeta)}{\mathcal{Q}'(\zeta)} \right\},$$

where $z \in UD, \zeta \in \partial UD \setminus E(\mathcal{Q})$ and $n \geq i \geq 2$.

Results and Discussion

Theorem 1: Let $\Omega \subseteq \mathbb{C}$ and $\Psi \in \delta_1[\Omega, \mathcal{Q}]$. If $f \in \Sigma(p, i)$ and $\mathcal{Q} \in \mathcal{Q}_1$ achieve the following requirements:

$$\operatorname{Re} \left\{ \frac{\zeta \mathcal{Q}''(\zeta)}{\mathcal{Q}'(\zeta)} \right\} \geq 0 \text{ and } \left| z \left(z^p \check{S}_{c,p}^{\zeta,a} f(z) \right)' \right| \leq n |\mathcal{Q}'(\zeta)|, \quad 6$$

then

$$\left\{ \Psi \left(z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z) \right); z \right\} \subset \Omega \quad 7$$

which leads to

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) \prec \mathcal{Q}(\zeta).$$

Proof: In unit disk UD , define

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) = \mathcal{M}(z) \quad (z \in UD). \quad 8$$

To differentiate equation Eq 8 with respect to z and using Eq 5 and its recurrence,

$$z^p \check{S}_{c,p}^{\zeta,a+1} f(z) = \mathcal{M}(z) - \frac{1}{a+\zeta} z \mathcal{M}'(z) \quad 9$$

Once again to differentiate equation Eq 9 through applying the recurrence relation Eq 5 and with regard to z ,

$$z^p \check{S}_{c,p}^{\zeta,a+2} f(z) = \mathcal{M}(z) - \frac{2}{a+\zeta+1} z \mathcal{M}'(z) + \frac{1}{(a+\zeta+1)(a+\zeta)} z^2 \mathcal{M}''(z) \quad 10$$

Additional calculations show that

$$z^p \check{S}_{c,p}^{\zeta,a+3} f(z) = \mathcal{M}(z) - \frac{3}{(a+\zeta+2)} z \mathcal{M}'(z) + \frac{3}{(a+\zeta+2)(a+\zeta+1)} z^2 \mathcal{M}''(z) - \frac{1}{(a+\zeta+2)(a+\zeta+1)(a+\zeta)} z^3 \mathcal{M}'''(z) \quad 11$$

Let

$$\left. \begin{aligned} a' &= t, & b' &= t - \frac{1}{a+\zeta} u, & c' &= t - \frac{2}{a+\zeta+1} u + \frac{1}{(a+\zeta+1)(a+\zeta)} v, \\ d' &= t - \frac{3}{(a+\zeta+2)} u + \frac{3}{(a+\zeta+2)(a+\zeta+1)} v - \frac{1}{(a+\zeta+2)(a+\zeta+1)(a+\zeta)} w \end{aligned} \right\} \quad 12$$

$$\frac{1}{(a+\zeta+2)(a+\zeta+1)(a+\zeta)} w \quad 13$$

Now, the transformation is defined

$\Upsilon(t, u, v, w; z): \mathbb{C}^4 \times UD \rightarrow \mathbb{C}$ by

$$\Upsilon(t, u, v, w; z) = \Psi(a', b', c', d'; z) = \Psi \left(t, t - \frac{1}{a+\zeta} u, t - \frac{2}{a+\zeta+1} u + \frac{1}{(a+\zeta+1)(a+\zeta)} v, t - \frac{3}{(a+\zeta+2)} u + \frac{3}{(a+\zeta+2)(a+\zeta+1)} v - \frac{1}{(a+\zeta+2)(a+\zeta+1)(a+\zeta)} w \right)$$

And by making use of the equations Eq 8 – Eq 11

$$\Upsilon(\mathcal{M}(z), z \mathcal{M}'(z), z^2 \mathcal{M}''(z), z^3 \mathcal{M}'''(z); z) = \Psi \left(z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z); z \right) \quad 14$$

Also, note that

$$\frac{v}{u} + 1 = \frac{c'(a+\zeta+1)+a'(a+\zeta)-b'(2a+2\zeta+1)}{a'-b'}$$

$\frac{w}{u}$

$$= \frac{3c'(a+\zeta+1)(a+\zeta) - a'(5a^2 + 5\zeta^2 + 10a\zeta - 3a - 3\zeta) - d'(a+\zeta+2)(a+\zeta+1) - 3b'(a+\zeta)(a+\zeta-2)}{a'-b'}$$

Theorem 1 is proved by relying on Lemma 1 because the requirements of admissibility for function $\Psi \in \delta_1[\Omega, \mathbb{Q}]$ of Definition 3 is equivalent to the requirements of admissibility for function $\Upsilon \in \Psi_\zeta[\Omega, \mathbb{Q}]$ in Definition 2.

When the function $\mathbb{Q}(z)$ has an unknown behavior on ∂UD , the following results will be an extension of Theorem 1.

Corollary 1: If $\Omega \subseteq \mathbb{C}$ and \mathbb{Q} is univalent in UD with $\mathbb{Q} \in \mathcal{Q}_1$. Let $\Psi \in \delta_1[\Omega, \mathbb{Q}_\partial]$ for some $\partial \in (0,1)$, where $\mathbb{Q}_\partial(z) = \mathbb{Q}(\partial z)$. If $f \in \Sigma(p, i)$ and \mathbb{Q}_∂ , satisfies the following condition:

$$Re \left\{ \frac{\Upsilon_{\mathbb{Q}_\partial}(z)}{\mathbb{Q}_\partial'(z)} \right\} \geq 0 \text{ and } \left| z \left(z^p \check{S}_{c,p}^{\zeta,a} f(z) \right)' \right| \leq n |\mathbb{Q}_\partial'(z)|, \quad 15$$

then

$$\left\{ \Psi \left(\begin{matrix} z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), \\ z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z); z \end{matrix} \right) : z \in UD \right\} \subset \Omega,$$

which implies

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < \mathbb{Q}(z).$$

Proof: According to Theorem 1

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < \mathbb{Q}_\partial(z),$$

and since $\mathbb{Q}_\partial(z) < \mathbb{Q}(z)$, then $z^p \check{S}_{c,p}^{\zeta,a} f(z) < \mathbb{Q}(z)$.

If $\Omega \neq \mathbb{C}$ is a family-connected domain, there is a conformal mapping $h: UD \rightarrow \mathbb{C}$ such that $h(UD) = \mathbb{C}$. The class $\delta_1[h(UD), \mathbb{Q}]$ is then denoted as

and

$\delta_1[h, \mathbb{Q}]$. The following two corollaries are direct implications of Theorem 1 as well as Corollary 1.

Corollary 2: If h is a univalent function in UD and let $\Psi \in \delta_1[h, \mathbb{Q}]$, assume that $\mathbb{Q} \in \mathcal{Q}_1$ satisfies Eq 6. Then,

$$\left\{ \Psi \left(\begin{matrix} z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), \\ z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z); z \end{matrix} \right) : z \in UD \right\} < h(z), \quad 16$$

which implies

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < \mathbb{Q}(z).$$

Corollary 3: If \mathbb{Q} is a univalent function in UD with $\mathbb{Q} \in \mathcal{Q}_1$ and $\Psi \in \delta_1[h, \mathbb{Q}_\partial]$ for some $\partial \in (0,1)$, where $\mathbb{Q}_\partial(z) = \mathbb{Q}(\partial z)$. If \mathbb{Q}_∂ meets the requirements of Eq 15, then the subordination relation Eq 16 implies that

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < \mathbb{Q}(z).$$

The next corollary explains the connection between the solution of the related 3^{rd} - order differential equation and the best dominant of a 3^{rd} -order differential subordination.

Corollary 4: If h is a univalent function in UD and Υ is given by Eq 14 where $\Psi \in \delta_1[h, \mathbb{Q}]$. If the differential equation

$$\Upsilon(\mathbb{Q}(z), z\mathbb{Q}'(z), z^2\mathbb{Q}''(z), z^3\mathbb{Q}'''(z); z) = h(z)$$

has a solution \mathbb{Q} with $\mathbb{Q} \in \mathcal{Q}_1$ that meets the requirements Eq 6, then subordination Eq 16 implies that

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < \mathbb{Q}(z),$$

and \mathcal{Q} is the best dominant of Eq 16.

Proof: Since

$$\Psi \left(\begin{array}{l} z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), \\ z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z); z \end{array} \right)$$

$$= \Upsilon(\mathcal{M}(z), z\mathcal{M}'(z), z^2\mathcal{M}''(z), z^3\mathcal{M}'''(z); z)$$

$$< \check{h}(z), \quad 17$$

then $\mathcal{M}(z)$ is a solution of Eq 17, and Corollary 2

$$\mathcal{M}(z) < \mathcal{Q}(z),$$

that is \mathcal{Q} is a dominant of Eq 17. So is,

$$\Psi \left(\begin{array}{l} z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), \\ z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z); z \end{array} \right)$$

$$\Psi \left(1 + \mathcal{B}e^{i\theta}, 1 + \frac{(a + \zeta - n)\mathcal{B}e^{i\theta}}{a + \zeta}, 1 + \frac{\mathcal{L} + (a + \zeta)[(a + \zeta + 1) - 2n]\mathcal{B}e^{i\theta}}{(a + \zeta + 1)(a + \zeta)}, 1 + \frac{3\mathcal{L}(a + \zeta) - \mathcal{N} + (a + \zeta + 1)(a + \zeta)[(a + \zeta + 2) - 3n]\mathcal{B}e^{i\theta}}{(a + \zeta + 2)(a + \zeta + 1)(a + \zeta)}; z \right) \notin \Omega,$$

whenever $z \in UD$, $Re\{\mathcal{L}e^{-i\theta}\} \geq n(n-1)\mathcal{B}$ and $Re\{\mathcal{N}e^{-i\theta}\} \geq 0$ for every $0 \leq \theta \leq \pi$ and $n \geq 2$. the following result can be derived from Theorem 1 using the concept of the family of admissible functions.

Corollary 5: If $\Omega \subseteq \mathbb{C}$ and $\delta_1[\Omega, \mathcal{B}]$. If

$$\left| z \left(z^p \check{S}_{c,p}^{\zeta,a} f(z) \right)' \right| \leq n\mathcal{B},$$

then

$$\Psi \left(\begin{array}{l} z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), \\ z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z); z \end{array} \right) \in \Omega \quad (z \in UD),$$

which leads to

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < 1 + \mathcal{B}e^{i\theta}.$$

$$= \Upsilon(\mathcal{M}(z), z\mathcal{M}'(z), z^2\mathcal{M}''(z), z^3\mathcal{M}'''(z); z)$$

$$< \check{h}(z)$$

$$= \Upsilon(\mathcal{Q}(z), z\mathcal{Q}'(z), z^2\mathcal{Q}''(z), z^3\mathcal{Q}'''(z); z),$$

that is \mathcal{Q} is the best dominant of Eq 17.

When $\mathcal{Q}(z) = 1 + \mathcal{B}e^{i\theta}$, $\mathcal{B} > 0$, by applying Theorem 1. According to Definition 3, the family of admissible functions $\delta_1[\Omega, \mathcal{Q}]$ is now denoted by $\delta_1[\Omega, \mathcal{B}]$ as follows:

Definition 4: If $\Omega \subseteq \mathbb{C}$ and let $\mathcal{B} > 0$. The family of admissible functions $\delta_1[\Omega, \mathcal{B}]$, which includes the functions $\Psi: \mathbb{C}^4 \times UD \rightarrow \mathbb{C}$ that satisfy the admissibility condition:

Applications Involving $\check{S}_{c,p}^{\zeta,a}$

If it is assumed that $\Omega = \{\tau \in \mathbb{C}: |\tau - 1| < \mathcal{B}\}$ and $\delta_1[\Omega, \mathcal{B}]$ is simply indicated by $\delta_1[\mathcal{B}]$, then Corollary 5 is reduced to the next corollary.

Corollary 6: Let $\Psi \in \delta_1[\mathcal{B}]$ and suppose that

$$\left| z \left(z^p \check{S}_{c,p}^{\zeta,a} f(z) \right)' \right| \leq n\mathcal{B},$$

then

$$\left| \Psi \left(\begin{array}{l} z^p \check{S}_{c,p}^{\zeta,a} f(z), z^p \check{S}_{c,p}^{\zeta,a+1} f(z), \\ z^p \check{S}_{c,p}^{\zeta,a+2} f(z), z^p \check{S}_{c,p}^{\zeta,a+3} f(z); z \end{array} \right) - 1 \right| < \mathcal{B} \quad (z \in UD),$$

which implies

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < 1 + \mathcal{B}z.$$

Corollary 7: Suppose that

$$\left| z \left(z^p \check{S}_{c,p}^{\zeta,a} f(z) \right)' \right| \leq nB,$$

then

$$\left| z^p \check{S}_{c,p}^{\zeta,a+1} f(z) - 1 \right| < B \quad (z \in UD),$$

which lead to

$$z^p \check{S}_{c,p}^{\zeta,a} f(z) < 1 + Bz.$$

Proof:

If it is put $\Psi(a', b', c', d'; z) = b'$, Corollary 6 leads to the conclusion.

If it is taken $c = -4$ and $(a)_\zeta = \frac{1}{(3/2)_\zeta}$ in Corollary 7, then

$$\check{S}_{-4,p}^{\zeta,a} f(z) = f(z)$$

$$\text{and } \check{S}_{-4,p}^{\zeta,a+1} f(z) = \frac{(a+\zeta-p)}{(a+\zeta)} f(z) - \frac{1}{(a+\zeta)} z f'(z)$$

The following results will be produced:

Example 1: If $f \in \Sigma(p, i)$ satisfies the following conditions

Conclusion

The purpose of using the operator $\check{S}_{c,p}^{\zeta,a}$, is to provide some results for the 3^{rd} – order differential subordination for analytic function. Investigating pertinent classes of admissible function leads to the results. The findings presented in this paper offer

Authors' Declaration

- Conflicts of Interest: None.

Authors' Contribution Statement

This work was carried out in collaboration with all authors. A.S. did the diagnosis of the cases then collected the samples, and did the tests. S.J. wrote and edited the manuscript with revised ideas. A.S. and H.H. conducted the analysis of the data with

References

$$\left| z \left(z^p \check{S}_{c,p}^{\zeta,a} f(z) \right)' \right| \leq nB$$

and

$$\left| \frac{(a+\zeta-p)}{(a+\zeta)} z^p f(z) - \frac{1}{(a+\zeta)} z^{p+1} f'(z) - 1 \right| < B,$$

then

$$\left| z^p f(z) - 1 \right| < B.$$

In addition, if it is taken $c = -4$ and $(a)_\zeta = \frac{1}{(3/2)_\zeta}$ in Corollary 7 and $p = i = 1$ the following results will be obtained:

Example 2: If $f \in \Sigma(p, i)$ satisfies the following inequality

$$\left| z^2 f'(z) + z f(z) \right| \leq nB$$

and

$$\left| \frac{(a+\zeta-1)}{(a+\zeta)} z f(z) - \frac{1}{(a+\zeta)} z^2 f'(z) - 1 \right| < B,$$

then

$$\left| z f(z) - 1 \right| < B.$$

fresh perspectives for further research and opportunities are provided for scholars to generalize the results to produce new results in the fields of meromorphic univalent and meromorphic multivalent function theory.

- Ethical Clearance: The project was approved by the local ethical committee at University of Tikrit.

revised ideas. All authors read and approved the final manuscript.

1. Cotîrlă L-I, Juma ARSJA. Properties of differential subordination and superordination for multivalent functions associated with the convolution operators. 2023;12(2):169. <https://doi.org/10.3390/axioms12020169>.
2. Çetinkaya A, Cotîrlă L-I. Briot–Bouquet differential subordinations for analytic functions involving the Struve function. Fractal and Fractional. 2022;6(10):540. <https://doi.org/10.3390/fractalfract6100540>.
3. Farzana HA, Jeyaraman MP, Bulboaca T. On Certain Subordination Properties of a Linear Operator. J Fract Calc. 2021; 12(2): 148-62. .
4. Toklu EJHJoM, Statistics. Radii of starlikeness and convexity of generalized Struve functions. 2020:1-18.: <https://doi.org/10.15672/hujms.518154> .
5. Shehab NH, Juma ARS. Third Order Differential Subordination for Analytic Functions Involving Convolution Operator. Baghdad Sci. J. 2022;19(3):0581-592. <https://doi.org/10.21123/bsj.2022.19.3.0581>.
6. Muhey UD, Srivastava HM, Mohsan RJHJoM, Statistics. Univalence of certain integral operators involving generalized Struve functions. 2018; 47(4): 821-33.
7. Ponnusamy S, Juneja O P. Third-order differential inequalities in the complex plane. Current Topics in Analytic Function Theory. 1992: 274-90. https://doi.org/10.1142/9789814355896_0023.
8. Tang H, Srivastava HM, Li S-H, Ma L-N, editors. Third-order differential subordination and superordination results for meromorphically multivalent functions associated with the Liu-Srivastava operator. Abstract and Applied Analysis; 2014: Hindawi. <https://doi.org/10.1155/2014/792175>.
9. Al-Khafaji TKM. Strong Subordination for E-valent Functions Involving the Operator Generalized Srivastava-Attiya. Baghdad Sci J. 2020; 17(2): 0509-514. <https://doi.org/10.21123/bsj.2020.17.2.0509>
10. Seoudy T. Second order differential subordination and superordination of Liu-Srivastava operator on meromorphic functions. Afr Mat. 2021; 32(7-8): 1399-408. <https://doi.org/10.1007/s13370-021-00907-4> .

التبعية التفاضلية من الدرجة الثالثة لدالة ستروف المعممة المرتبطة بالدوال الميرومورفية (الدوال التحليلية باستثناء عدد منته من النقاط)

سهى جمعة حماد¹، عبدالرحمن سلمان جمعة²، حسن حسين ابراهيم³

¹ قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة تكريت، تكريت، العراق.

² قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة الانبار، الرمادي، العراق.

³ قسم الرياضيات، كلية علوم الحاسوب والرياضيات، جامعة تكريت، تكريت، العراق.

الخلاصة

سابقاً تناولت العديد من الاعمال دراسة التبعية التفاضلية من الدرجة الاولى وبعد فترة وجيزة تناولت دراسات اخرى التبعية التفاضلية من الدرجة الثانية في قرص الوحدة . مؤخراً تم تقديم التبعية التفاضلية من الدرجة الثالثة من قبل بوانسامي واخرون في عام 1992 و انتونيو ميلر عام 2011 . تبحث هذه الورقة في صنف اوسع بكثير من مترجمات التبعية التفاضلية من الدرجة الثالثة . عرف المؤلفون المعايير الخلصة بصنف من الاوبريترات المسموح بها وهذا يعني ضمان وجود التبعية التفاضلية من الدرجة الثالثة. الدوال الميرومورفية في D هي دوال تحليلية في المجال D باستثناء الرواسب اذا كانت $D = \mathbb{C}$ فان الدالة ميرومورفية وهي دوال يمكن تمثيلها كحاصل قسمة دالتين. دالة ستروف لها تطبيقات في قضايا الموجات السطحية و موجات الماء والديناميكا الهوائية غير المستقرة و الاتجاه البصري ونظرية عدم الاستقرار المقاوم ظهرت دالة ستروف مؤخرًا في عدد من أنظمة الجسيمات . فكرة التبعية التفاضلية في لغة \mathbb{C} هي تعميم للمتباينات في لغة \mathbb{R} , وقد بدأت في عام 1981 من خلال اعمال ميلر و موكانو و ريد. في هذا المقال يتم معاينة الفئات المناسبة من الدوال المقبولة ويتم انشاء خصائص التبعية التفاضلية من الدرجة الثالثة باستخدام المؤثر $S_{\zeta, p}^{\alpha}$ للدوال متعددة التكافؤ التحليلية باستثناء عدد منته من النقاط (الاقطاب) المرتبطة بوظيفة ستروف المعممة . في هذه الدراسة هناك حاجة لعرض العديد من المفاهيم منها التبعية , الفوقية , المسيطر , افضل الساند , الالتواء (او منتج هادامارد) , الدالة متعددة التكافؤ التحليلية باستثناء عدد منته من النقاط (الاقطاب) , دالة ستروف بالاضافة الى مفهوم المضروب المزاح (او رمز بوشهامر) و الدوال المسموح بها.

الكلمات المفتاحية: الدوال المسموح بها، الدالة التحليلية، الالتفاف، الدالة التحليلية باستثناء عدد منته من النقاط (الاقطاب)، دالة ستروف، التبعية التفاضلية من الدرجة الثالثة.