

On Monotonically T₂-spaces and Monotonically normal spaces

*Radhi I.M.Ali**

*Jalal Hatem Hussein**

Date of acceptance 22/9/2008

Abstract

In this paper we show that if $\prod X_i$ is monotonically T₂-space then each X_i is monotonically T₂-space, too. Moreover, we show that if $\prod X_i$ is monotonically normal space then each X_i is monotonically normal space, too. Among these results we give a new proof to show that the monotonically T₂-space property and monotonically normal space property are hereditary property and topologically property and give an example of T₂-space but not monotonically T₂-space.

Keywords: topological space, continuous function, monotonically T₂-spaus, monotonically normal space.

1. Introduction

The property of monotonically T₂-space first appeared by R.E. Buck, some weaker monotone separation and basis properties are presented in [1]. Unfortunately Buck's definition is not precise so we give a precise definition of monotonically T₂-space and find several properties of such concept and other topological concepts. In order to make this work as self-contained as possible we give the following lemmas.

Lemma 1.1: [6] Let X, Y is two topological spaces and $f: X \rightarrow Y$ is a closed injective function then f is open function.

Proof: Let W be any open set in X , $X \setminus W$ is closed set. Hence $f(X \setminus W)$ is closed set in Y . Since f is injective $f(X \setminus W) = Y \setminus f(W)$ which implies that $f(W)$ is open set in Y . Therefore f is open function \square

Lemma 1.2: [6] Let X, Y is two topological spaces and $f: X \rightarrow Y$ is a continuous closed injective function and let B be a subset of Y then

$$f^{-1}(f(B)) = B$$

Proof: Since f is continuous we have $f^{-1}(f(B)) \subseteq B$.

Let $x \in B$, i.e. $f(x) \in f(B)$. If W is any open set in X containing x , then $f(W)$ is open set in Y containing $f(x)$ which implies that $f(W) \cap f(B) \neq \emptyset$ and have $W \cap f^{-1}(f(B)) \neq \emptyset$, therefore $x \in f^{-1}(f(B)) \square$

2. Monotonically T₂-space

Let (X, τ_X) be a topological space, we begin with precise definition of monotonically T₂-space.

Definition 2.1: Let $\Delta = \{(x, x) : x \in X\}$ and denote by $(X \times X)^* = (X \times X) \setminus \Delta$

A topological space X is monotonically T₂-space if there is a function

$$g : (X \times X)^* \rightarrow \tau_X$$

* Department of mathematics- College of Science for Women- Baghdad University

Assigning to each order pair (x,y) in $(X \times X)^*$ an open neighborhood $g(x, y) \subset X$ of x such that

- (i) $g(x, y) \cap g(y, x) = \phi$,
- (ii) For each subset M of X and if $x \in \overline{U\{g(y, x) : y \in M\}}$ then $x \in \overline{M}$.

We will call such a function g monotone T_2 -operator on X . Of course, every monotonically T_2 -space is T_2 -space, but later we find an example of T_2 -space but not monotonically T_2 -space.

Theorem 2.2: If (X, τ_X) is a topological space, (Y, τ_Y) is a topological space which is monotonically T_2 -space and $f: X \rightarrow Y$ continuous closed injective function then X is a monotonically T_2 -space.

Proof: Let g' be monotone T_2 -operator on Y , i.e.

$$g': (Y \times Y)^* \rightarrow \tau_Y$$

Satisfies (i) and (ii) of definition (2.1). In order to define a monotone T_2 -operator on X , let $(x,y) \in (X \times X)^*$ then $(f(x), f(y)) \in (Y \times Y)^*$ because f is injective. So there is an open neighborhood $g'(f(x),f(y)) \subset Y$ of $f(x)$. Since f is continuous from X into Y $f(g'(f(x),f(y)))$ is an open neighborhood of x subset of X .

Define $g: (X \times X)^* \rightarrow \tau_X$ as follows $g(x,y) = f^{-1}(g'(f(x),f(y)))$ for all $(x,y) \in (X \times X)^*$.

We note that

$$g(x,y) \cap g(y,x) = f^{-1}((g'(f(x),f(y))) \cap f^{-1}(g'(f(y),f(x))))$$

$$= f^{-1}(g'(f(x),f(y)) \cap g'(f(y),f(x)))$$

$$= f^{-1}(\phi) = \phi.$$

Let M be any subset of X and $x \in \overline{U\{g(y,x) : y \in M\}}$ i.e. $x \in \overline{U\{f^{-1}(g'(f(y),f(x))) : y \in M\}} = f^{-1}(U\{g'(f(y),f(x)) : y \in M\})$ $\therefore x \in f^{-1}(U\{g'(f(y),f(x)) : y \in M\})$ which implies that

$f(x) \in U\{g'(f(y),f(x)) : y \in M\}$. i.e. $f(x) \in U\{g'(f(y),f(x)) : f(y) \in f(M)\}$. Hence $f(x) \in \overline{f(M)}$ because Y is monotonically T_2 -space.

$\therefore f(x) \in \overline{f(M)}$ because f is closed function, which implies $x \in M$ because f is injective function. Thus X is monotonically T_2 -space and g is monotone T_2 -operator on X . \square

Corollary 2.3: If topological space X is a homeomorphic to monotonically T_2 -space Y then X is monotonically T_2 -space.

Proof: Since a Homeomorphism function is a continuous closed injective function then the result can be deduced from theorem 2.2. \square

Theorem 2.4 [2]: If the topological space is monotonically T_2 -space then X is regular space.

We are going to give an example of T_2 -space but not monotonically T_2 -space.

Example 2.5 [6]: Let X denote the closed interval $[0, 1]$ and D the subset $\{1/n : n=1, 2, 3, \dots\}$. Define on X the smallest topology which contains every open set of $X \setminus \{0\}$ as a subspace of the real line \mathbb{R} and every set B_a , $(0 < a \leq 1)$, defined by $B_a = \{t \in X : t < a \text{ and } t \notin D\}$. The space X is T_2 -space but not regular which implies it is not monotonically T_2 -space.

Theorem 2.6[2]: If X is a monotonically T_2 -space and A is a subspace of X then A is monotonically T_2 -space.

In the following theorem we are going to show that if the product $X \times Y$ is monotonically T_2 -space then X and Y are monotonically T_2 -spaces.

Theorem 2.7: Let X, Y be two topological spaces. If the product space $X \times Y$ is monotonically T_2 -space then X, Y are monotonically T_2 -spaces.

Proof: For each fixed $y \in Y$, $X \times \{y\}$ is a subspace of $X \times Y$ then by theorem 2.6 $X \times \{y\}$ is a monotonically T_2 -space but $X \times \{y\}$ homeomorphic to space X hence, by corollary 2.3, X is monotonically T_2 -space. Using same argument we can show that Y is monotonically T_2 -space, too. \square
Use mathematical induction and apply theorem 2.7 we can get the following result \square

Corollary 2.8: Let X_i be indexed family of topological spaces. If the product space $\prod X_i$ is monotonically T_2 -space then each X_i is monotonically T_2 -space.

3. Monotonically Normal spaces

The property of monotone normality first appears without name, in Lemma 2.1 of C.R. Borge's paper on startifiable spaces [3]. In [4], Zenor and others gave properties and characterizations of monotonically normal spaces.

Definition 3.1 A T_1 -space X is monotonically normal if there is a function G which assigns to each order pair (H, K) of disjoint closed subsets of X an open set $G(H, K)$ such that
(i) $H \subset G(H, K) \subset G(H, K) \subset X \setminus K$
(ii) If (H', K') is a pair of disjoint closed subsets of X such that $H \subset H'$ and $K \supset K'$, then $G(H, K) \subset G(H', K')$.
The function G is called a monotone normality operator for X .

Lemma 3.2: [4] Any monotonically normal has a monotone normality operator G satisfying $G(H, K) \cap G(K, H) = \emptyset$ for any pair (H, K) of closed sets. Furthermore, each of the

following properties is equivalent to monotone normality of a space X :

(a) There is a function G which assigns to each order pair (S, T) of separated subsets of X an open set $G(S, T)$ satisfying

(i) $S \subset G(S, T) \subset G(S, T) \subset X \setminus T$.

(ii) If (S', T') is a pair of separated sets having $S \subset S'$ and $T \supset T'$ then $G(S, T) \subset G(S', T')$.

(b) There is a function F which assigns to each order pair (p, C)

with C closed set and $p \in X \setminus C$, an open set $F(p, C)$

satisfying

(i) $p \in F(p, C) \subset X \setminus C$.

(ii) If D is closed and $C \supset D$ then $F(p, C) \subset F(p, D)$.

(iii) If $p \neq q$ are points of X then $F(p, \{q\}) \cap F(q, \{p\}) = \emptyset$.

Remarks 3.3: [4] (a) The property described in lemma 3.2 (a) was originally called complete monotone normality and because condition (i) in (a) the space is T_5 -space. Hence the space which is normal but it is not T_5 -space is a normal space but not monotonically normal space, see [5].

Theorem 3.4: If X is monotonically normal and A is a closed subspace of X then A is monotonically normal space, too.

Proof : Let G be a monotone normality operator for X and H_1, K_1 be any two disjoint closed sets in A , so $H = H_1 \cap A, K = K_1 \cap A$ are two disjoint closed sets in X . Hence there is an open set $G(H, K)$ in X , we can define a function G_1 assigns to each order pair (H_1, K_1) an open set $G_1(H_1, K_1)$ in A such that $G_1(H_1, K_1) = G(H, K) \cap A$,

It is easy to show that G_1 is a monotone normality operator for A . \square

Theorem 3.5: If (X, τ_X) is a topological space, (Y, τ_Y) is a

monotone normal space with G is a monotone normality operator for Y and $f: X \rightarrow Y$ continuous closed injective function then X is a monotonically normal space, too.

Proof: Let H, K be any two disjoint closed sets in X . Since f is closed and injective function, $f(H), f(K)$ are two disjoint closed sets in Y and hence there is an open set $G(f(H), f(K))$ in Y assigns with $(f(H), f(K))$. $f(G(f(H), f(K)))$ is an open set in X therefore we can define a function G' assigns with H, K an open set $G'(H, K)$ in X as follows

$$G'(H, K) = f(G(f(H), f(K)))$$

Since G is a monotone normality operator in Y and f is closed we obtain from

$$f(H) \subset G(f(H), f(K)) \subset G(f(H), f(K)) \subset Y \setminus f(K),$$

$$H \subset f(G(f(H), f(K))) \subset f(G(f(H), f(K))) = f(G(f(H), f(K))) \subset f(Y \setminus f(K)) = X \setminus K.$$

Moreover, let (H', K') be a pair of disjoint closed sets in X having

$H \subset H'$ and $K \supset K'$. Since f is closed and injective function we have $f(H) \subset f(H')$ and $f(K) \supset f(K')$. Hence $G(f(H), f(K)) \subset G(f(H'), f(K'))$ implies $f(G(f(H), f(K))) \subset f(G(f(H'), f(K')))$ i.e. $G'(H, K) \subset G'(H', K')$

$\therefore G'$ is a monotone normality operator for X \square

Corollary 3.6: The monotonically normal property of space is topological property.

Proof: Homeomorphism is continuous closed injective function and use theorem 3.5 \square

In the following theorem we are going to show that if the product space $X \times Y$ is a monotonically normal space then X, Y are monotonically normal spaces.

Theorem 3.7: let X and Y be two topological spaces. If the product space $X \times Y$ is monotonically normal space then X, Y are monotonically normal spaces.

Proof: For each fixed $y \in Y$, $X \times \{y\}$ is closed subspace of $X \times Y$ and then by theorem 3.4 $X \times \{y\}$ is monotonically normal space. But $X \times \{y\}$ homeomorphic to space X hence X is monotonically normal space. By using same argument we can show that Y is monotonically normal space \square

Use mathematical induction and apply theorem 3.7 we can get the following result.

Corollary 3.8: Let X_i indexed family of topological spaces. If the product space $\prod X_i$ is monotonically normal space then X_i is monotonically normal space.

References:

1. Buck, R.E., 1996 Some weaker monotone separation and basis properties, *Topology and its Applications*. 69:1-12.
2. Buck, R.E., 1993 Monotone separation and basis properties, Ph.D. Thesis, University of Pittsburgh, Pittsburgh, PA.
3. Borges, C.J.R., 1966 On startifiable spaces, *Pacific J.Math.* 17:1-16.
4. Heath, R.W., Lutzer D.J. and Zenor P.L., 1973 Monotonically normal spaces, *Trans, Amer, Math. Soc.* 178: 481-493.
5. Shkerin, S.A. 2004. *Topology and its Applications*. 136:129-134.
6. Steen, L. A. and Seebach J. A. Jr., 1970. *Counterexamples in Topology*. Holt, Rinehart and

Winston, Inc., New York,
210pp.

7. Hu, S.T., 1969 Element of
general topology, 3rd Ed,

Holaen. Day, Inc, California,
214pp.

الفضاءات T_2 -الرتيبية والفضاءات العادية الرتيبية

جلال حاتم حسين*

راضي إبراهيم محمد*

* جامعة بغداد - كلية العلوم للبنات - قسم الرياضيات

الخلاصة:

أثبتنا في هذا البحث انه اذا كان $\prod X_i$ فضاءاً T_2 رتيباً فأن كل من X_i فضاء T_2 رتيب اضافة الى ذلك اثبتنا انه اذا كان $\prod X_i$ فضاءاً عادياً رتيباً فأن كل من X_i فضاء عادي رتيب ايضاً. من بين هذه النتائج اعطينا اثباتاً جديداً بأن خاصية كون الفضاء T_2 رتيب بانها خاصية وراثية وتوبولوجية وأن خاصية كون الفضاء الرتيب العادي خاصية وراثية وتوبولوجية واعطينا مثلاً لفضاء T_2 لكنه ليس فضاء T_2 رتيباً.