

# Preparation and characterization and biological activity of new complexes for some transition elements with Schiff bases derived from one antipyrine substitute

Zeena U. Jasim , Amal M. Ali , Heba A. Mohammed , Maher A. Mohammed\* 

Department of Chemistry, Education College for Girls, Mosul University, Mosul, Iraq.

\*Corresponding Author.

Received 27/09/2023, Revised 07/01/2024, Accepted 09/01/2024, Published Online First 20/10/2024



© 2022 The Author(s). Published by College of Science for Women, University of Baghdad.

This is an open access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

A group of divalent complexes was prepared in the formula  $[M(L)Cl_2]$  when  $M = Co^{+2}, Ni^{+2}, Cu^{+2}, Zn^{+2}$ , and  $Cd^{+2}$   $L = (Z)-4-((1-(H-indol-3-yl)ethylidene)amino)1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one$ . The Schiff base ligand is prepared by reacting 4-aminoantipyrine with 3-acetylindol. The prepared ligand and complexes were identified by elemental analysis, an infrared spectrometer, the nuclear magnetic resonance spectrum, molar electrical conductivity, electronic spectra, magnetic susceptibility, atomic absorption of the elements, and biological efficiencies. The measurements showed that the synthesized ligand interacts with the central metal ion through the oxygen carbonyl group and nitrogen azomethine atoms, that the coordination of the prepared complexes has a square planar nickel complex and the tetrahedral geometry of other complexes, and that these complexes are non-electrolytic. The synthesized compounds have good antibacterial activity against Gram-negative bacteria, represented by *Pseudomonas savastanoi*, and Gram-positive bacteria, represented by *Bacillus subtilis*.

**Keywords:** 4-aminoantipyrin substitute, Biological activity, Schiff base ligand, Square planar complexes, Transition metals.

## Introduction

Schiff bases are chemical substances that are typically created by the condensation of an aldehyde or ketone with a primary amine and typically occur under acid, base, or heat catalysis<sup>1, 2</sup>. 4-Aminoantipyrine (AAP), a pyrazolone derivative, has drawn a lot of interest from medicinal chemists since it has so many potential uses. pharmacological effects such as analgesic, anti-inflammatory, antiviral, and antibacterial. Two nitrogen atoms, the carbonyl and amine groups, which are present in the cyclic structure of AAP, have a significant impact on its biological action. As donor ligands, the carbonyl and amine groups of AAP help to create different coordination assemblies and chelate structures. The study of AAP derivative complexes is fascinating

since the development of the organic-metal chelate results in significant changes to the compound's biological characteristics<sup>3-6</sup>. The novel complexes with L ligands were synthesized in this research and investigated using a variety of spectroscopic methods. These complexes have been prepared by the reaction of many transition metal salts with Schiff bases. Mononuclear complexes with different stoichiometry and structure were characterized. This type of produced complex was found depending on the reaction medium. Moreover, the complexes antibacterial effectiveness was investigated using Hinton Mueller agar medium<sup>7,8</sup>. In this work, we aim to prepare some transition metal complexes (II) with Schiff base ligands and characterize them to show

probable applications for these complexes in the future.

## Materials and Methods

The Fluka or Merck companies supplied all of the reagent-grade chemicals utilized in this experiment. Without any further purification, they were used totally as they were. Using the Stuart-SMP30 melting point apparatus, the melting point temperature was calculated. The atomic absorption spectrophotometer of type Shimadzu used spectral methods to determine the elemental contents. Lovibond Con200 was used to test the molar conductance in DMF ( $10^{-3}$  M) at 25 °C. Using KBr pellets at 400 to 4000  $\text{cm}^{-1}$ , Shimadzu FTIR was used to record the IR spectra of the ligand and its complexes. UV/visible spectra measurements were made by the T80 UV/VIS Spectrometer. Magnetic susceptibility measurements were made by the magnetic susceptibility balance of Sherwood Scientific, Cambridge, UK. Using deuterated DMSO as a solvent, the  $^1\text{H}$  NMR was captured at 400 MHz

on the BioSpin GmbH shield. The agar and bacteria were supplied by the department of biology at the College of Science at the University of Mosul.

### Synthesis of Ligand (L)

In 30 mL of hot ethanol, dissolve 4-aminoantipyrine (6.09 g, 0.03 mol) and a hot solution of 3-acetylindol (4.77 g, 0.03 mol) and mix. Glacial acetic acid was then added in small amounts to the mixture and put under reflux for 3 hours. The resulting solid was purified by recrystallization, filtered, washed with ether, and then dried <sup>9, 10</sup>. The orange Schiff base product yield: 85%, m.p. 68–70 °C, Anal. Calc. for  $\text{C}_{21}\text{H}_{20}\text{N}_4\text{O}$ : C; 73.25; H; 5.81; N; 16.27; found: C; 73.02; H; 5.68; N; 16.07%. The synthesis of Schiff base ligand (L) is shown in Fig. 1.




Figure 1. Synthesis of Schiff base ligand (L)

### Complexes' synthesis

The following general process <sup>11, 12</sup> was used to prepare each complex: Schiff base ligand (L) (1.72 g, 0.005 mol), dissolved in ethanol, was added to a heated solution of metal(II)chloride ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  (1.19g,0.005mol),  $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$  ( 1.18g,0.005mol),

$\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$  ( 0.85,0.005mol),  $\text{ZnCl}_2$  ( 0.68,0.005mol ) and  $\text{CdCl}_2$  (0.91,0.005mol)) in a mole ratio of 1:1 L:M. For six hours, the mixture was refluxed, and as the complexes cooled, they were filtered out, washed with ethanol and ether, and then dried.

## Results and Discussion

The metal (II) complexes in this investigation were soluble in DMF and other solvents but insoluble in water and stable in air. Various spectroscopic techniques revealed the structure of the produced compound. Elemental analysis (C.H.N.), as well as melting points, supported the predicted molecular composition of the compound. The molar

conductance of the prepared complexes was measured at a concentration of  $10^{-3}$  M using DMF solvent, which gave a conductivity ranging between 12–22  $\text{Ohm} \cdot \text{cm}^2 \cdot \text{mol}^{-1}$  which is consistent with non-electrolyte complexes<sup>13</sup>. Characterization, analytical, and molar conductance data of the complexes listed in table1.

**Table 1. Data on the characteristics, analysis, and molar conductance of complexes**

no.	(Formula)	(Color)	Yield %	(m.p) (°C)	$(\Delta_M)$ $\text{Ohm}^{-1} \cdot \text{cm}^2 \cdot \text{mol}^{-1}$	(calculate(found)%)			
						(C%)	(H%)	(N%)	(M%)
1	[Co(L)Cl <sub>2</sub> ]	Gray	67	179-181	12	53.18 (53.07)	4.22 (4.12)	11.81 (11.54)	12.43 (12.19)
2	[Ni(L)Cl <sub>2</sub> ]	Brownish Yellow	94	159-160	17	53.14 (52.68)	4.22 (4.10)	11.82 (11.54)	12.39 (12.12)
3	[Cu(L)Cl <sub>2</sub> ]	Brown	86	112-113	22	52.67 (52.43)	4.18 (3.89)	11.70 (11.57)	13.28 (13.12)
4	[Zn(L)Cl <sub>2</sub> ]	Red	64	117-119	18	52.47 (52.26)	4.16 (3.86)	11.66 (11.31)	13.61 (13.38)
5	[Cd(L)Cl <sub>2</sub> ]	Gray	90	166-168	30	47.79 (47.26)	3.79 (3.41)	10.62 (10.30)	21.31 (21.11)


### Infrared Spectra

The spectra of the ligand showed the recognizable bands at 1620  $\text{cm}^{-1}$ , 1730  $\text{cm}^{-1}$ , 3163  $\text{cm}^{-1}$ , 2927  $\text{cm}^{-1}$ , and 3116  $\text{cm}^{-1}$ , and the bands were referred to as  $\nu$  C=N azomethine,  $\nu$  C=O carbonyl,  $\nu$  N-H,  $\nu$  C-H aliphatic, and  $\nu$  C-H aromatic, respectively<sup>14,15</sup>. In all the metal complexes, the stretching frequencies of the azomethine and carbonyl groups shifted

downward, which indicated that the N and O atoms were involved in coordination. The spectra of IR of the complexes referred to new bands in 510–455  $\text{cm}^{-1}$ , 565–509  $\text{cm}^{-1}$  regions, which can be assigned to  $\nu$  (M–N) and  $\nu$  (M–O) vibrations<sup>16-19</sup>. Ligand and their complexes' selected IR bands  $\text{cm}^{-1}$  listed in Table 2 and Fig. 2 show the FTIR spectra of the ligand (L).

**Table 2. Ligand and their complexes' selected IR bands  $\text{cm}^{-1}$**

Compound no.	$\nu$ (N-H)	$\delta$ (N-H)	$\nu$ (C=N)	$\nu$ (C=O)	$\nu$ (N-N)	$\nu$ (M-N)	$\nu$ (M-O)
L	3163	1238	1620	1730	1016	–	–
1	3166	1240	1571	1614	1020	509	556
2	3168	1240	1596	1616	1020	510	565
3	3165	1240	1523	1618	1022	459	555
4	3161	1241	1521	1616	1021	459	509
5	3167	1240	1520	1616	1020	455	557



**Figure 2. FTIR spectra of the ligand (L)**

### Electronic Spectra and Magnetic Moment Measurement


The transitions  $\pi \rightarrow \pi^*$  and  $n \rightarrow \pi^*$  are attributed to the bands in the 37750, 33444  $\text{cm}^{-1}$  range of the

ligand's electronic spectra.<sup>20</sup> The Co(II) d<sup>7</sup> complex's electronic spectra exhibit absorption bands at 17182 cm<sup>-1</sup> due to (ν<sub>3</sub>), which is attributed to the transition <sup>4</sup>A<sub>2</sub> (F)→<sup>4</sup>T<sub>1</sub>(P) and 4.46 B.M. practical magnetic moment values of the cobalt complex, which is higher than the theoretical value of 3.82 B.M. due to the orbital contribution<sup>21</sup>. The nickel (II) d<sup>8</sup> complex has electronic spectrum bands that are ascribed to transitions <sup>1</sup>A<sub>1g</sub> → <sup>1</sup>A<sub>2g</sub> (ν<sub>1</sub>) and <sup>1</sup>A<sub>1g</sub> → <sup>1</sup>B<sub>1g</sub> (ν<sub>2</sub>), 20618 cm<sup>-1</sup> and 24213 cm<sup>-1</sup>, respectively, and a dia magnetic moment that was measured<sup>22</sup>. In Cu(II) d<sup>9</sup> complexes, the band at 15290 cm<sup>-1</sup> is due to <sup>2</sup>T<sub>2</sub> (D) → <sup>2</sup>E (P), as well as the practical magnetic moment of the Cu(II) complex of 2.12 B.M., which is higher than the theoretical value


of 1.73 B.M. due to the second-order orbital contribution<sup>23</sup>. These results were compatible with the tetrahedral structure of the Co(II) and Cu(II) ions, while the Ni(II) complex supports the square-planar structure. The magnetic moment measurements of the zinc (II) d<sup>10</sup> and cadmium (II) d<sup>10</sup> complexes showed that they are diamagnetic, and the peaks in the electronic spectra are attributed to charge transfer transitions and suggest that the metal ions are surrounded by a tetrahedral structure<sup>24,25</sup>. Table 3 shows the electronic transition and magnetic data of the ligand and complexes, and Figs. 3 and 4 show the UV spectra of the ligand and complex [Ni(L)Cl<sub>2</sub>], respectively.

**Table 3 . Electronic transition and magnetic moment for metal complexes and Ligand**

Number	Compounds	λ <sub>max</sub> (nm)	Band (cm <sup>-1</sup> )	Assignments	magnetic moment	Structures
	ligand (L)	265	37750	π → π*	--	--
		299	33444	n → π*		
1	[Co(L)Cl <sub>2</sub> ]	582	17182	<sup>4</sup> A <sub>2</sub> (F) → <sup>4</sup> T <sub>1</sub> (P)	4.46	<b>Tetrahedral</b>
		339	29498	C.T		
2	[Ni(L)Cl <sub>2</sub> ]	485	20618	<sup>1</sup> A <sub>1g</sub> → <sup>1</sup> A <sub>2g</sub>	Dia	<b>Squer planar</b>
		413	24213	<sup>1</sup> A <sub>1g</sub> → <sup>1</sup> B <sub>1g</sub>		
		340	29411	C.T		
3	[Cu(L)Cl <sub>2</sub> ]	654	15290	<sup>2</sup> T <sub>2</sub> → <sup>2</sup> E	2.12	<b>Tetrahedral</b>
		329	30395	C.T		
4	[Zn(L)Cl <sub>2</sub> ]	305	32786	C.T	--	<b>Tetrahedral</b>
5	[Cd(L)Cl <sub>2</sub> ]	303	33033	C.T	--	<b>Tetrahedral</b>



**Figure3. Electronic Spectrum of ligand (L)**



**Figure4. Electronic Spectrum of [Ni(L) Cl<sub>2</sub>]**

### <sup>1</sup>H-NMR Spectra

The <sup>1</sup>H-NMR spectrum of the prepared ligand (L) Fig. 6 gave the following signals: -NH at 11.9 ppm, -N=C-CH<sub>3</sub> at 2.13 ppm, =C-CH<sub>3</sub> at 2.74 ppm, -N-N-CH<sub>3</sub> at 3.84 ppm, 7.1–8.3 ppm due to the aromatic proton, These signs did not change in the prepared complexes, which confirms their steric structure for

the prepared compound<sup>26-28</sup>. <sup>1</sup>H-NMR spectrum of the ligand (L) is shown in Fig. 5.




Figure 5. <sup>1</sup>H-NMR Spectrum of the ligand (L)




Figure 6. The structure of the prepared complexes a [M(L)Cl<sub>2</sub>] b [Ni(L)Cl<sub>2</sub>]

### Biological activity of the prepared compounds

The bacteriological activity of the prepared ligand and its metal complexes was conducted on two types of bacteria that are pathogenic to humans. The first type was Gram-negative bacteria, represented by *Pseudomonas savastanoi*, and the second type was Gram-positive bacteria, represented by *Bacillus subtilis*. The solutions in our study were prepared at a concentration of 10 mg/ml dissolved in the solvent (DMSO), and 100 microliters were placed inside the holes in the culture medium. The dishes were closed with a special tape, and the dishes were placed inside the incubator for 24 hours at a temperature of 37 °c, after which a ruler was used to measure the diameter of inhibition to show that the sensitivity of the studied compounds depending on the diameter of the

inhibition. One type of antibiotic (Ciprodar) was used. The prepared ligand showed higher effectiveness against both types of bacteria than the prepared complexes. The reason for this is the presence of alkyl groups and donor atoms within the structure of the active ligand, which led to the formation of complexes with the elements present inside the cell that the bacterial cell needs, such as cobalt and copper, and the loss of these elements leads to cell death. The prepared complexes also showed good biological activity, and the reason for this may be attributed to the coordination of surrounding the cell wall that contains a high percentage of fats the ligand with the central metal ion through the nitrogen and oxygen donor atoms. Bacterial cells are resistant to some chemical compounds because of the presence of a thick membrane<sup>29-31</sup>. The antibacterial activity of the ligands and their complexes (10 µg/1 ml) is shown in Tables 4 and Figs. 7 and 8.

Table 4. Antibacterial activity of the ligands and their complexes (10 µg/1 ml)

prepared compound	<i>Bacillus subtilis</i>	<i>Pseudomonas savastanoi</i>
Control	20	12
L	14	15
[Co(L)Cl <sub>2</sub> ]	10	11
[Ni(L)Cl <sub>2</sub> ]	8	10
[Cu(L)Cl <sub>2</sub> ]	13	14
[Zn(L)Cl <sub>2</sub> ]	10	12
[Cd(L)Cl <sub>2</sub> ]	9	10






Figure 7. Differences in biological activity against bacteria for ligand and complexes

## Conclusion

Through several physicochemical, Electronic Spectra, Infrared Spectra, <sup>1</sup>H-NMR spectra, and magnetic susceptibilities measurements, it was found that the ligand is bidentate and it is linked through the nitrogen and oxygen donor atoms to the central metal ion and completes the coordination number with a chlorine ion to become the tetrahedral geometry of the cobalt(II), copper(II), zinc(II) and

## Acknowledgment

The researchers expressed their thanks and appreciation to the Chemistry Department, College of Education for Girls, University of Mosul, for




Figure 8. Inhibitory region of ligands and complexes against bacteria

cadmium (II) complexes, while the nickel(II) complex is a square planar geometry. The prepared ligand showed higher activity than the complexes against the two types of bacteria used in this research, which may be due to its ligand structure containing azomethine and carbonyl groups and its ability to coordinate with metal ions.

providing the necessary facilities for the research work.

## Authors' Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been included with the necessary permission for re-publication, which is attached to the manuscript.
- No animal studies are present in the manuscript.
- No human studies are present in the manuscript.
- No potentially identified images or data are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee at University of Mosul.

## Authors' Contribution Statement

The role of first authors Z.U. J. Prepared ligands and complexes, did IR test and data analysis, The role of second authors A. M. A. (UV visible and <sup>1</sup>H NMR tests , measure molarity conductivity and study biological activity) for the complexes, the role of the

third author H. A. M. writing the manuscript, drafting and designing the Ms and the forth author M.A. M. Conception, analysis, revision and proofreading, acquisition of data, data analysis .

## References

- 1- Wannas NM, Al-Hamdani AAS, Al Zoubi, W. Spectroscopic characterization for new complexes with 2, 2'-(5, 5-dimethylcyclohexane-1, 3-diyliidene) bis (azan-1-yl-1-ylidene) dibenzoic acid. *J Phys Org Chem.* 2020; 33(11): 1-12. <https://doi.org/10.1002/poc.4099>
- 2- Bozejewicz D, Osmiałowski B, Kaczorowska MA, Witt K. 2, 6-Bis ((benzoyl-R) amino) pyridine (R= H, 4-Me, and 4-NMe<sub>2</sub>) Derivatives for the Removal of Cu (II), Ni (II), Co (II), and Zn (II) Ions from Aqueous Solutions in Classic Solvent Extraction and a Membrane Extraction. *Membranes.* 2021; 11(4): 1-9. <https://doi.org/10.3390/membranes11040233>
- 3- Al-Shemary RKR, Alkam HH, AlWan WM. Implementations of Transition Metal Complexes of Schiff Base Derived Ampyrone. *Der Pharm Lett.* 2021; 13(6): 17-48. <https://2u.pw/LFjCdrS>.
- 4- Rasyda YA, Widowati MK, Marliyana SD, Rahardjo SB. Synthesis, Characterization and Antibacterial Properties of Nickel (II) Complex with 4-Aminoantipyrine Ligand. *Indones J Chem.* 2021; 21(2): 391-399. <https://doi.org/10.22146/ijc.56552>
- 5- Ibraheem IH, Sadiq AS, Al-Tameemi M, Alias MF. Synthesis, Spectral Identification, Antibacterial Evaluation and Theoretical Study of Co, Fe, Rh and Pd Complexes for 2-benzoylthiobenzimidazol. *Baghdad Sci J.* 2022; 19(6): 1326-1326. <https://dx.doi.org/10.21123/bsj.2022.6704>
- 6- Ghosh P, Dey SK, Ara MH, Karim R, Islam ABM. A review on synthesis and versatile applications of some selected Schiff bases with their transition metal complexes. *Egypt J Chem.* 2019; 62(Special Issue (Part 2) Innovation in Chemistry: 523-547. <https://dx.doi.org/10.21608/ejchem.2019.13741.1852>
- 7- Ashrafuzzaman MD, Camellia FK, Mahmud AA, Pramanik MD, Nahar K, HAQUE M. Bioactive mixed ligand metal complexes of Cu (II), Ni (II), and Zn (II) ions: synthesis, characterization, antimicrobial and antioxidant properties. *J Chil Chem Soc.* 2021; 66(3): 5295-5299 <http://dx.doi.org/10.4067/S0717-97072021000305295>
- 8- Hamza IS, Mahmmod WA, Al-Hamdani AA, Ahmed SD, Allaf AW, Al, et al. Synthesis, characterization, and bioactivity of several metal complexes of (4-Amino-N-(5-methyl-isaxazol-3-yl)-benzenesulfonamide. *Inorg Chem Commun.* 2022; 144(109776): 1-23. <https://doi.org/10.1016/j.inoche.2022.109776>
- 9- Janabi ZMA, Alsalami HSJ, Al-Khafaji ZS, Hussien SA. Increasing of the corrosion resistance by preparing the trivalent nickel complex. *Egypt J Chem.* 2022; 65(6): 193-198. <https://doi.org/10.21608/ejchem.2021.100733.4683>
- 10- Nabhan KJ, Mahdi AS, Al-Zaidi BH, Ismail AH, Nasif ZN. New tetra-dentate schiff base ligand N<sub>2</sub>O<sub>2</sub> and its complexes with some of metal ions: preparation, identification, and studying their enzymatic and biological activities. *Baghdad Sci J.* 2022; 19(1): 0155-0155. <http://dx.doi.org/10.21123/bsj.2022.19.1.0155>
- 11- Hashim DJ, Mahdi SM. Preparation, Characterization, and Biological Study of New Halogenated Azo-Schiff Base Ligands and Their Complexes. *J Med Chem. Sci.* 2022; 6: 1555-1576. <https://doi.org/10.26655/JMCHEMSCI.2023.7.8>
- 12- Al-Shaheen AJ, Al-Bayati MR, Ali AM. Synthesis and characterization some of metal (II) complexes with 2-(6-methoxy naphthalene-2-yl) propionic acid (2-hydroxy benzyliidene) hydrazone. *Int J Enhance Sci Technol Eng.* 2017; 6(11): 10-16.

- 13- Chandra S, Raizada S, Tyagi M, Sharma PK. Spectroscopic and biological approach of Ni (II) and Cu (II) complexes of 2-pyridinecarboxaldehyde thiosemicarbazone. *Spectrochim Acta A Mol Biomol Spectrosc.* 2008; 69(3): 816-821. <https://doi.org/10.1016/j.saa.2007.05.033>.
- 14- Alattar RA, Hassan ZM, Abass SK, Ahmad LM. Synthesis, characterization and study the photodecolorization of Schiff base Fe (III) complex in ZnO/Uv-A light system. *AIP Conf Proc.* 2020; 2290(1): 030032-10. <https://doi.org/10.1063/5.0027611>
- 15- Numan,AT, Atiyah EM, Fayad AA, Namuq IB. Synthesis, Characterization and Study of Antibacterial Activity of a New Schiff Base Ligand and Its Complexes with Co (II), Ni (II), Cu (II), Cd (II) and Hg (II) Metal Ions. *Baghdad Sci J.* 2016; 13(2): 0153-0153. <http://dx.doi.org/10.21123/bsj.2016.13.2.2NCC.0351>
- 16- Saranya J, Kirubavath SJ, Chitra S, Zarrouk A, Kalpana K, Lavany K, et al. Tetradentate Schiff Base complexes of transition metals for antimicrobial activity. *Arab J Sci Eng.* 2020; 45: 4683-4695. <https://doi.org/10.1007/s13369-020-04416-7>
- 17- Shaalan N, Preparation, Spectroscopy, Biological Activities and Thermodynamic Studies of New Complexes of Some Metal Ions with 2-[5-(2-Hydroxy-Phenyl)-1, 3, 4-Thiadiazol-2-Ylimino]-Methyl-Naphthalen-1-Ol. *Baghdad Sci J.* 2022; 19(4): 829-837. <http://dx.doi.org/10.21123/bsj.2022.19.4.0829>
- 18- Jamil W, Solangi S, Ali M, Taha KMKM, Khuhawar M Y. Syntheses, characterization, in vitro antiglycation and DPPH radical scavenging activities of isatin salicylhydrazidehydrazone and its Mn (II), Co (II), Ni (II), Cu (II), and Zn (II) metal complexes. *Arabian Journal of Chemistry.* 2019; 12(8):2262-2269 <http://dx.doi.org/10.1016/j.arabjc.2015.02.015>
- 19- Ali A, Elasala GS, Mohamed EA, Kolkaila SA. Spectral, thermal studies and biological activity of pyrazinamide complexes. *Heliyon.* 2019; 5(11): 1-10. <https://doi.org/10.1016/j.heliyon.2019.e02912>
- 20- Reda S M, Al-Hamdani A A S. Mn (II), Fe (III), Co (II) and Rh (III) complexes with azo ligand: Synthesis, characterization, thermal analysis and bioactivity. *Baghdad Sci J.* 2023; 20(3): 642-660. <https://dx.doi.org/10.21123/bsj.2022.7289>
- 21- Kareem MJ, Al-Hamdani AAS, Ko YG, Al Zoubi W, Mohammed SG. Synthesis, characterization, and determination antioxidant activities for new Schiff base complexes derived from 2-(1H-indol-3-yl)-ethylamine and metal ion complexes. *J Mol Struct.* 2021; 1231: 129669. <https://doi.org/10.1016/j.molstruc.2020.129669>
- 22- Al-Hassani R A. Synthesis, Characterization, Antimicrobial and Theoretical Studies of V (IV), Fe (III), Co (II), Ni (II), Cu (II), and Zn (II) Complexes with Bidentate (NN) Donor Azo Dye Ligand. *Baghdad Sci J.* 2016; 13(4): 793-805. <http://dx.doi.org/10.21123/bsj.2016.13.4.0793>
- 23- Sabah AA, Ameen AM, Al-Daher A. Metal Complexes of Bis (2, 6-diamine pyridine 2, 5-hexanedione) Macrocyclic Schiff-Base Ligand: Preparation, Characterization and thermal study. *Iraqi J Sci.* 2022; 63(5): 1885-1893. <https://doi.org/10.24996/ij.s.2022.63.5.3>
- 24- Al-Hamdani AA, Shaalan ND, Bakir SR, Mohammed MS. Preparation, spectral characterization, structural study, and evaluation of antibacterial activity of Schiff base complexes for VOII, CrIII, MnII, ZnII, CdII and CeIII. *Baghdad Sci J.* 2015; 12(2): 350-361.
- 25- Al-qasii NAR, Ali TB, Zaied M. Synthesis and Characterization of a Novel Azo-Dye Schiff Base and Its Metal Ion Complexes Based on 1, 2, 4-Triazole Derivatives. *Indones J Chem.* 2023; 23(6): 1555-1566. <https://doi.org/10.22146/ijc.83509>
- 26- Ali AM, Mohammed HA, Jasim ZU, Abed RR. Characterization and Antibacterial Evaluation of New Complexes of Nicotinamide semicarbazone Manganese (II), Zinc (II), Silver (I) Synthesizes. *Res J Pharm Technol.* 2023; 16(2): 539-544. <https://doi.org/10.52711/0974-360X.2023.00092>
- 27- Tyagi M, Chandra S, Akhtar J, Chand D. Modern spectroscopic technique in the characterization of biosensitive macrocyclic Schiff base ligand and its complexes: Inhibitory activity against plantpathogenic fungi. *Spectrochim Acta A Mol Biomol Spectrosc.* 2014; 118: 1056-1061. <https://doi.org/10.1016/j.saa.2013.09.124>
- 28- Dakore SD ,Kamble VT. synthesis of Biological Active N<sub>2</sub>O<sub>2</sub> Type of Navel Schiff base Metal Complexes Derived from 4- Amino Antipyrin using Tea. *J Chem Pharm Res.* 2016; 8(11): 211-215.
- 29- Sherif SH, Kure DA, Moges EA, Argaw B. Synthesis., Characterization, and Antibacterial, Activity, Evaluation of 4-amino Antipyrine. Derivatives and Their Transition Metal Complexes. *Am J Biosci Bioeng.* 2021; 9(1): 8-12. <https://doi.org/10.11648/j.bio.20210901.12>
- 30- Mohanambal D, Sridev G, Arul antony S, Angayarkani R. review: a recent review in applications of heterocyclic compounds as antimicrobial agent. *Asian J Pharm Clin Res.* 2018; 11(12): 66-74. <https://doi.org/10.22159/ajpcr.2018.v11i12.26890>
- 31- Kareem IK, Waddai FY, Abbas G J. Synthesis, Characterization and biological activity of Some Transition Metal Complexes with New Schiff Base Ligand Type (NNO) Derivative from Benzoin. *J Pharm Sci Res.* 2019; 11(1): 119-124.



## تحضير وتشخيص ودراسة الفعالية البيولوجية لمعقدات جديدة لبعض العناصر الانتقالية مع قواعد شيف مشتقة من احدى معوضات الأنتي بايرين

زينة أسامة جاسم ، أمل محسن علي، هبة عزيز محمد، ماهر عبد الرزاق محمد

قسم الكيمياء، كلية التربية للبنات، جامعة الموصل، الموصل، العراق.

### الخلاصة

حضرت مجموعة من المعقدات ذات الصيغة  $[M(L)Cl_2]$  حيث  $M = Co^{+2}, Ni^{+2}, Cu^{+2}, Zn^{+2}, Cd^{+2}$  من ليكاند قاعدة شيف مشتق من المركب 4-امينو انتيبيرين مع 3-استايل اندول كما تم تشخيص الليكاند والمعقدات المحضرة بالتحليل الدقيق للعناصر وأطياف الأشعة تحت الحمراء وطيف الرنين النووي المغناطيسي وقياسات الحساسية المغناطيسية وتحليل المحتوى الفلزي بطريقة الامتصاص الذري فضلا عن دراسة الفعالية البيولوجية للمركبات المحضرة. اثبتت القياسات المستخدمة ان الليكاند المحضر يتناسق مع ايون الفلز المركزي من خلال ذرة اوكسجين مجموعة الكاربونيل وذرة نتروجين مجموعة الازوميثين ويتخذ معقد النيكل المحضر الشكل الفراغي مربع مستوي اما بقية المعقدات فتتخذ الشكل الفراغي رباعي السطوح، كما تبين ان المعقدات المحضرة غير موصلة الكتروليتيا واثبتت المركبات المحضرة فعالية عالية كمضاد بكتريا لبعض أنواع بكتريا كرام الموجبة والسالبة.

**الكلمات المفتاحية:** معوضات 4-امينو انتي بايرين، الفعالية البيولوجية، ليكاند قاعدة شيف، معقدات المربع المستوي، الفلزات الانتقالية.