

Evaluation of Antibiofilm and Cytotoxic Activity of Microalgae Isolated from Different Sites of betwata- Erbil/Iraq

Sewgil Saaduldeen Anwer^{1,2*}  , Ayad Kareem Ali³  , Nur Bozbeyoglu Kart⁴  ,
Dogukan Mutlu⁵  , Refika Ceyda Beram⁵  , Volkan Kuzucu⁵  , Sevki Arslan⁵  ,
, Nazime Mercan Dogan⁵  

¹Clinical Biochemistry, College of Health Sciences, Hawler Medical University, Erbil, Iraq.

²Nursing department, Faculty of Nursing, Tishk International University, Erbil, Iraq.

³Community Health- Koya, Technical Institute, Erbil Polytechnic University, Erbil, Iraq.

⁴Plant and Animal Production Department, Tavas Vocational High School, Pamukkale University, Denizli, Türkiye.

⁵Biology Department, Faculty of Science, Pamukkale University, Denizli, Türkiye.

*Corresponding author.

Received 02/10/2023, Revised 11/12/2023, Accepted 13/12/2023, Published Online First 20/04/2024



© 2022 The Author(s). Published by College of Science for Women, University of Baghdad.

This is an Open Access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

This *in vivo* study is aimed to assess the antibiofilm and cytotoxic potential of selected microalgae isolates. Different concentration of microalgal isolates used for each of antibiofilm and cytotoxic studies. A total extract of 10 µL MTT was added to each well with LB broth, plates were read using microtiter plate spectrophotometer at 490 nm and 630 nm, respectively. The cell viability was determined in CCD-18°C cells (3×10^3 cells/well) by MTT assay at 590 nm. The microalgae isolates showed antibiofilm affect after 48 hrs on *Pseudomonas aeruginosa* PA01 and this effect was observed to be higher than those of *S. aureus*. *Chlorella* sp. which showed the maximum antibiofilm activity while *Arthrospira platensis* had no biofilm inhibition activity. *Chlorella* sp. inhibited about 82.67% of the biofilm at 5 mg/mL concentration. Moreover, the antibiofilm effects of *Chara* sp. and *Spyrogyra* sp.1 reached 79.01% and 76.31% at 1 mg/mL, respectively. Among all microalgal isolates, a moderate inhibitory effect was observed on *Pseudomonas aeruginosa* biofilm with *Chlorella sorokiniana*. Cell viability was not changed significantly at higher doses extracts. More than 73% of the cells were viable in all concentration of the algae extracts. All of the microalgal samples were found to have biofilm inhibition activity. The findings suggested that future development of microalgal samples as a means of inhibiting *Pseudomonas aeruginosa* biofilms is possible. Colonic epithelial cells may become cytotoxic in response to algae extracts. Further research is necessary to verify that extracts are safe for human consumption, which may have been hypothesized.

Keywords: Antibiofilm, Cell Line, Cytotoxicity, Cell viability, Microalgae.

Introduction

The occurrence of many biofilm-based human infections and their multiple antimicrobial resistance is a major concern in medicine and human health.

The treatment of infections caused by biofilms with antimicrobial drugs has prompted several studies since biofilms act a significant role in infectious

diseases. However, due to the development of multidrug-resistant bacteria, antibiotic therapy in infections brought on by foreign bodies is insufficient^{1,2}. Bacterial anti-biofilm actions are thus expected to manage bacterial populations in an ecological niche in a natural way³. Compounds with antibacterial action generated by cyanobacteria and microalgae have been reported in several studies. Nevertheless, there are little investigations on the antibiofilm activity of the extracts and/or chemicals these microbes create. Several studies have demonstrated that microalgae play an important role in the pharmaceutical business because of their potential to create secondary metabolites with a wide variety of pharmacological actions, such as anti-inflammatory, antioxidant, and antimicrobial properties, antibacterial, cytotoxic, antiviral, and anticoagulant. Antibacterial activity from microalgae extracts has been examined by several researchers due to the rising number of reported cases of antibiotic resistance⁴.

Microalgae are photosynthetic organisms that account for a large amount of freshwater and marine phytoplankton⁵. The rise of antibiotic-resistant bacteria necessitates the investigation of novel microalgal bioactive chemicals that are cost-effective, safe, and potent⁶. Many studies showed that natural products obtained from plant, algae, microorganisms etc. has great diversity of

pharmacological activities due to their secondary metabolites contents⁷. Antibody-based antitoxins are a treatment option for neutralizing toxins in people who have been exposed to them or who have been infected with a toxin-producing microbe. The antibiotic activity of microalgae and cyanobacteria has been attributed to compounds belonging to several chemical classes, such as alkaloids, aromatic compounds, acetogenins, cyclophanes and paracyclophanes, dicarboximides, indanes, indoles, pH, lactones, and lipids⁸.

Many bioactive molecules obtained from microalgae have cytotoxic effects on cancer cells. However, the fact that a substance has shown cytotoxicity to cancer cells in anticancer drug research is not an adequate result. For a substance to have drug potential, it is important that it exhibits selective cytotoxicity⁹.

Concisely, secondary metabolites from microalgae and cyanobacteria are highly regarded and are being extensively explored as replacements in biomedical research because to their potential biological and pharmacological qualities. Numerous scientific reports have been mentioned and are continuously being investigated in this regard. This in vivo study's objective is to assess the antibiofilm and cytotoxic potential of microalgae and cyanobacteria that were isolated from various locations between Erbil and Iraq.

Materials and Methods

Bacteria and Algal Strains:

Staphylococcus aureus ATCC 29213 and *Pseudomonas aeruginosa* PA01 were used in the present study. Both bacterial strains were sub-cultured in Nutrient Broth (NB) and incubated in shaker incubator at 37 °C, for 24 hrs. with agitation speed 150 rpm. Each culture was then streaked on a nutrient agar medium and stained with gram stain to confirm purity of strains.

Different algal strains obtained from betwata-Erbil transferred to College of Health Sciences, Microbiology lab; sub cultured on BG11 medium. Furthermore, each of algal colony was identified under microscope based on morphological forms and then transferred to BG11 liquid medium, after 14 days the fresh biomass was dried overnight at 50 °C

, and was used for further processing (antibiofilm and cytotoxic activities)¹⁰.

Determination of Biofilm Inhibition Effect

The quantity of each living bacteria *S. aureus* ATCC 29213 and *P. aeruginosa* PA01 in the biofilm was monitored by methoxynitrosulfophenyl-tetrazolium carboxanilide (XTT) assay¹¹⁻¹³. The microalgal isolates at different concentrations (0.125-5.0 mg/mL), and Lysogeny broth (LB) broth were added to a 96 well plate to a final volume of 100 µl and incubated at 37°C for 24-48 hrs. After that, the wells were washed 2 to 3 times with 100 µl Phosphate buffered saline (PBS) and 50 µl XTT reaction solutions. The plate was incubated at 37°C for 5 hrs., later the plates were read in a microtiter plate spectrophotometer at 490, 630 nm, respectively. The media sterility control without biofilm was

represented as zero indicates, whereas the presence of biofilm creation is revealed by a positive value.

Fluorescence Microscope Analysis

For microscopy analysis, the coverslips prepared determining the method described by^{14,15}. The mixture of SYBR Green and Propidium iodide (SYBR Green/PI; z ratio of 1: 3 in 100 μ L dH₂O) was used and the coverslips were analyzed by Fluorescence Microscope (100x magnification).

Cytotoxic Effect

Cell Culture

The human fibroblast cell line CCD-18Co (CVCL_2379) was obtained from ATCC (American Type Culture Collection). The cells were cultivated in Dulbecco's Modified Eagle Medium (DMEM) with 10% fetal bovine serum (FBS, Serox) and 1% penicillin/streptomycin mixture (100x, Capricorn) in 5% CO₂ in humidified air at 37°C in a 75cm² culture flasks (Nest). After the cells reached 90% confluence, the culture medium was replaced every 2–3 days¹⁶.

Viability Assay

The MTT reagent (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) was purchased and used by following the procedure of the manufacturer. The cell line CCD-18°C cells (3x10³ cells/well) were seated in 96-well plates and incubated for 24 hrs. Then, the medium was refreshed with that containing concentration of each extract. The final concentrations of each microalgal strains were 0, 125, 250, 500, 750, and 1000 μ g/mL. The experiment was carried out three times, with three replicates included for each treatment. The medium was taken out after 48 hrs, and each well received 10 mL of MTT solution (5 mg/mL) in 100 mL of media¹⁷. To dissolve the purple-colored formazan products, the solution was withdrawn after 4 hrs and 50 μ L of dimethyl sulfoxide (DMSO, Carlo Erba) was added. A microplate reader was used to measure absorbance at 590 nm after the plates had been incubated for 30 min at 37 °C. The percentage of cell viability was calculated by using following formula:

Results and Discussion

The observation of samples under microscope showed the following characters Table 1 and Fig. 1 Also, molecular analyze of two species confirmed

Percentage of cell viability = (OD value in the experiment group – OD value in the blank well)/(OD value in the control group – OD value in the blank well) \times 100%¹⁸.

Molecular Characterization of Cyanobacteria and Microalgae

For molecular identification, Thermo Scientific GeneJET Genomic DNA Purification Kit was used for DNA isolation of the isolates. The procedure was applied in the user manual of the relevant kit. The amount and purity of DNA obtained after DNA isolation was determined by spectrophotometric measurements in Thermo Scientific Nanodrop 2000 (USA).

CYA781R – CYA359F¹⁹ primers were used in the PCR studies and Amplification was done on the gene areas that were intended for species identification. PCR reactions were performed with Solis Biodyne (Estonia) FIREPol® DNA Polymerase Taq polymerase enzyme. PCR products were sequenced using the same primer pairs on the ABI 3730XL Sanger sequencing instrument (Applied Biosystems, Foster City, CA) with the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA). Sanger sequencing was performed by a commercial firm (BM LABOSIS, Ankara, Turkey). The obtained DNA sequences were first edited using BioEdit software and then compared with other sequences in the NCBI (National Center for Biotechnology Information) Gene Bank (NCBI database) using the nucleotide BLAST algorithm. For identification at the species level, the criterion of the sequence in question was more than 99% similar to the species in the gene bank.

Statistical Analysis


Vassar Stats, an online statistical software tool, was used to conduct the statistical analyses. The data was presented as means, with standard error of the means included (SEM). The student t-test used to compare the data and ($p \leq 0.05$) was chosen for statistical significance.

that one of the cultures belong to *Chlorella* genus and shows that *Chlorella sorokiniana* with similarity of 100% (Accession number: MK177540.1). Another

culture was determined to be *Arthrospira platensis* with a 100 % (Accession number: MT426015.1) similarity rate.

Table 1. Morphological structure of algae


Filament or unicellular	Akinet	Hormogonia	Heterocyst	Colour	Chloroplast	Cell shape	Microalgae strains
Filament (Trichomes cylindrical, sheath absent, coiled into a more or less regular spiral, the apex of Trichome usually not attenuated)	Absent	present	Absent	Colour is variable, blue-green	Chlorophyl 1- a	Spiral	<i>Chlorella</i>
Filament spiral with Short chains of cells (of up to 10 cells), known as hormogonia, are produced as a result of the trichome's fragmentation at the necridia.	Absent	present	Absent	Colour is variable, blue-green	Chlorophyl 1- a	Spiral	<i>Arthrospira</i>
Unicellular single-celled alga	Absent	Absent	Absent	Green algae	Chlorophyl 1-b	Spherical	<i>Chlorella</i>
Filament (Filaments long & unbranched, cells cylindrical short to very long)	Absent	Absent	Absent	Green algae	Chlorophyl 1-b	cells cylindrical 1 short to very long	<i>Spirogyra</i>
Filament (Filaments long & unbranched, cells cylindrical short to very long)	Absent	Absent	Absent	Green algae	Chlorophyl 1-b	cells cylindrical 1 short to very long	<i>Spirogyra</i>
Filamentous algae	Absent	Absent	Absent	Light green	Discoid shape	The axis branches globule seen above the nuclei	<i>Chara</i>



Chlorella sp.



Arthrospira platensis



Chlorella soro kiniana



Figure 1. Morphological structures algal strains¹⁰

The inhibition of metabolic activity was detected by XTT Assay and the percentage of metabolic inhibition was evaluated as biofilm inhibition effect. The antibiofilm activity of algal samples against both pathogenic bacteria appeared to be dosage related. The results of antibiofilm activity of aqueous extracts were given as mean \pm standard deviation (SD) for three independent experiments. Statistically there was significant difference compared to the control ($*p \leq 0.05$).

In the first 24 hrs. of incubation, the samples did not show any effect on gram negative *P. aeruginosa*. The samples showed an effect on *P. aeruginosa* 48 hrs. In other words, the microalga and *Cyanobacteria* samples showed antibiofilm affect after 48 hrs on *P. aeruginosa* PA01 and this effect was observed to be higher than those of *S. aureus*. Against *P. aeruginosa*

biofilms, *Chlorella* sp. showed the maximum antibiofilm activity while *Arthrospira platensis* had no biofilm inhibition activity. *Chlorella* sp. inhibited about 82.67% of the biofilm at 5 mg/mL concentration. Moreover, the antibiofilm effects of *Chara* sp. and *Spyrogyra* sp.1 reached 79.01% and 76.31% at 1 mg/mL, respectively. Among all microalgal samples, the moderate inhibitory effect was observed on *P. aeruginosa* biofilm with *Chlorella sorokiniana*. In general, it was observed that *S. aureus* were more resistant than *P. aeruginosa* against all samples, Fig. 2. Among the algal samples, only *Chara* sp., *Chlorella* sp. and *Chlorella sorokiniana* samples were found to have a significant biofilm inhibition. This difference in their antibiofilm activities may be related to both differences of microalgal and *Cyanobacteria* samples and the difference in bacterial behavior.




Figure 2. Antibiofilm activity of extracts. a. *Chlorella sorokiniana*, b. *Arthrospira platensis*, c. *Spyrogyra* sp.1, d. *Spyrogyra* sp.2, e. *Chara* sp., f. *Chlorella* sp.

We now face a huge public health issue since the majority of traditional antimicrobial drugs are no longer effective at this crucial moment when infections have developed diverse resistance routes that have allowed them to surpass our capacity to effectively manage them. In this context, the quest for natural remedies with unique methods to prevent and/or treat life-threatening illnesses might be inspired by the aquatic environment, a place of great biodiversity.

The findings of this investigation revealed that six unprocessed algal extracts exhibit antibacterial properties against *P. aeruginosa* and *S. aureus*. According to the findings, all algal extracts had a substantial amount of antibiofilm activity. High antibiofilm activity showed by *Chlorella* against *P. aeruginosa* within 48 hrs while *Arthrospira platensis* showed low antibiofilm activity. The researchers in ²⁰ used different solvents for antibiofilm activity and they concluded that each of solvents showed different antibiofilm activity this discrepancy in the results is likely due to a number of variables that affected the antimicrobial effectiveness of marine algae, including the season and place of algal species, various phases of plant development, extraction ability.

The acetone extract of *Pleurastrum minutum* demonstrated the highest antimicrobial activity among the tested extracts and showed moderate

antioxidant activity. Conversely, the methanol extract of the same microalgae exhibited the highest antioxidant activity and moderate antimicrobial activity. These findings underscore the significance of the choice of solvent in the extraction process, as it plays a crucial role in selectively extracting specific bioactive compounds with diverse properties ²¹.

Anwer and coauthor's isolated phenolic compound and fatty acid properties of some microalgae species and they found that microalgae with a high antioxidant capacity may be investigated for various industrial and pharmaceutical uses ²².

It is known that cell-to-cell intercellular interaction is an important for biofilm formation and structure. For an effective intercellular communication, bacterial cells need to be positioned in close proximity with their neighbors. After the treated with micro-algal and *Cyanobacteria* samples, the bacterial biofilm structure imaged and the close interaction of cells were degraded. The photograph of microscopy images of control group (untreated with samples) and cells treated with micro-algal samples were given in Fig. 3. The general biofilm structure after the treated with algal samples was not a continuous and no dense EPS-matrix. Moreover, the micro-colonies of cell aggregates encapsulated in EPS-matrix and the water channels were degraded.




Figure 3. General Biofilm structure before and after the treated with algal samples

To evaluate the cell viability, all extracts were tested by using the MTT assay method. The results for the percentage of cell viability after treatment for 48 hrs., are shown in Fig. 4. As can be seen in Fig. 4, the cell viability was not changed at higher doses and more than 73% of the cells was viable in all concentration of the algae extracts. These results also clearly showed that all algae extracts cannot showed cytotoxic effect on colon cells. It may have

speculated that extracts are safe for human usage and further tests will be required to test this hypothesis.




Figure 4. Effects of samples on the CCD-18Co cells after 48 h of treatment. Results were given as mean \pm standard deviation (SD) for three independent experiments. Statistically, there was significant difference compared to the control (* $p \leq 0.05$).

CCD-18Co is well-known human fibroblast cell line isolated from normal colon tissue. It is widely used in experimental pharmacology and cancer studies. Moreover, it is important to describe whether secondary metabolites have toxicological effects to gastrointestinal system or not²³⁻²⁵. Thus, CCD-18Co cell line was used throughout in this study.

The results indicated a minimal decline in the viability even in higher doses of algae extracts after 48 hrs. incubation. These results also clearly showed that all algae extracts cannot showed cytotoxic effect on colon cells. Therefore, it may be speculated that they are safe to use many purposes including antioxidant source. Algae have been shown to contain large amounts of cytotoxic substances with

Conclusion

Antibiotic resistance spreads among bacteria in the world and biofilm formation is effective in spreading this resistance. All algal strains showed antibiofilm activity. *Chlorella* sp. showed maximum antibiofilm activity. It was observed that all microalgal samples had biofilm inhibition activity. And these findings were also confirmed by fluorescent microscopy analysis. Bacterial biofilms will cause a lot of damage in the future, both in the health sector and in

anticancer, antitumor, antiproliferative, and antimetastatic characteristics such fucoidans, laminarians, and terpenoids^{26, 27}. In a study done by^{28, 29}, it was found that the viability of cells decreased as extract concentrations increase. Bechelli and coauthors³⁰ implied that algae extracts may be able to suppress primary leukemia blasts and AML cell lines, as well as normal hematopoietic cells. As demonstrated by³¹, the methanol extract showed more cytotoxicity oxidative stress than the chloroform extract in the MCF-7 cell line and the tested algae may have potential use for treatment of cancer. These results showed that algae extracts may have the potential to be cytotoxic to cancer cells including gastrointestinal cancers without affecting normal cells. But, further experiments will be required to verify this condition.

Due to its quick growth, ease of large-scale cultivation compared to other microalgae, and valuable nutritional contents, chlorella is one of the microalgae that has been investigated extensively. According to research by³², oral administration of *Chlorella pyrenoidosa* effectively and dose-dependently prevented increases in blood aspartate aminotransferase (AST), alanine aminotransferase (ALT), and glutamate aminotransferase (GAT) and alkaline phosphatase (ALP) activities in male rats with d-galactosamine-induced hepatic lesions. It also appeared to alleviate hepatic cell necrosis. Additionally, they discovered that *C. pyrenoidosa* reduced lipid peroxide levels and restored normal triglyceride levels in rat serum. When compared to earlier research, the current study revealed limited impacts on cell viability; this could be because the current study utilized a low concentration^{33,34}.

the economic field. In order to prevent this, it is important to find new antibiofilm substances. Although the importance of microalgae among new antibiofilm products is increasing, it is necessary to investigate their mechanism of action in detail. Also, the results indicated that microalgal samples have the potential to be developed as an agent to inhibition of *P. aeruginosa* biofilms in the future.

Acknowledgment

The study was carried out at Pamukkale University, Denizli- Turkey and Hawler Medical University, Erbil-Iraq.

Authors' Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and Images, that are not ours, have been included with the necessary permission for re-publication, which is attached to the manuscript.
- No animal studies are present in the manuscript.
- No human studies are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee in Hawler Medical University

Authors' Contribution Statement

S.S.A. and A. A. contributed in Isolation and identification of Microalgae, R. C. B. was responsible for molecular identification, S. A. and D. M. analyzed cytotoxic activity of samples, N. M. D., N. B. K. and made antibiofilm and V. K. used

fluorescence microscopy for analysis. All authors have substantial contributions to the final manuscript and approved this submission. All authors read the manuscript carefully and approve the final version of their MS.

References

1. Schulze A, Mitterer F, Pombo J P, Schild S. Biofilms by bacterial human pathogens: Clinical relevance - development, composition and regulation - therapeutical strategies. *Microb Cell*. 2021 Feb; 8(2): 28–56. <https://doi.org/10.15698/mic2021.02.741>
2. Prestinaci F, Pezzotti P, Pantosti A. Antimicrobial resistance: A global multifaceted phenomenon. *Pathog Glob Health*. 2015 Sep; 109(7): 309–318. <https://doi.org/10.1179/2047773215Y.0000000030>
3. Schoenborn AA, Yannarell SM, Wallace ED, Clapper H, Weinstein I C, Shank EA. Defining the expression, production, and signaling roles of specialized metabolites during bacillus subtilis differentiation. *J Bacteriol*. 2021 Oct; 203(22): e0033721. <https://doi.org/10.1128/JB.00337-21>
4. Karunakaran E, Biggs CA. Mechanisms of bacillus cereus biofilm formation. An investigation of the physicochemical characteristics of cell surfaces and extracellular proteins. *Appl. Microbiol Biotechnol*. 2011 Feb; 89(4): 1161–1175. <https://doi.org/10.1007/s00253-010-2919-2>
5. Shaima AF, Yasin NH, Ibrahim NH, Takrif MS, Gunasekaran DI, Mahmud YY. Unveiling antimicrobial activity of microalgae *Chlorella sorokiniana* (UKM2), *Chlorella* sp. (UKM8) and *Scenedesmus* sp. (UKM9). *Saudi J Biol Sci*. 2022 Oct; 29(2): 1043–1052. <https://doi.org/10.1016/j.sjbs.2021.09.069>
6. Lauritano C, Andersen J, Hansen E, Albrigtsen ME, Esposito F, Khanssen K, et al. Bioactivity screening of microalgae for antioxidant, anti-inflammatory, anticancer, anti-diabetes, and antibacterial activities. *Front Mar Sci*. 2016 May; 3(68): 1–2. <https://doi.org/10.3389/fmars.00068>
7. Barrera DJ, Rosenberg JN, Chiu JG, Chang YN, Debatis M, Ngoi SM, et al. Algal chloroplast produced camelid VH antitoxins are capable of neutralizing botulinum neurotoxin. *Plant Biotechnol J*. 2015 Jan; 13(1): 117-124. <https://doi.org/10.1111/pbi.12244>
8. Tian F, Li J, Nazir A, Tong Y. Bacteriophage - a promising alternative measure for bacterial biofilm control. *Infect Drug Resist*. 2021 Jan; 20(14): 205–217. <https://doi.org/10.2147/IDR.S290093>
9. Arslan A, Yılancıoğlu K, Kuşoğlu S, Albayrak G. Chemical constituent of Isochrysis galbana microalgae extract and its cytotoxic activities on leukemic cell lines. *Istanbul J Pharm*. 2022 Apr; 52 (1): 64-68. <https://doi.org/10.26650/IstanbulJPharm.2022.105733>
10. Anwer S. Direct analysis of algae for determination of major and minor elements using energy dispersive x-ray fluorescence (EDXRF), *Kuwait Journal of Science*, 2023 Sep; 50 (4): 697-702, <https://doi.org/10.1016/j.kjs.2023.04.012>.
11. Shen F, Ge C, Yuan P. Metabolomics Study Reveals Inhibition and Metabolic Dysregulation in Staphylococcus aureus Planktonic Cells and Biofilms Induced by Carnosol. *Front Microbiol*. 2020 Sep; 18: 538572. <https://doi.org/10.3389/fmicb.2020.538572>
12. Sivaranjani M, Gowrishankar S, Kamaladevi A, Pandian SK, Balamurugan K, Ravi AV. Morin

- inhibits biofilm production and reduces the virulence of *Listeria monocytogenes* an in vitro and in vivo approach. *Int J Food Microbiol.* 2016 Nov; 21: 237: 73–82.
<https://doi.org/10.1016/j.ijfoodmicro.2016.08.021>
13. Feng J, Yee R, Zhang S, Tian L, Shi W, Zhang WH, et al. A Rapid Growth-Independent Antibiotic Resistance Detection Test by SYBR Green/Propidium Iodide Viability Assay. *Front Med.* 2018 May; 5(127): 1-11.
<https://doi.org/10.3389/fmed.2018.00127>
14. Haney EF, Trimble MJ, Cheng JT, Vall Q, Hancock RE W. Critical assessment of methods to quantify biofilm growth and evaluate antibiofilm activity of host defence peptides. *Biomolecules.* 2018 May; 8(2):29. <https://doi.org/10.3390/biom8020029>
15. Sahin C, Mutlu D, Nasirli F, Mahmoudi G, Zubkov FI, Arslan S, et al. New iridium bis-terpyridine complexes: synthesis, characterization, antibiofilm and anticancer potentials. *Biometals.* 2021 Apr; 34: 701–713. <https://doi.org/10.1007/s10534-021-00307-y>
16. David S, Perkins RS, Fronczek FR, Kasiri S, Mandal SS, Srivastava RS. Synthesis, characterization, and anticancer activity of ruthenium-pyrazole complexes. *J Inorg Biochem.* 2012 Jun; 111:33-9.
<http://doi.org/10.1016/j.jinorgbio.2012.02.022>
17. Nedji N, Loucif-Ayad W. Antimicrobial activity of Algerian propolis foodborne pathogens and its quantitative chemical composition. *Asian Pacific Journal of Tropical Disease,* 2014 4(6): 433–437.
[http://doi.org/10.1016/S2222-1808\(14\)60601-0](http://doi.org/10.1016/S2222-1808(14)60601-0)
18. Wu Z, McGoogan JM. 2020. Characteristics of and important lessons from the Coronavirus Disease 2019 (COVID-19) outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *Jama-Journal of the American Medical Association* 2020 Apr 323(13): 1239–1242.
<https://doi.org/10.1001/jama.2020.2648>
19. Nübel U, Garcia-Pichel F, Muyzer G. PCR primers to amplify 16S rRNA genes from cyanobacteria. *Appl. Environ. Microbiol.* 1997 Aug; 63(8): 3327-3332.
<https://doi.org/10.1128/aem.63.8.3327-3332.1997>
20. Guchhait KC, Manna T, Barai M, Karmakar M, Nandi SK, Jana D, et al. Antibiofilm and anticancer activities of unripe and ripe *Azadirachta indica* (neem) seed extracts. *BMC Complement Med Ther.* 2022 Feb; 22(42): 1-18. <https://doi.org/10.1186/s12906-022-03513-4>
21. Elshobary E, El-Shenody A, Ashour M, Zabed M, Qi X. Antimicrobial and antioxidant characterization of bioactive components from *Chlorococcum minutum*. *Food Biosci.* 2020 Jan; 35: 100567.
<https://doi.org/10.1016/j.fbio.2020.100567>
22. Anwer SS, Sadiq KH, Muhammad KR, Aladdin LM. Phenolic compound and fatty acid properties of some microalgae species isolated from Erbil City. *Braz J Biol.* 2022 Jan; 21; 82: e256927.
<https://doi.org/10.1590/1519-6984.256927>
23. Gadhi A, El-Sherbiny M, Al-Sofyani A, Ba-Akdah M, Satheesh S. Antibiofilm Activities of Extracts Of The Macroalga *Halimeda* Sp. From The Red Sea. *J Mar Sci Technol.* 2018; 26(6): 838-846.
[https://doi.org/10.6119/JMST.201812_26\(6\).0008](https://doi.org/10.6119/JMST.201812_26(6).0008)
24. Ibraheem S, Sadiq AS, Al-Tameemli M, Alias MF. Synthesis, Spectral Identification, Antibacterial Evaluation and Theoretical Study of Co, Fe, Rh and Pd Complexes for 2-benzoylthiobenzimidazol. *Baghdad Sci J.* 2022 Dec; 19(6): 1326-1334.
<https://dx.doi.org/10.21123/bsj.2022.6704>
25. Griffin C, Karnik A, McNulty J, Pandey S. Pancratistatin selectively targets cancer cell mitochondria and reduces growth of human colon tumor xenografts. *Mol Cancer Ther.* 2011 Jan; 10(1): 57-68. <https://doi.org/10.1158/1535-7163.MCT-10-0735>. PMID: 21220492.
26. Toma JJ, Aziz F. Antibacterial Activity of Three Algal Genera against some Pathogenic Bacteria. *Baghdad Sci J.* 2023 Feb; 20(1): 32-40.
<https://dx.doi.org/10.21123/bsj.20022.6818>
27. Takawa M, Masuda K, Kunizaki M, Daigo Y, Takagi K, Iwai Y, et.al. Validation of the histone methyltransferase EZH2 as a therapeutic target for various types of human cancer and as a prognostic marker. *Cancer Sci.* 2011; 102(7): 1298-1305.
<https://doi.org/10.1111/j.1349-7006.2011.01958.x>
28. Mendanha, D.; Vieira de Castro, J.; Moreira, J.; Costa, B.M.; Cidade, H.; Pinto, M.; Ferreira, H.; Neves, N.M. A New Chalcone Derivative with Promising Antiproliferative and Anti-Invasion Activities in Glioblastoma Cells. *Molecules* 2021, 26, 3383.
<https://doi.org/10.3390/molecules26113383>
29. Zong A, Cao H, Wang F. Anticancer polysaccharides from natural resources: a review of recent research. *Carbohydr Polym.* 2012 Nov; 90(4): 1395–1410.
<https://doi.org/10.1016/j.carbpol.2012.07.026>
30. Mosaddegh M, Gharanjik BM, Naghibi F, Esmaeili S, Pirani A, Eslami Tehrani B, et al. A survey of cytotoxic effects of some marine algae in the Chabahar coast of Oman Sea. *Res j pharmacogn.* 2014; 1: 27–31.
31. Bechelli J, Coppage M, Rosell K, Liesveld J. Cytotoxicity of algae extracts on normal and malignant cells. *Leuk Res Treatment.* 2011 Jan; 2011: 373519.
<https://doi.org/10.4061/2011/373519>
32. Kurt O, Özdal-Kurt F, Akçora CM, Özkut M, Tuğlu, MI. Neurotoxic, cytotoxic, apoptotic and antiproliferative effects of some marine algae. Extracts on the NA2B cell line. *Biotech. Histochem.* 2018 January; 93(1): 59–69.
<https://doi.org/10.1080/10520295.2017.1381992>
33. El-Sheekh M, Abu-Faddan M, Abo-Shady A, Nassar MZA, Labib W. Molecular identification, biomass, and biochemical composition of the marine Chlorophyte *Chlorella* sp. MF1 isolated from Suez Bay. *J Genet Eng Biotechnol.* 2020 Jul; 18(1): 27.
<https://doi.org/10.1186/s43141-020-00044-8>

34. Yarmohammadi S, Hosseini R, Foshati S, Moradi M, Hemati N, Moradi S, et al. Effect of *Chlorella vulgaris* on Liver Function Biomarkers: a Systematic Review

and Meta-Analysis. Clin Nutr Res. 2021 Jan; 10(1):83-94. <https://doi.org/10.7762/cnr.2021.10.1.83>

تقييم التضاد الحيوي والنشاط السمي للخلايا من قبل الطحالب الدقيقة التي تم عزلها من مواقع مختلفة في بيتواتا- أربيل/العراق

سيوكل سعدالدين انور^{1,2}، اياد كريم علي³، نور بوزيبي كورت⁴، دوغوكان موتلو⁵، رفيقة جيدا بيرام⁵، فولكان كوزوجوا⁵، شوقي ارسلان⁵، نزيمة مرجان دوغان⁵

¹قسم الكيمياء السريرية، كلية العلوم الصحية، جامعة هتولير الطبية، أربيل، العراق.

²قسم التمريض، كلية التمريض، جامعة تيشك الدولية، أربيل، العراق.

³قسم صحة المجتمع، كويبا، المعهد التقني، جامعة أربيل التقنية، أربيل، العراق.

⁴معهد تافاس الثانوية المهنية، جامعة باموكالي، دنيزلي، تركيا.

⁵بأبولوجي، كلية العلوم، جامعة باموكالي، دنيزلي، تركيا.

الخلاصة

تمتلك الطحالب الدقيقة نظراً لسهولة زراعتها ونموها السريع وتجدها، امكانات كمصدر للمركبات النشطة بيولوجياً للاغراض الصناعية والطبية. تهدف هذه الدراسة التي اجريت بشكل (in vivo) الى تقييم القدرة المضادة للاغشية الحيوية والسموم الحيوية لطحالب الدقيقة مختارة وعزلات البكتريا الزرقاء. تم اختيار تراكيز مختلفة من عزلات الطحالب الدقيقة في كل من دراسات المضادات الحيوية والخلاية السامة. تمت اضافة مستخلص اجمالي قدره 10 ميكروليتر من MTT الى كل بئر باستخدام مرق LB, وتمت قراءة اللوحات باستخدام مقياس الطيف الضوئي للوحة ميكروتيتر عند 490 نانوميتر و 630 نانوميتر. على التوالي. تم تحديد بقاء قدرة الخلية في خلايا-CCD 18C° (3x10³ cells/well) بواسطة اختبار MTT عند 590 نانوميتر. اظهرت عزلات الطحالب الدقيقة تأثير مضاد حيوي بعد 48 ساعة على *Pseudomonas aeruginosa* PA01 ولوحظ ان هذا التأثير اعلى من تأثيره على *Chlorella sp.* *S. aureus*. اظهر الحد الاقصى لنشاط المضاد الحيوي بينما لم يكن لدى *Arthrospira platensis* اي نشاط تثبيط للاغشية الحيوية. *Chlorella sp.* تثبط حوالي 82.67% من الغشاء الحيوي في تركيز 5 ملغم/مل. علاوة على ذلك فان التأثيرات المضادة للاغشية الحيوية ل (*Spyrogyra sp.* و *Chara sp.*) وصلت بنسبة الى 79.01% و 76.31% عند 1 ملغم/مل على التوالي. من بين جميع العزلات الطحالب الدقيقة، لوحظ وجود تأثير مثبط معتدل على الاغشية الحيوية ل *Pseudomonas aeruginosa* مع *Chlorella sorokiniana*. لم تتغير قابلية الخلية للبقاء بشكل ملحوظ في مستخلصات جرعات عالية. اكثر من 73% من الخلايا كانت قابلة للحياة في جميع تركيزات مستخلصات الطحالب. وجد بان جميع عينات الطحالب الدقيقة لها نشاط تثبيط الاغشية الحيوية. تشير النتائج الى ان التطوير المستقبلي لعينات الطحالب الدقيقة كوسيلة لتثبيط الاغشية الحيوية ل *Pseudomonas aeruginosa* امر ممكن. قد تصبح الخلايا الطلائية القولونية خلية سامة استجابة لمستخلصات الطحالب. من الضروري اجراء المزيد من الابحاث للتحقق من ان المستخلصات آمنة للاستهلاك البشري، وهو ما يكون مفترضاً.

الكلمات المفتاحية: المضادات الحيوية، الخط الخلوي، السمية الخلوية، حيوية الخلية، الطحالب الدقيقة.